

8-28-2009

Contact Process in a Wedge

J. Theodore Cox

Syracuse University, jtcox@syr.edu

Nevena Maric

University of Missouri - St Louis

Rinaldo Schinazi

University of Colorado at Colorado Springs

Follow this and additional works at: <http://surface.syr.edu/mat>

 Part of the [Mathematics Commons](#)

Recommended Citation

Cox, J. Theodore; Maric, Nevena; and Schinazi, Rinaldo, "Contact Process in a Wedge" (2009). *Mathematics Faculty Scholarship*. Paper 42.

<http://surface.syr.edu/mat/42>

This Article is brought to you for free and open access by the Department of Mathematics at SURFACE. It has been accepted for inclusion in Mathematics Faculty Scholarship by an authorized administrator of SURFACE. For more information, please contact surface@syr.edu.

Contact process in a wedge

J. Theodore Cox* Nevena Marić†
Syracuse University *University of Missouri - St. Louis*

Rinaldo Schinazi‡
University of Colorado - Colorado Springs

August 28, 2009

Abstract

We prove that the supercritical one-dimensional contact process survives in certain wedge-like space-time regions, and that when it survives it couples with the unrestricted contact process started from its upper invariant measure. As an application we show that a type of weak coexistence is possible in the nearest-neighbor “grass-bushes-trees” successional model introduced in [3].

Key words: contact process, grass-bushes-trees

AMS Classification: Primary: 60K35; Secondary: 82B43

1 Introduction

The contact process of Harris (introduced in [5]) is a well known model of infection spread by contact. The one-dimensional model is a continuous time Markov process ξ_t on $\{0, 1\}^{\mathbb{Z}}$. For $x \in \mathbb{Z}$, $\xi_t(x) = 1$ means the individual at site x is infected at time t while $\xi_t(x) = 0$ means the individual is healthy. Infected individuals recover from their infection after an exponential time with mean 1, independently of everything else. Healthy individuals become infected at a rate proportional to the number of infected neighbors. Alternatively, individuals (1's) die at rate one and give birth onto neighboring empty sites (0's) at rate λ . If we let $n_i(x, \xi) = \sum_{y:|y-x|=1} 1\{\xi(y) = i\}$, and $\lambda \geq 0$ the infection parameter, then the transitions at x in state ξ are

$$1 \rightarrow 0 \text{ at rate } 1 \quad \text{and} \quad 0 \rightarrow 1 \text{ at rate } \lambda n_1(x, \xi). \quad (1)$$

When convenient we will identify $\xi \in \{0, 1\}^{\mathbb{Z}}$ with $\{x : \xi(x) = 1\}$, and use the notation $\|\xi\|_i = \sum_x 1\{\xi(x) = i\}$.

*Supported in part by NSF Grant No. 0803517

†Supported in part by NSF Grant No. 0803517

‡Supported in part by NSF Grant No. 0701396

Let ξ_t^0 denote the contact process with initial state $\xi_0^0 = \{0\}$. The critical value λ_c is defined by

$$\lambda_c = \inf\{\lambda \geq 0 : P(\xi_t^0 \neq \emptyset \text{ for all } t \geq 0) > 0\}. \quad (2)$$

It is well known that $0 < \lambda_c < \infty$, and that in the supercritical case $\lambda > \lambda_c$ there is a unique stationary distribution ν for ξ_t , called the upper invariant measure, with the property

$$\nu(\xi : \|\xi\|_1 = \infty) = 1.$$

There are also well-defined ‘‘edge speeds.’’ Let ξ_0^- (ξ_0^+) be the initial state given by $\xi_0^- = \mathbb{Z}^-$ ($\xi_0^+ = \mathbb{Z}^+$), and define the edge processes

$$r_t = \max\{x : \xi_t^-(x) = 1\} \text{ and } l_t = \min\{x : \xi_t^+(x) = 1\}. \quad (3)$$

There is a strictly increasing function $\alpha : (\lambda_c, \infty) \rightarrow (0, \infty)$ such that for $\lambda > \lambda_c$

$$\lim_{t \rightarrow \infty} \frac{r_t}{t} = \alpha(\lambda) \text{ and } \lim_{t \rightarrow \infty} \frac{l_t}{t} = -\alpha(\lambda) \text{ a.s.} \quad (4)$$

All of the above facts are contained in Chapter VI of [6] and Part I of [7].

We are interested in contact processes for which the infection is restricted to certain space-time regions. For $\mathcal{W} \subset \mathbb{Z} \times [0, \infty)$ define the \mathcal{W} -restricted contact process $\xi_t^{\mathcal{W}}$ as follows. First, set $\xi_t^{\mathcal{W}}(x) = 0$ for all $(x, t) \notin \mathcal{W}$. Second, for $(x, t) \in \mathcal{W}$, replace (1) with

$$1 \rightarrow 0 \text{ at rate } 1 \quad \text{and} \quad 0 \rightarrow 1 \text{ at rate } \lambda \sum_{y:|y-x|=1} \xi(y)1_{\mathcal{W}}(y, t), \quad (5)$$

so that infection spreads only between sites in the wedge. We will give an explicit *graphical construction* of $\xi_t^{\mathcal{W}}$ in Section 2.

For $0 < \alpha_l < \alpha_r < \infty$ and $M \geq 0$ define the ‘‘wedges’’ $\mathcal{W} = \mathcal{W}(\alpha_l, \alpha_r, M)$ by

$$\mathcal{W} = \{(x, t) \in \mathbb{Z} \times [0, \infty) : \alpha_l t \leq x \leq M + \alpha_r t\}. \quad (6)$$

In view of (4), we will impose the conditions

$$\lambda > \lambda_c \text{ and } 0 < \alpha_l < \alpha_r < \alpha(\lambda). \quad (7)$$

Our first result is that survival in wedges is possible.

Theorem 1. *Assume (7) holds, $\mathcal{W} = \mathcal{W}(\alpha_l, \alpha_r, M)$, and $\xi_0^{\mathcal{W}} = [0, M] \cap \mathbb{Z}$. Then*

$$\lim_{M \rightarrow \infty} P(\xi_t^{\mathcal{W}} \neq \emptyset \text{ for all } t \geq 0) = 1. \quad (8)$$

When $\xi_t^{\mathcal{W}}$ survives it looks like the unrestricted contact process in equilibrium. To state this more precisely, let

$$r_t^{\mathcal{W}} = \max\{x : \xi_t^{\mathcal{W}}(x) = 1\} \text{ and } l_t^{\mathcal{W}} = \min\{x : \xi_t^{\mathcal{W}}(x) = 1\}, \quad (9)$$

and let ξ_t^{ν} denote the contact process started in its upper invariant measure ν .

Theorem 2. Assume (7), $\mathcal{W} = \mathcal{W}(\alpha_l, \alpha_r, M)$, and $\xi_0^{\mathcal{W}} = [0, M] \cap \mathbb{Z}$. On the event $\{\xi_t^{\mathcal{W}} \neq \emptyset \text{ for all } t \geq 0\}$,

$$\lim_{t \rightarrow \infty} \frac{r_t^{\mathcal{W}}}{t} = \alpha_r \text{ and } \lim_{t \rightarrow \infty} \frac{l_t^{\mathcal{W}}}{t} = \alpha_l \text{ a.s.} \quad (10)$$

Furthermore, $\xi_t^{\mathcal{W}}$ and ξ_t^{ν} can be coupled so that on the event $\{\xi_t^{\mathcal{W}} \neq \emptyset \text{ for all } t \geq 0\}$,

$$\xi_t^{\mathcal{W}}(x) = \xi_t^{\nu}(x) \text{ for all } x \in [l_t^{\mathcal{W}}, r_t^{\mathcal{W}}] \text{ for all large } t \text{ a.s.} \quad (11)$$

Remark 3. By standard arguments using exponential estimates, $|\xi_t^{\nu} \cap [at, bt]| \rightarrow \infty$ as $t \rightarrow \infty$ with probability one for any $a < b$ (see Theorem VI.3.33 in [6]). Therefore Theorem 2 implies that when $\xi_t^{\mathcal{W}}$ survives, $|\xi_t^{\mathcal{W}}| \rightarrow \infty$ a.s.

Theorem 1 can be used to obtain information about the “grass-bushes-trees” model (GBT) of [3]. In this model sites are either empty (0), occupied by a bush (1) or occupied by a tree (2). Both 1’s and 2’s turn to 0’s at rate one. The 2’s give birth at rate λ_2 on top of 1’s and 0’s. The 1’s give birth at rate λ_1 on top of 0’s only, and hence are at a disadvantage compared to 2’s. The state space for the process is $\{0, 1, 2\}^{\mathbb{Z}}$, and the nearest-neighbor version of the model makes transitions at x in state ζ

$$0 \rightarrow \begin{cases} 1 & \text{at rate } \lambda_1 n_1(x, \zeta) \\ 2 & \text{at rate } \lambda_2 n_2(x, \zeta) \end{cases} \quad 1 \rightarrow \begin{cases} 0 & \text{at rate 1} \\ 2 & \text{at rate } \lambda_2 n_2(x, \zeta) \end{cases} \quad 2 \rightarrow 0 \text{ at rate 1.} \quad (12)$$

A natural question to ask is whether or not coexistence of 1’s and 2’s is possible. It was shown in [3] that coexistence is possible for a non-nearest neighbor version of the model and appropriate λ_i , where coexistence meant that ζ_t had a stationary distribution μ such that

$$\mu\left(\zeta : \|\zeta\|_i = \infty \text{ for } i = 1, 2\right) = 1. \quad (13)$$

It was also shown in [3] that there is no stationary distribution satisfying (13) in the nearest-neighbor case for *any* choice of the λ_i . Moreover, if there are infinitely many 2’s initially then for each site there is a last time at which a 1 can be present. Nevertheless, it is a consequence of Theorem 1 and the construction used in its proof that a form of weak coexistence is possible, even starting from a single 1 and infinitely many 2’s.

Corollary 4. Let ζ_t be the GBT process with initial state ζ_0 , where $\zeta_0(x) = 2$ for $x < 0$, $\zeta_0(0) = 1$ and $\zeta_0(x) = 0$ for $x > 0$. For all $\lambda_c < \lambda_2 < \lambda_1$,

$$P\left(\lim_{t \rightarrow \infty} \|\zeta_t\|_1 = \infty\right) > 0. \quad (14)$$

The 2’s spread to the right at rate $\alpha(\lambda_2)$, ignoring the 1’s, while the 1’s try to spread to the right at the faster rate $\alpha(\lambda_1)$. The 1’s will be killed by 2’s invading from the left, but Theorem 1 shows that they can survive with positive probability by moving off to the right in the space-time region free of 2’s.

Remark 5. (1) *With a little more work one can use Theorem 2 to say more about the set of 1's in ζ_t since it dominates wedge-restricted contact processes with positive probability.* (2) *Non-oriented percolation in various subsets of \mathbb{Z}^d has been studied by others (e.g. see [4] and [1]), but as far as we are aware our results on oriented percolation are new.*

In Section 2 we give the standard graphical construction due to Harris, then prove Theorem 1 in Section 3, Theorem 2 in Section 4, and Corollary 4 in Section 5.

2 The graphical representation

For $x \in \mathbb{Z}$ let $\{T_n^x : n \geq 1\}$ be the arrival times of a Poisson process with rate 1, and for all pairs of nearest-neighbor sites x, y let $\{B_n^{x,y} : n \geq 1\}$ be the arrival times of a Poisson process with rate λ . The Poisson processes $T^x, B^{x,y}$, $x, y \in \mathbb{Z}$, are all independent. At the times T_n^x we put a δ at site x to indicate a death at x , and at the times $B_n^{x,y}$ we draw an arrow from x to y , indicating that a 1 at x will give birth to a 1 at y . For $0 \leq s < t$ and sites x, y we say that there is an active path up from (x, s) to (y, t) if there is a sequence of times $t_0 = s \leq t_1 < t_2 < \dots < t_n \leq t_{n+1} = t$ and a sequence of sites $x_0 = x, x_1, \dots, x_n = y$ such that

1. for $i = 1, 2, \dots, n$, $|x_i - x_{i-1}| = 1$ and there is an arrow from x_{i-1} to x_i at time t_i
2. for $i = 0, \dots, n$, the time segments $\{x_i\} \times [t_i, t_{i+1}]$ do not contain any δ 's

By default there is always an active path up from (y, t) to (y, t) . For a space-time region $\mathcal{W} \subset \mathbb{Z} \times [0, \infty)$ we define $\xi_t^{\mathcal{W}}$, the contact process restricted to \mathcal{W} , as follows. Given an initial state $\xi_0 \subset \{x : (x, 0) \in \mathcal{W}\}$, set $\xi_t(y) = 0$ for all $(y, t) \notin \mathcal{W}$. If there is a site x with $\xi_0(x) = 1$ and an active path up from $(x, 0)$ to (y, t) lying entirely in \mathcal{W} set $\xi_t^{\mathcal{W}}(y) = 1$, otherwise set $\xi_t^{\mathcal{W}}(y) = 0$. For $\mathcal{W} = \mathbb{Z} \times [0, \infty)$ we will write ξ_t and refer to it as the unrestricted process.

We may also construct the GBT process ζ_t with the above Poisson processes and the help of some additional independent coin flips. Fix $\lambda_c < \lambda_2 < \lambda_1$, and suppose $\lambda = \lambda_1$ in the construction just given. Independently of everything else, label the arrows determined by the $B_n^{x,y}$ with a "1-only" sign with probability $(\lambda_1 - \lambda_2)/\lambda_1$. Call an active path up from (x, s) to (y, t) a 2-path if none of its arrows are 1-only arrows. Given ζ_0 , we may now construct ζ_t as follows. First, for all $t > 0$ and $x \in \mathbb{Z}$, put $\zeta_t(x) = 2$ if for some site y with $\zeta_0(y) = 2$ there is an active 2-path up from $(y, 0)$ to (x, t) . Next, for all other (x, t) put $\zeta_t(x) = 1$ if for some site y with $\zeta_0(y) = 1$ there is an active path up from $(y, 0)$ to (x, t) with the property that no vertical segments in the path contain a point (z, u) such that $\zeta_u(z) = 2$. Otherwise set $\zeta_t(x) = 0$. A little thought shows that ζ_t is the GBT process with the rates given in (12). The process of 2's is a contact process with infection parameter λ_2 , and in the absence of 2's, the process of 1's is a contact process with infection parameter λ_1 .

3 Proof of Theorem 1

The space-time regions \mathcal{Y}_{jk} . We will modify somewhat the standard approach of constructing a mapping from appropriate space-time regions of the construction just given to an oriented-percolation model with the property that survival of the percolation process implies survival of the contact process. We will call the regions \mathcal{Y}_{jk} , they will be defined using the parallelograms of Section VI.3 of [6].

Let \mathcal{L} be the lattice $\mathcal{L} = \{(j, k) \in \mathbb{Z}^2 : k \geq 0 \text{ and } j + k \text{ is even}\}$ with norm $\|(j, k)\| = 1/2(|j| + |k|)$. Fix $0 < \beta < \alpha/3$ and $M > 0$ so that $M\beta/2$ and $M\alpha$ are integers. Later we will set $\alpha = \alpha(\lambda)$ and take β small. For $(j, k) \in \mathcal{L}$, L_{jk} and R_{jk} are the “large” space-time parallelograms in $\mathbb{Z} \times [0, \infty)$ given by:

$$L_{jk} = M(j(\alpha - \beta), k) + L_{00}, \quad R_{jk} = M(j(\alpha - \beta), k) + R_{00}$$

where

$$L_{00} = \{(x, t) \in \mathbb{Z} \times [0, M(1 + \beta/\alpha)] : M\beta/2 \leq x + \alpha t \leq 3M\beta/2\}$$

$$R_{00} = \{(x, t) \in \mathbb{Z} \times [0, M(1 + \beta/\alpha)] : -3M\beta/2 \leq x - \alpha t \leq -M\beta/2\}.$$

We will also need the “small” parallelograms

$$L_{jk}^{small} = M(j(\alpha - \beta), k) + L_{00}^{small}, \quad R_{jk}^{small} = M(j(\alpha - \beta), k) + R_{00}^{small}$$

where

$$L_{00}^{small} = \{(x, t) \in \mathbb{Z} \times [0, M\frac{3\beta}{2\alpha}] : M\beta/2 \leq x + \alpha t \leq 3M\beta/2\}$$

$$R_{00}^{small} = \{(x, t) \in \mathbb{Z} \times [0, M\frac{3\beta}{2\alpha}] : -3M\beta/2 \leq x - \alpha t \leq -M\beta/2\}.$$

It is important to note that $L_{00}^{small} \subset L_{00}$, $R_{00}^{small} \subset R_{00}$, and

$$R_{jk} \cap L_{jk} = R_{jk} \cap L_{jk}^{small} = R_{jk}^{small} \cap L_{jk},$$

as shown in Figure 1.




Figure 1: Large parallelograms L_{00} and R_{00} . The shaded region is L_{00}^{small} .

We can now define the new objects \mathcal{Y}_{jk} which will be used to construct our oriented percolation process. As is the case with the parallelograms, the \mathcal{Y}_{jk} will be certain

translates of \mathcal{Y}_{00} , and depend on two fixed integers ℓ, d which satisfy $\ell \geq 2$ and $d \geq 0$ with $\ell > d$. We will form \mathcal{Y}_{00} by sticking together ℓ big right parallelograms, connected with appropriate small left parallelograms, and then two branches of d and $d + 1$ big left parallelograms connected by small right parallelograms. Figure 2 shows examples of \mathcal{Y}_{00} with parameters $\ell = 5$ and $d = 0, 1, 2$. It seems simplest to define \mathcal{Y}_{00} in stages, beginning with $\mathcal{Y}_{00}^0 = R_{00}$.

1. Attach ℓ big right parallelograms with ℓ small parallelograms to connect them:

$$\mathcal{Y}_{00}^1 = \mathcal{Y}_{00}^0 \cup \left(\bigcup_{i=1}^{\ell} (R_{ii} \cup L_{ii}^{small}) \right).$$

2. Attach one big left parallelogram: $\mathcal{Y}_{00}^2 = \mathcal{Y}_{00}^1 \cup L_{\ell, \ell}$.

3. If $d = 0$ set $\mathcal{Y}_{00} = \mathcal{Y}_{00}^2$. If $d \geq 1$, attach another big left parallelogram:

$$\mathcal{Y}_{00}^3 = \mathcal{Y}_{00}^2 \cup L_{\ell+1, \ell+1}.$$

4. If $d = 1$, attach another big left and small right parallelogram:

$$\mathcal{Y}_{00}^4 = \mathcal{Y}_{00}^3 \cup (L_{\ell-1, \ell+1} \cup R_{\ell-1, \ell+1}^{small})$$

and set $\mathcal{Y}_{00} = \mathcal{Y}_{00}^4$. If $d \geq 2$, attach two branches, to reach ‘‘height’’ $\ell + d + 1$, of big left parallelograms with small right parallelograms as connectors:

$$\mathcal{Y}_{00}^4 = \mathcal{Y}_{00}^3 \cup \left(\bigcup_{i=0}^{d-1} (L_{\ell-i, \ell+i} \cup R_{\ell-i, \ell+i}^{small}) \cup (L_{\ell+1-i, \ell+1+i} \cup R_{\ell+1-i, \ell+1+i}^{small}) \right).$$

5. If $d \geq 2$, attach a final big left parallelogram and small right parallelogram:

$$\mathcal{Y}_{00}^5 = \mathcal{Y}_{00}^4 \cup L_{\ell-d, \ell+d} \cup R_{\ell-d, \ell+d}^{small}$$

and put $\mathcal{Y}_{00} = \mathcal{Y}_{00}^5$.

Having defined \mathcal{Y}_{00} we set

$$\mathcal{Y}_{jk} = M([k(\ell - d) + j](\alpha - \beta), k(\ell + d + 1)) + \mathcal{Y}_{00}, \quad (j, k) \in \mathcal{L}.$$

The percolation variables U_{jk} . Let \mathcal{O}_{jk} be the event that for every parallelogram \mathcal{P} in \mathcal{Y}_{jk} there is an active path in the graphical representation of the contact process which stays entirely in \mathcal{P} and connects some point in the bottom edge of \mathcal{P} to some point in the the top edge of \mathcal{P} . Thus on \mathcal{O}_{jk} there is some point in the bottom edge of \mathcal{Y}_{jk} with the property that there are active paths in \mathcal{Y}_{jk} connecting this point to the top edge of every parallelogram in \mathcal{Y}_{jk} , and in particular to the top edges of the two top parallelograms \mathcal{Y}_{jk} . This means that on \mathcal{O}_{jk} there is a point in the bottom edge of \mathcal{Y}_{jk} and active paths in \mathcal{Y}_{jk} connecting this point to the bottom edges of both $\mathcal{Y}_{j-1, k+1}$ and $\mathcal{Y}_{j+1, k+1}$.

It is a consequence of Lemma VI.3.17 in [6] that $P(\mathcal{O}_{00})$ is close to 1 for large M .




Figure 2: \mathcal{Y}_{00} with $\ell = 5$, $d = 0, 1, 2$.

Lemma 6. For $0 < \beta < \alpha/3$, $\lim_{M \rightarrow \infty} P(\mathcal{O}_{00}) = 1$.

Proof: As in [6] let \mathcal{E}_{jk} to be the event that there is an active path in the graphical representation of the contact process which goes from the bottom edge of R_{jk} to the top edge, always staying entirely within R_{jk} , and also that there is an active path from the bottom edge of L_{jk} to the top edge, always staying entirely within L_{jk} . It is clear that the probability of connecting the bottom edge of a small parallelogram to its top edge by an active path staying in the parallelogram is bounded below by $P(\mathcal{E}_{00})$. By Lemma 3.17 in [6], for $0 < \beta < \alpha/3$, $\lim_{M \rightarrow \infty} P(\mathcal{E}_{00}) = 1$. In the construction of \mathcal{Y}_{00} there are most $h = 2\ell + 4d$ (if $d \geq 1$) or $h = 2\ell + 1$ (if $d = 0$) parallelograms used. It follows from positive correlations that $P(\mathcal{O}_{00}) \geq P(\mathcal{E}_{jk})^h$, and thus $\lim_{M \rightarrow \infty} P(\mathcal{O}_{00}) = 1$ ■

For $(j, k) \in \mathcal{L}$ let $U_{jk} = 1_{\mathcal{O}_{jk}}$. Then $P(U_{jk} = 1) = P(\mathcal{O}_{00})$ does not depend on (j, k) . Furthermore, the U_{jk} are 1-dependent, meaning that if $I \subset \mathcal{L}$ is such that $\|(j, k) - (j', k')\| > 1$ for all $(j, k) \neq (j', k') \in I$, then the $U_{jk}, (j, k) \in I$ are independent. This is because the corresponding space-time regions $\mathcal{Y}_{jk}, \mathcal{Y}_{j'k'}$ are disjoint. Using the U_{jk} we may construct a 1-dependent oriented percolation process in the usual way. A path in \mathcal{L} is a sequence $(j_1, k_1), \dots, (j_n, k_n)$ of points of \mathcal{L} which satisfies $k_{i+1} = k_i + 1$ and $j_{i+1} = j_i \pm 1$ for all $1 \leq i \leq n - 1$. The path is said to be open if $U_{j_i, k_i} = 1$ for each $1 \leq i \leq n - 1$. It is clear from the properties of the \mathcal{O}_{jk} that if $(j_1, k_1), \dots, (j_n, k_n)$ is an open path in \mathcal{L} then there must an active path in the graphical representation from the bottom edge of \mathcal{Y}_{j_1, k_1} to the bottom edge of \mathcal{Y}_{j_n, k_n} .

If we let Ω_∞ be the event that there is an infinite open path in \mathcal{L} starting at $(0, 0)$, then by Lemma 6 above and Theorem VI.3.19 of [6],

$$\lim_{M \rightarrow \infty} P(\Omega_\infty) = 1. \quad (15)$$

Survival of $\xi_t^{\mathcal{W}}$. Let $\mathcal{Y} = \mathcal{Y}(\ell, d, M) = \bigcup_{k=0}^{\infty} \bigcup_{j=-k}^k \mathcal{Y}_{jk}$. On Ω_∞ there must be an infinite active path in the graphical representation starting at some $(x, 0)$, $x \in [-3M\beta/2, -M\beta/2]$, which lies entirely in \mathcal{Y} . Thus if \mathcal{W} is any space-time region such that $\mathcal{Y} \subset \mathcal{W}$, and $\xi_t^{\mathcal{W}}$ is the \mathcal{W} -restricted contact process starting from $\{x : (x, 0) \subset \mathcal{W}\}$, then $\xi_t^{\mathcal{W}} \neq \emptyset \forall t \geq 0$ on Ω_∞ . We will prove the following.

Claim. Assume (7) holds and $\alpha = \alpha(\lambda)$. Then there exists $0 < \beta < \alpha/3$ and integers ℓ', d' such that for all $M > 0$,

$$\mathcal{Y}(\ell', d', M/\alpha(\ell' + 3)) \subset \mathcal{W}(\alpha_l, \alpha_r, M) - (M/(\ell' + 3), 0). \quad (16)$$




Figure 3: $\mathcal{Y}_{00}, \mathcal{Y}_{1,1}, \mathcal{Y}_{-1,1}$

Given (16), it follows from translation invariance and (15) that

$$P(\xi_t^{\mathcal{W}(\alpha_l, \alpha_r, M)} \neq \emptyset \forall t \geq 0) \geq P(\Omega_\infty) \rightarrow 1 \text{ as } M \rightarrow \infty,$$

proving (8).

To prove (16) we first suppose that ℓ, d , are positive integers with $d < \ell$ and $M > 0$. For $(j, k) \in \mathcal{L}$, the left upper corner of L_{jk} is $(M(j(\alpha - \beta) - \alpha - \beta/2), M(k + 1 + \beta/\alpha))$, and the right bottom corner of L_{jk} is $(M(j(\alpha - \beta) + 3\beta/2), Mk)$. A little thought shows that \mathcal{Y} must be contained in the space-time region bounded by the following two lines and the x -axis. The first line connects the leftmost point of the top edge of \mathcal{Y}_{00} with the leftmost point of the top edge of $\mathcal{Y}_{-1,1}$, which are the left upper corner of $L_{\ell-d, \ell+d}$ and the left upper corner of $L_{2(\ell+d)-1, 2(\ell+d)+1}$, namely, the points $(M((\ell - d)(\alpha - \beta) - \alpha - \beta/2), M(\ell + d + 1 + \beta/\alpha))$ and $(M(2(\ell - d)(\alpha - \beta) - 2\alpha + \beta/2), M(2(\ell + d + 1) + \beta/\alpha))$. The slope of this line is

$$s_l = \frac{\ell + d + 1}{\ell - d - 1} \frac{1}{\alpha - \beta} \quad (17)$$

and it contains the point $(x_l, 0)$ where $x_l = -M(3\beta/2 + \beta/\alpha s_l)$. The second line connects the rightmost point of \mathcal{Y}_{00} with the rightmost point of $\mathcal{Y}_{1,1}$, the bottom right corner of $L_{\ell+1, \ell+1}$ and the bottom right $L_{2(\ell+1)-d, 2(\ell+1)+d}$, namely, the points $(M((\ell+1)(\alpha - \beta) + 3\beta/2), M(\ell+1))$ and $(M((2(\ell+1) - d)(\alpha - \beta) + 3\beta/2), M(2(\ell+1) + d))$. The slope of this line is

$$s_r = \frac{\ell + d + 1}{\ell - d + 1} \frac{1}{\alpha - \beta} \quad (18)$$

and it contains the point $(x_r, 0)$ where $x_r = M((\ell + 1)(\alpha - \beta - 1/s_r) + 3\beta/2)$.

This analysis shows that $\mathcal{Y}(\ell, d, M)$ is contained in the wedge $\mathcal{W}(1/s_l, 1/s_r, M') + (x_l, 0)$, where $M' = x_r - x_l$. A little algebra shows that $-M\alpha < x_l < x_r < M\alpha(\ell + 2)$, and thus

$$\mathcal{Y}(\ell, d, M) \subset \mathcal{W}(1/s_l, 1/s_r, M\alpha(\ell + 3)) - (M\alpha, 0). \quad (19)$$

We now set $s_\ell = 1/\alpha_\ell, s_r = 1/\alpha_r$ and solve (17) and (18) for d and ℓ , obtaining

$$\ell = \frac{s_r(s_\ell(\alpha - \beta) + 1)}{s_\ell - s_r}, \quad d = \frac{s_\ell(s_r(\alpha - \beta) - 1)}{s_\ell - s_r}. \quad (20)$$


Unfortunately, ℓ, d need not be integers. To deal with this problem we first note that if $s_l \geq s'_l > s_r$ then for any M , the wedge $\mathcal{W}(\alpha_l, \alpha_r, M)$ contains the narrower wedge $\mathcal{W}(1/s'_\ell, 1/s_r, M)$. If we can find s'_ℓ and $0 < \beta < \alpha/3$ such that

$$\ell' = \frac{s_r(s'_\ell(\alpha - \beta) + 1)}{s'_\ell - s_r} \text{ and } d' = \frac{s'_\ell(s_r(\alpha - \beta) - 1)}{s'_\ell - s_r} \quad (21)$$

are both integers, then (16) follows from (19).

We can find s'_ℓ, β as follows. Let $m_0 = 3/\alpha s_r$ and take any integer $m > m_0$ such that $s_r \frac{m}{m-1} < s_l$. Put $s'_l = s_r \frac{m}{m-1}$, so that $s_l > s'_l > s_r$. Since $m > 3/\alpha s_r, 1/3\alpha m s_r > 1$ and the interval $(\frac{2}{3}\alpha m s_r, \alpha m s_r)$ must contain at least one integer. Since $\alpha s_r > 1$, the right endpoint of this interval is greater than m . Choose any integer $c \geq m$ from the interval and put $\beta = \alpha - \frac{c}{m s_r}$. Then $0 < \beta < \alpha/3$ and $s_r(\alpha - \beta) = c/m$. A little algebra shows that ℓ', d' given in (21) are the integers $\ell' = c + m - 1, d' = c - m$, and we are done.

Figure 4: Wedge containing \mathcal{Y}



4 Proof of Theorem 2

We begin by analyzing the rightmost particle. Let $\mathcal{W}(\alpha_r, M) = \{(x, t) : t \geq 0, x \in (-\infty, M + \alpha_r t] \cap \mathbb{Z}\}$ and consider the restricted contact process $\xi_t^{\mathcal{W}(\alpha_r, M)}$ with initial

state $\xi_0^{\mathcal{W}(\alpha_r, M)} = (-\infty, M] \cap \mathbb{Z}$. Let \bar{r}_t be the right-edge process for $\xi_t^{\mathcal{W}}$, $\bar{r}_t = \max\{x : \xi_t^{\mathcal{W}(\alpha_r, M)}(x) = 1\}$. We claim that for every M ,

$$\lim_{t \rightarrow \infty} \frac{\bar{r}_t}{t} = \alpha_r \quad a.s. \quad (22)$$

By construction and (4), $\limsup_{t \rightarrow \infty} \bar{r}_t/t \leq \alpha_r$. For the lower bound, fix $0 < \varepsilon < \alpha_r$ and define the region $\mathcal{W}_\varepsilon = \mathcal{W}(\alpha_r - \varepsilon, \alpha_r, M)$ and restricted contact process $\xi_t^{\mathcal{W}_\varepsilon}$ with initial state $\xi_0^{\mathcal{W}_\varepsilon} = [0, M] \cap \mathbb{Z}$. Then $\xi_t^{\mathcal{W}_\varepsilon} \subset \xi_t^{\mathcal{W}(\alpha_r, M)}$, which implies that on the event $\{\xi_t^{\mathcal{W}_\varepsilon} \neq \emptyset \forall t \geq 0\}$, $\liminf_{t \rightarrow \infty} \bar{r}_t/t \geq \alpha_r - \varepsilon$. Theorem 1 now implies we must have $\liminf_{t \rightarrow \infty} \bar{r}_t/t \geq \alpha_r - \varepsilon$ a.s., completing the proof of (22).

It is a consequence of the nearest-neighbor interaction mechanism that for any $\alpha_l < \alpha_r$ and M , with $\mathcal{W} = \mathcal{W}(\alpha_l, \alpha_r, M)$,

$$\xi_t^{\mathcal{W}}(x) = \xi_t^{\mathcal{W}(\alpha_r, M)}(x) \forall x \in [l_t^{\mathcal{W}}, r_t^{\mathcal{W}}] \text{ on } \{\xi_t^{\mathcal{W}} \neq \emptyset\}.$$

This implies $r_t^{\mathcal{W}} = \bar{r}_t$ on $\{\xi_t^{\mathcal{W}} \neq \emptyset\}$, and so by (22), $\lim_{t \rightarrow \infty} r_t^{\mathcal{W}}/t = \alpha_r$. We omit the similar argument proving $\lim_{t \rightarrow \infty} l_t^{\mathcal{W}}/t = \alpha_l$.

For (11), let $\xi_t^{\mathbb{Z}}$ denote the unrestricted process with initial state $\xi_0^{\mathbb{Z}} = \mathbb{Z}$, and let ξ_t^ν be the unrestricted process constructed as in Section 2 with initial state ξ_0^ν which has law ν , independent of the Poisson processes. We observe again that the nearest-neighbor interaction implies

$$\xi_t^{\mathbb{Z}}(x) = \xi_t^\nu(x) \forall x \in [l_t^{\mathcal{W}}, r_t^{\mathcal{W}}] \text{ on } \{\xi_t^{\mathcal{W}} \neq \emptyset \forall t \geq 0\}.$$

Standard exponential estimates for $P(\xi_t^{\mathbb{Z}}(x) \neq \xi_t^\nu(x)) = P(\xi_t^{\mathbb{Z}}(x) = 1) - P(\xi_t^\nu(x) = 1)$, a ‘‘filling in’’ argument and Borel-Cantelli (see Theorem I.2.30 of [7]) imply that for any $A > 0$,

$$P(\xi_t^{\mathbb{Z}} = \xi_t^\nu \text{ on } [-At, At] \text{ for all large } t) = 1$$

Combining the above with (10) gives (11).

5 Proof of Corollary 4

We will make use of the graphical construction in Section 2 and define independent events $\Omega_1, \Omega_2, \Omega_3$, each with positive probability, and such that $\|\zeta_t\|_1 \rightarrow \infty$ as $t \rightarrow \infty$ on their intersection.

First, since $\alpha(\lambda)$ is strictly increasing we may choose $\alpha(\lambda_2) < \alpha_l < \alpha_r < \alpha(\lambda_1)$. Fix $M > 2$ and write \mathcal{W} for $\mathcal{W}(\alpha_l, \alpha_r, M)$. The first event is

$$\Omega_1 = \{\text{there is no active 2-path from any } (x, 0), x < 0, \text{ to any point of } \mathcal{W}(\alpha_l, \alpha_r, M)\}.$$

Since the process of 2’s is a contact process with parameter λ_2 , and $\alpha(\lambda_2) < \alpha_l$, it follows from (4) that Ω_1 has positive probability.

For the second event, choose $x_0 \in \mathbb{Z}$ and $t_0 > 0$ such that $x_0 = \alpha_l t_0$ and $(x, t_0) \subset \mathcal{W}$ for all $x \in [x_0, x_0 + M] \cap \mathbb{Z}$. Since $M > 2$ the event,

$$\Omega_2 = \{\text{there is an active path in } \mathcal{W} \text{ from } (0, 0) \text{ to each of } (x, t_0), x \in [x_0, x_0 + M] \cap \mathbb{Z}\}$$

has positive probability.

For the third event, define, for $t \geq t_0$,

$$A_t = \{y : \text{there is an infinite active path in } \mathcal{W} \text{ from } (x, t_0) \text{ to } (y, t) \\ \text{for some } x \in [x_0, x_0 + M] \cap \mathbb{Z} \}$$

and put $\Omega_3 = \{|A_t| \rightarrow \infty \text{ as } t \rightarrow \infty\}$. It follows from Theorems 1 and 2 that Ω_3 has positive probability.

The events Ω_i are independent since they are defined in terms of our Poisson processes over disjoint space-time regions. Furthermore, it is easy to see from Remark 3 that $\|\zeta_t\|_1 \rightarrow \infty$ on their intersection, so we are done.

References

- [1] Chayes, J.T. and Chayes, L. (1986) Critical points and intermediate phases on wedges of \mathbb{Z}^d , *Journal of Physics A: Mathematical and General* **19**, 3033-3048.
- [2] Durrett, R. (1980) On the growth of one-dimensional contact process, *Ann. Probab.* **8**, 890-907.
- [3] Durrett, R. and Swindle, G. (1991) Are there bushes in a forest? *Stoch. Proc. Appl.* **37**, 19-31.
- [4] Grimmett, G. (1983) Bond percolation on subsets of the square lattice, and the transition between one-dimensional and two-dimensional behavior, *Journal of Physics A: Mathematical and General* **16**, 599-604.
- [5] Harris, T.E. (1974) Contact interactions on a lattice, *Ann. Probab.* **2**, 969-988.
- [6] Liggett, T. M. (1985) *Interacting Particle Systems*, Springer-Verlag, Berlin.
- [7] Liggett, T. M. (1985) *Stochastic Interacting Systems: Contact, Voter and Exclusion Processes*, Springer-Verlag, Berlin.

J. Theodore Cox

Department of Mathematics, Syracuse University Syracuse, NY 13244
e-mail: jtcox@syr.edu

Nevena Marić

Department of Mathematics and CS, University of Missouri- St. Louis St. Louis, MO 63121
e-mail: maricn@umsl.edu

Rinaldo Schinazi

Department of Mathematics, University of Colorado-Colorado Springs Colorado Springs, CO 80933
e-mail: rschinaz@uccs.edu