

12-15-2010

Warped Product Einstein Metrics Over Spaces with Constant Scalar Curvature

Chenxu He

Lehigh University, he.chenxu@lehigh.edu

Peter Petersen

University of California - Los Angeles, petersen@math.ucla.edu

William Wylie

University of Pennsylvania, wylie@math.upenn.edu

Follow this and additional works at: <http://surface.syr.edu/mat>

 Part of the [Mathematics Commons](#)

Recommended Citation

He, Chenxu; Petersen, Peter; and Wylie, William, "Warped Product Einstein Metrics Over Spaces with Constant Scalar Curvature" (2010). *Mathematics Faculty Scholarship*. Paper 131.

<http://surface.syr.edu/mat/131>

This Article is brought to you for free and open access by the Department of Mathematics at SURFACE. It has been accepted for inclusion in Mathematics Faculty Scholarship by an authorized administrator of SURFACE. For more information, please contact surface@syr.edu.

WARPED PRODUCT EINSTEIN METRICS OVER SPACES WITH CONSTANT SCALAR CURVATURE

CHENXU HE, PETER PETERSEN, AND WILLIAM WYLIE

ABSTRACT. In this paper we study warped product Einstein metrics over spaces with constant scalar curvature. We call such a manifold rigid if the universal cover of the base is Einstein or is isometric to a product of Einstein manifolds. When the base is three dimensional and the dimension of the fiber is greater than one we show that the space is always rigid. We also exhibit examples of solvable four dimensional Lie groups that can be used as the base space of non-rigid warped product Einstein metrics showing that the result is not true in dimension greater than three. We also give some further natural curvature conditions that characterize the rigid examples in higher dimensions.

1. INTRODUCTION

A $(\lambda, n + m)$ -Einstein manifold (M^n, g, w) is a complete Riemannian manifold, possibly with boundary, and a smooth function w on M satisfying:

$$(1.1) \quad \begin{aligned} \text{Hess}w &= \frac{w}{m}(\text{Ric} - \lambda g) \\ w &> 0 \text{ on } \text{int}(M), \\ w &= 0 \text{ on } \partial M. \end{aligned}$$

When $m = 1$, we make the additional assumption that $\Delta w = -\lambda w$.

$(\lambda, n + m)$ -Einstein metrics are also called *m-Quasi Einstein* manifolds and the case where ∂M is empty was studied earlier in [CSW] and [KK]. We also studied this equation in [HPW1] and showed that many of the results from these earlier works generalize to the case where the boundary is non-empty.

The $(\lambda, n + m)$ -Einstein equation has a natural geometric interpretation. Namely, if $m > 1$ is an integer, then (M^n, g, w) is a $(\lambda, n + m)$ -Einstein manifold if and only if there is a smooth $(n + m)$ -dimensional warped product Einstein metric (with no boundary) with base space M (see [HPW1, Proposition 1.1]). Thus we can study warped product Einstein metrics by analyzing the $(\lambda, n + m)$ -Einstein equation on the lower dimensional base space. While the motivation requires m to be an integer, there is no reason to restrict to this case in any of our results. When $m = 1$, solutions are also called the *static metrics* and have been studied thoroughly for their connection to general relativity and the positive mass theorem [Co]. The equation is also closely related to considerations in optimal transport [Vi] and comparison geometry [WW].

2000 *Mathematics Subject Classification.* 53B20, 53C30.

The second author was supported in part by NSF-DMS grant 1006677.

The third author was supported in part by NSF-DMS grant 0905527.

Many Einstein metrics can be constructed as warped products. In fact the first non-trivial example of an Einstein metric, the Schwarzschild metric, is a 4-dimensional doubly warped product metric on $\mathbb{R}^2 \times \mathbb{S}^2$. In our context it can be viewed in two ways, either as a $(0, 2 + 2)$ -Einstein metric on \mathbb{R}^2 , or as a $(0, 3 + 1)$ -Einstein metric on $[0, \infty) \times \mathbb{S}^2$. Much more recently C. Böhm constructed interesting warped product Einstein metrics on spheres and product of spheres. These examples give rotationally symmetric $(\lambda, n + m)$ -Einstein metrics on hemispheres and spheres respectively when $n = 3, 4, 5, 6, 7$ (see [Bö1]). Also see [LPP, Bö2] for more examples.

The Böhm metrics are in contrast to the solutions to the analogous *gradient Ricci soliton* equation,

$$\text{Ric} + \text{Hess}f = \lambda g.$$

There is a well known classification theorem of three dimensional gradient Ricci solitons with $\lambda > 0$ following from the works of Ivey[Iv], Hamilton[Ha], and Perelman[Per]. Thus the Böhm metrics show that there are more examples of $(\lambda, 3 + m)$ -Einstein metrics than gradient Ricci solitons. It is then natural to ask whether one can classify $(\lambda, 3 + m)$ -Einstein metrics under additional natural curvature assumptions. Before we state our first result in this direction, we require a definition.

Definition 1.1. A $(\lambda, n + m)$ -Einstein manifold (M, g, w) is called *rigid* if it is Einstein or its universal cover is a product of Einstein manifolds.

There is an explicit list of the possible rigid examples, in particular we will see that there are at most two different Einstein constants in the product, one of which is λ , see Section 2. Our first result is that, in dimension three, constant scalar curvature characterizes rigidity.

Theorem 1.2. *A $(\lambda, 3 + m)$ -Einstein manifold with $m > 1$ has constant scalar curvature if and only if it is rigid. In particular, if the manifold has no boundary, then it is a quotient of \mathbb{S}^3 , $\mathbb{S}^2 \times \mathbb{R}$, \mathbb{R}^3 , $\mathbb{H}^2 \times \mathbb{R}$, or \mathbb{H}^3 with the standard metrics.*

Remark 1.3. In [Se] H. Seshadri considered compact $(\lambda, 3 + 1)$ -Einstein manifold where the total space admits a non-trivial circle action. Under certain further conditions on the circle action, he showed that the total space is either the standard \mathbb{S}^4 with the base 3-disk \mathbb{D}^3 or the Riemannian product $\mathbb{S}^2 \times \mathbb{S}^2$ with the base $[-\frac{\pi}{2}, \frac{\pi}{2}] \times \mathbb{S}^2$.

Note, on the other hand, that every $(\lambda, n + 1)$ -Einstein metric has constant scalar curvature. While constant scalar curvature may appear to be a strong assumption, we also exhibit examples showing that this theorem is not true in higher dimensions.

Theorem 1.4. *For each $m > 0$, there are 4-dimensional solvable Lie groups with left invariant $(\lambda, 4 + m)$ -Einstein metrics that are not rigid.*

Remark 1.5. These examples are also in stark contrast to the gradient Ricci soliton case, where all homogeneous gradient Ricci solitons are rigid. As m goes to infinity these examples converge to a Riemannian product $\mathbb{R} \times H^3$ where H is a homogeneous (non-gradient) Ricci soliton, see Theorem 6.10.

Remark 1.6. We also note that the left invariant metrics we construct on solvable Lie groups are not solvsoliton metrics unless they are isometric to $\mathbb{R} \times H$ or are rigid.

We are then led to the question of whether we can classify $(\lambda, n + m)$ -Einstein metrics with constant scalar curvature in higher dimensions under natural additional assumptions. This problem is also addressed in [CSW] where it is shown that every compact $(\lambda, n + m)$ -Einstein metric with constant scalar curvature which does not have boundary must be a trivial λ -Einstein metric. They also show that when $\lambda = 0$ the metric must be Ricci flat, our next result is the extension of this result to the non-empty boundary case.

Theorem 1.7. *Suppose (M, g, w) is a complete $(0, n + m)$ -Einstein manifold with constant scalar curvature, then (M, g) is either Ricci flat without boundary or it is isometric to $(F \times [0, \infty), g_F + dr^2)$ where (F, g_F) is Ricci flat and $w = w(r)$ is a linear function.*

Classification theorems for gradient Ricci solitons with constant scalar curvature were also considered by the last two authors in [PW3]. They show that a gradient Ricci soliton is rigid if it has constant scalar curvature and all the sectional curvatures in the direction of the gradient of the potential function are zero. Our next result appears to be a stronger result for $(\lambda, n + m)$ -Einstein metrics.

Theorem 1.8. *Suppose (M, g, w) is a complete $(\lambda, n + m)$ -Einstein manifold with constant scalar curvature and $\lambda > 0 (< 0)$, if $\text{Ric}(\nabla w, \nabla w) \leq 0 (\geq 0)$, then M is isometric to the product $\mathbb{R} \times N$ where N is a λ -Einstein manifold.*

Remark 1.9. It turns out that the assumption about Ricci curvature is equivalent to the scalar curvature being bounded between $n\lambda$ and $(n - 1)\lambda$. Thus this theorem can also be viewed as a generalization of Proposition 3.6 in [CSW] which states that for $\lambda < 0$ if the scalar curvature is constant then it is bounded below by $n\lambda$, and is equal to $n\lambda$ if and only if the metric is λ -Einstein.

This theorem shows that only a limited number of rigid $(\lambda, n + m)$ -Einstein metrics have vanishing sectional curvatures in the “radial” direction. This is in contrast with the gradient soliton case where all rigid examples have vanishing radial curvatures. Our next theorem is a characterization of all rigid $(\lambda, n + m)$ -Einstein metrics in terms of scalar curvature and the sectional curvatures in the radial direction.

Theorem 1.10. *Suppose (M, g, w) is a complete simply connected non-trivial $(\lambda, n + m)$ -Einstein manifold with constant scalar curvature and $\lambda \neq 0$. If the Riemann curvature tensor does not grow exponentially, and the radial sectional curvature satisfies*

$$(1.2) \quad R(X, \nabla w, \nabla w, X) = \frac{|\nabla w|^2}{m} (\lambda |X|^2 - \text{Ric}(X, X)),$$

then the manifold is rigid.

Remark 1.11. We only need to assume the curvature growth condition in the case when the fiber in the warped product construction is Ricci flat.

In [HPW1] we defined two natural tensors, P and Q , which are modifications of the Ricci tensor and the Riemann curvature tensor. The equation in (1.2) is equivalent to the radial flatness of Q , i.e., $Q(X, \nabla w, \nabla w, X) = 0$.

Theorem 1.10 also has the same corollary as the result in the gradient Ricci soliton case. Note in the corollary below, we do not need the curvature growth assumption.

Corollary 1.12. *If (M, g, w) is a complete simply connected non-trivial $(\lambda, n+m)$ -Einstein manifold with harmonic curvature, then it is rigid.*

The paper is organized as follows. We classify the rigid examples in Section 2. In Section 3, we derive some identities for the tensors P and Q on $(\lambda, n+m)$ -Einstein manifolds with constant scalar curvature. In Section 4 we turn our attention to proving the classification theorems. We first prove Theorems 1.7 and 1.8. Theorem 1.2, the three dimensional classification, is then an application of these two results. In Section 5 we discuss Theorem 1.10 and in Section 6 we construct the non-rigid metrics on solvable Lie groups. Further investigations in the homogeneous case will be the subject of future paper. In Appendix A, we prove a result for the integrability of eigen-distribution of Ricci tensor that is used in the proof of Theorem 1.10, see Theorem A.1.

Acknowledgment: The major part of this paper was done when the first author was a visiting scholar at Department of Math, University of Pennsylvania. He would like to thank the university for their hospitality.

2. RIGID EXAMPLES

In this section, we give a classification of rigid $(\lambda, n+m)$ -Einstein manifolds. First recall

Definition 2.1. A $(\lambda, n+m)$ -Einstein manifold (M, g, w) is called *rigid* if it is Einstein or its universal cover is a product of Einstein manifolds.

In the gradient Ricci soliton case, if the metric splits, then each factor is a gradient soliton with the same value of λ and none of them has to be trivial. However in the $(\lambda, n+m)$ -Einstein case if the metric splits, then there are at most two Einstein constants and one factor is trivial.

Lemma 2.2. *Suppose (M, g, w) is a $(\lambda, n+m)$ -Einstein metric such that the metric splits as a product*

$$(M, g) = (M_1, g_1) \times (M_2, g_2)$$

Then one of the manifolds, say (M_1, g_1) , is a trivial $(\lambda, n+m)$ -Einstein manifold.

Proof. We write the $(\lambda, n+m)$ -Einstein equation in the $(1, 1)$ -tensor form:

$$\text{Ric} - \frac{m}{w} \nabla \nabla w = \lambda I.$$

The operator $E \mapsto \frac{m}{w} \nabla_E \nabla w$ preserves the splitting of the metric $(M, g) = (M_1 \times M_2, g_1 + g_2)$. Using the local coordinates x^j and writing $\nabla w = \alpha^j \partial_j$, we have $\nabla w = X_1 + X_2$ where X_i is a vector field on M_i , $i = 1, 2$. Choose a fixed point $(p, q) \in M_1 \times M_2$, then on the submanifold $\{p\} \times M_2$, we have

$$m \nabla X_2(x_2) = w(p, x_2) (\text{Ric}_{g_2} - \lambda I).$$

Therefore if (M_2, g_2) is not a λ -Einstein manifold, then $w(p, x_2)$ does not depend on the value of p , i.e., $X_1 \equiv 0$, which implies (M_1, g_1) is a trivial $(\lambda, n+m)$ -Einstein manifold. \square

The non-trivial $(\lambda, n + m)$ -Einstein manifolds which are also Einstein were classified in [CSW] and extended to manifolds with boundary in [HPW1]. The Einstein constant which is not λ is given by

$$(2.1) \quad \rho = \frac{(n-1)\lambda - \text{scal}}{m-1},$$

and all examples are listed in the next proposition.

Proposition 2.3. *Suppose that (M, g, w) is a non-trivial $(\lambda, n + m)$ -Einstein manifold which is also Einstein, then up to multiples of w and g , it is isometric to one of the examples in Table 1.*

M	g	w	λ	ρ	μ
$[-\frac{\pi}{2}, \frac{\pi}{2}]$	dr^2	$w(r) = \cos(r)$	m	0	$m-1$
$[0, \infty)$	dr^2	$w(r) = r$	0	0	$m-1$
$[0, \infty)$	dr^2	$w(r) = \sinh(r)$	$-m$	0	$m-1$
$(-\infty, \infty)$	dr^2	$w(r) = e^r$	$-m$	0	0
$(-\infty, \infty)$	dr^2	$w(r) = \cosh(r)$	$-m$	0	$-(m-1)$
\mathbb{D}^n	$dr^2 + \sin^2(r)g_{\mathbb{S}^{n-1}}$	$w(r) = \cos(r)$	$n+m-1$	$n-1$	$m-1$
$[0, \infty) \times F$	$dr^2 + g_F$	$w(r) = r$	0	0	$m-1$
$[0, \infty) \times N$	$dr^2 + \cosh^2(r)g_N$	$w(r) = \sinh(r)$	$-(n+m-1)$	$-(n-1)$	$m-1$
$(-\infty, \infty) \times F$	$dr^2 + e^{2r}g_F$	$w(r) = e^r$	$-(n+m-1)$	$-(n-1)$	0
\mathbb{H}^n	$dr^2 + \sinh^2(r)g_{\mathbb{S}^{n-1}}$	$w(r) = \cosh(r)$	$-(n+m-1)$	$-(n-1)$	$-(m-1)$

TABLE 1. Non-trivial $(\lambda, n + m)$ -Einstein manifolds that are also Einstein. Here \mathbb{S}^{n-1} has Ricci curvature $n-2$, F is Ricci flat and N has Ricci curvature $-(n-2)$.

Remark 2.4. The first five $(\lambda, 1 + m)$ -Einstein structures in dimension one can be viewed as degenerate cases of the last five examples. For instance, in Example 8 if N is a point, then we have Example 3.

Combining these two results allows us to easily classify all rigid $(\lambda, n + m)$ -Einstein manifolds.

Proposition 2.5. *A non-trivial complete rigid $(\lambda, n + m)$ -Einstein manifold (M, g, w) is one of the examples in Table 1, or its universal cover \widetilde{M} splits as*

$$\begin{aligned} \widetilde{M} &= (M_1, g_1) \times (M_2, g_2) \\ w &= (c, w_2), \end{aligned}$$

where c is a constant, (M_1, g_1) is a trivial $(\lambda, n + m)$ -Einstein manifold and (M_2, g_2, w_2) is one of the examples in Table 1.

Proof. If M is Einstein, then by Proposition 2.3 it is one of the examples in Table 1. Otherwise the metric splits as a product of Einstein manifolds and Lemma 2.2 implies that the potential function w also splits and one of the factors, M_1 have

constant potential function and Ricci curvature λ . Now M_2 is both Einstein and $(\lambda, n+m)$ -Einstein and so applying Proposition 2.3 again shows that M_2 is in Table 1. \square

3. PRELIMINARIES

In this section we collect the formulas from [KK], [CSW] and [HPW1] which we will use later in the proof of the theorems. We apply some of these identities to give a classification of the possible forms of w on a $(\lambda, n+m)$ -Einstein manifold with $m > 1$ and constant scalar curvature. Finally we derive some properties about the critical point set of w .

Recall from [HPW1] that for a $(\lambda, n+m)$ -Einstein manifold with $m \neq 1$, we define

$$\begin{aligned}\rho(x) &= \frac{(n-1)\lambda - \text{scal}}{m-1} \\ P &= \text{Ric} - \rho g \\ Q &= R + \frac{2}{m}P \odot g + \frac{\rho - \lambda}{m}g \odot g\end{aligned}$$

where R is the (0,4) Riemann curvature tensor, and for any two symmetric (0,2)-tensor s and r , $s \odot r$ is the Kulkarni-Nomizu product defining a (0,4)-tensor

$$\begin{aligned}(s \odot r)(X, Y, Z, W) &= \frac{1}{2}(r(X, W)s(Y, Z) + r(Y, Z)s(X, W)) \\ &\quad - \frac{1}{2}(r(X, Z)s(Y, W) + r(Y, W)s(X, Z)).\end{aligned}$$

Note that we use the convention for the Riemann tensor that makes $R(X, Y, Y, X)$ have the same sign as the sectional curvature of the plane spanned by X and Y . The Kulkarni-Nomizu tensor is defined to be consistent with this choice.

Up to a dimensional constant, P can be viewed as the Ricci tensor associated with Q , i.e., if $\{E_i\}_{i=1}^n$ is an orthonormal frame, then

$$(3.1) \quad \sum_{i=1}^n Q(X, E_i, E_i, Y) = \frac{n+m-2}{m}P(X, Y).$$

Also note that the trace of P and ρ are related by the equation

$$(3.2) \quad \text{tr}P = (n-1)\lambda - (n+m-1)\rho.$$

When the scalar curvature is constant, ρ is constant on M and P is just the Ricci tensor shifted by a constant multiple of the metric. In this case the formulas in [CSW] and [HPW1] simplify significantly. The first set of formulas we will need involve P and the derivatives of the scalar curvature. Recall that for a $(\lambda, n+m)$ -Einstein manifold with $m > 1$,

$$\begin{aligned}\frac{w}{2}\nabla\rho &= P(\nabla w) \\ \frac{1}{2}\Delta(\text{scal}) + \frac{m+1}{2w}g(\nabla\text{scal}, \nabla w) &= (\lambda - \rho)\text{tr}(P) - |P|^2 \\ \text{div}(w^{m+1}P) &= 0.\end{aligned}$$

The first two equations are just formulas (3.11) and (3.12) in [CSW] rewritten in our notation, for the third identity see Proposition 5.6 in [HPW1]. When the scalar curvature is constant these identities give us the following formulas.

Proposition 3.1. *Let (M, g, w) be a $(\lambda, n + m)$ -Einstein manifold with constant scalar curvature and $m > 1$, then*

$$\begin{aligned} P(\nabla w) &= 0 \\ |P|^2 &= (\lambda - \rho)\text{tr}P = \text{constant}. \\ \text{div}(P) &= 0. \end{aligned}$$

Proof. The first two are obvious from the equations above because scal , ρ , and $\text{tr}P$ are all constant. For the third fact, note that $\text{div}(w^{m+1}P) = 0$ is equivalent to

$$\text{div}(P) = -\frac{(m+1)}{w}P(\nabla w).$$

□

There are two important corollaries of these formulas which we will find useful.

Corollary 3.2 ([CSW], Proposition 3.6). *Let (M, g, w) be a $(\lambda, n + m)$ -Einstein manifold with constant scalar curvature and $m > 1$, assume in addition that $\lambda \neq 0$, then the scalar curvature is bounded by $n\lambda$ and $n\rho$. Moreover if $\text{scal} = n\lambda$ or $n\rho$, then the manifold is Einstein.*

Proof. From the equation $|P|^2 = (\lambda - \rho)\text{tr}P$, we have

$$\left|P - \frac{\text{tr}P}{n}g\right|^2 = (\text{tr}P) \left(\lambda - \frac{\text{scal}}{n}\right) = -\frac{1}{n}(\text{scal} - n\rho)(n\lambda - \text{scal}).$$

Then the statement follows easily. □

Corollary 3.3. *Let (M, g, w) be a $(\lambda, n + m)$ -Einstein manifold with constant scalar curvature and $m > 1$. If $\lambda > 0 (< 0)$, then $\lambda - \rho > 0 (< 0)$.*

Proof. Suppose that $\lambda < 0$ and $\rho \leq \lambda$, then by (3.2), $\text{tr}(P) > 0$. Then the formula

$$|P|^2 = (\lambda - \rho)\text{tr}P \leq 0$$

shows that M is Einstein and $\rho = \lambda$. However, this case never occurs in Table 1 with $m > 1$ so we have a contradiction.

A similar argument shows $\rho - \lambda < 0$ if $\lambda > 0$. In fact, constant scalar curvature is not necessary for the conclusion in the $\lambda > 0$ case, see Proposition 5.4 in [HPW1]. □

The other set of formulas we will use involve the covariant derivatives of the tensors P and Q , and are proven as Proposition 6.2 in [HPW1].

$$(3.3) \quad \frac{w}{m}((\nabla_X P)(Y, Z) - (\nabla_Y P)(X, Z)) = -Q(X, Y, Z, \nabla w) - \frac{1}{m}(g \odot g)(X, Y, Z, P(\nabla w))$$

In the case of constant scalar curvature these give us the following identities.

Proposition 3.4. *Suppose (M, g, w) is a $(\lambda, n+m)$ -Einstein manifold with constant scalar curvature and $m > 1$, then*

$$\begin{aligned} \frac{w}{m} ((\nabla_X P)(Y, Z) - (\nabla_Y P)(X, Z)) &= \frac{w}{m} ((\nabla_X \text{Ric})(Y, Z) - (\nabla_Y \text{Ric})(X, Z)) = -Q(X, Y, Z, \nabla w), \\ \frac{w}{m} (\nabla_{\nabla w} P)(X, Y) &= -\left(\frac{w}{m}\right)^2 (\lambda - \rho) P(X, Y) + \left(\frac{w}{m}\right)^2 g(P(X), P(Y)) \\ &\quad + Q(\nabla w, X, Y, \nabla w). \end{aligned}$$

Proof. The first equation comes from combining the equation (3.3) with the fact that $P(\nabla w) = 0$ and ρ is constant. The second equation then follows from combining the first equation with the following formula

$$\begin{aligned} (\nabla_X P)(\nabla w, Y) &= P(\nabla_X \nabla w, Y) \\ &= \frac{w}{m} P((\text{Ric} - \lambda I)(X), Y) \\ &= \frac{w}{m} P((P + (\rho - \lambda)I)(X), Y), \end{aligned}$$

where we have assumed that X and Y are arbitrary parallel fields. \square

We obtain the following corollary from considering the last identity at a critical point of w .

Corollary 3.5. *If ∇w vanishes at $p \in M$, then*

$$P \circ (P - (\lambda - \rho)I) = 0.$$

The other very important formula we will utilize is the identity from [KK] which states that

$$(3.4) \quad w\Delta w + (m-1)|\nabla w|^2 + \lambda w^2 = \mu = \text{const.}$$

By tracing the $(\lambda, n+m)$ -Einstein equation, we also have

$$\Delta w = \frac{w}{m} (\text{scal} - n\lambda).$$

Letting $\bar{\mu} = \frac{\mu}{m-1}$ after a calculation the equation (3.4) then becomes

$$(3.5) \quad \bar{\mu} = \bar{k}w^2 + |\nabla w|^2, \quad \text{where} \quad \bar{k} = \frac{\lambda - \rho}{m}.$$

Since \bar{k} is constant when the scalar curvature is constant this tells us that the only possibilities for the form of the function w are the functions appearing in the rigid examples.

Proposition 3.6. *Suppose (M, g, w) is a non-trivial $(\lambda, n+m)$ -Einstein manifold with constant scalar curvature and $m > 1$.*

- *If $\bar{k} > 0$, then $\bar{\mu} > 0$ and*

$$w = \sqrt{\frac{\bar{\mu}}{\bar{k}}} \cos(\sqrt{\bar{k}}r),$$

for a distance function r .

- *If $\bar{k} = 0$, then $\bar{\mu} > 0$ and $w = \sqrt{\bar{\mu}}r$ for a distance function r .*

- If $\bar{k} < 0$, then the general form of w is

$$w = C_1 \exp\left(\sqrt{-\bar{k}r}\right) + C_2 \exp\left(-\sqrt{-\bar{k}r}\right)$$

where r is a distance function. More specifically we can express this as

$$\begin{aligned} w &= \exp\left(\sqrt{-\bar{k}r}\right), \text{ when } \bar{\mu} = 0, \\ w &= \sqrt{\frac{\bar{\mu}}{\bar{k}}} \cosh\left(\sqrt{-\bar{k}r}\right), \text{ when } \bar{\mu} < 0, \\ w &= \sqrt{-\frac{\bar{\mu}}{\bar{k}}} \sinh\left(\sqrt{-\bar{k}r}\right), \text{ when } \bar{\mu} > 0. \end{aligned}$$

Remark 3.7. Note that Corollary 3.3 shows that \bar{k} and λ always have the same sign.

Proof. Recall that a distance function is simply a smooth function whose gradient always has unit length. The proof uses the equation

$$\bar{k}w^2 + |\nabla w|^2 = \bar{\mu}.$$

When $\bar{k} = 0$ and $\bar{\mu} > 0$ we note that $\frac{w}{\sqrt{\bar{\mu}}}$ is a distance function.

When $\bar{\mu} = 0$ and $\bar{k} < 0$ the function $\log \frac{w}{\sqrt{-\bar{k}}}$ is a distance function.

When $\bar{k} > 0$ we note that $\bar{\mu} > 0$ and we can rewrite the formula as

$$\frac{1}{\bar{k}} \frac{|\nabla w|^2}{\left(\sqrt{\frac{\bar{\mu}}{\bar{k}}}\right)^2 - w^2} = 1$$

showing that $\sqrt{\bar{k}} \arccos\left(\frac{w}{\sqrt{\frac{\bar{\mu}}{\bar{k}}}}\right)$ is a distance function.

Finally when $\bar{k} < 0$ there are two cases depending on the sign of $\bar{\mu}$. The formula is then rewritten as

$$\frac{1}{-\bar{k}} \frac{|\nabla w|^2}{\left(\sqrt{-\frac{\bar{\mu}}{\bar{k}}}\right)^2 + w^2} = 1$$

when $\bar{\mu} > 0$, or

$$\frac{1}{\bar{k}} \frac{|\nabla w|^2}{\left(\sqrt{\frac{\bar{\mu}}{\bar{k}}}\right)^2 - w^2} = 1$$

when $\bar{\mu} < 0$. And we get the specific expression using arcsinh or arccosh. \square

Finally in this section we discuss two propositions about the critical point set of w that will be used in the proof of Theorem 1.10.

Proposition 3.8. *Suppose (M, g, w) is a $(\lambda, n + m)$ -Einstein manifold with constant scalar curvature and the set of critical points of w is non-empty. Then all connected components have the same dimension. Furthermore let N be a connected component, then normal vectors to N are 0 eigenvectors for P and tangent vectors are $(\lambda - \rho)$ eigenvectors for P .*

Proof. There are only two nontrivial cases where w has critical points

$$\begin{aligned} w &= C \cos\left(\sqrt{\bar{k}r}\right), \text{ or} \\ w &= C \cosh\left(\sqrt{-\bar{k}r}\right). \end{aligned}$$

By scaling we can further assume that $C = 1$ and additionally that $N^l \subset \{r = 0\}$.

We know that at N

$$\Delta r = \frac{n-l-1}{r} + O(1).$$

Thus

$$\begin{aligned} \Delta w &= (-\bar{k}r + O(r^3)) \left(\frac{n-l-1}{r} + O(1) \right) + (-\bar{k} + O(r^2)) \\ &= -\bar{k}(n-l) + O(r). \end{aligned}$$

The $(\lambda, n+m)$ -Einstein equation gives us

$$\begin{aligned} \Delta w &= \frac{w}{m} (\text{scal} - n\lambda) \\ &= \frac{w}{m} (\text{tr}P - n(\lambda - \rho)). \end{aligned}$$

So at N we have

$$\frac{1}{m} (\text{tr}P - n(\lambda - \rho)) = -\bar{k}(n-l)$$

showing that

$$\begin{aligned} \text{tr}P &= -(\lambda - \rho)(n-l) + n(\lambda - \rho) \\ &= l(\lambda - \rho). \end{aligned}$$

Corollary 3.5 implies that at N

$$P \circ (P - (\lambda - \rho)I) = 0.$$

Thus 0 and $\lambda - \rho$ are the only possible eigenvectors at N . Since $\frac{\nabla w}{|\nabla w|}$ converges to normal vectors to N it follows that any normal vector to N is a 0 eigenvector for P . As

$$\text{tr}P = l(\lambda - \rho)$$

the tangent vectors to N must be $\lambda - \rho$ eigenvectors for P . \square

Proposition 3.9. *Suppose a $(\lambda, n+m)$ -Einstein manifold (M, g, w) satisfies the following identity*

$$P \circ (P - (\lambda - \rho)I) = 0$$

everywhere on M , then for N , the set of critical points of w , we have

- (1) N is totally geodesic.
- (2) ∇P vanishes at N .

Proof. The equation

$$m\nabla\nabla w = w(P - (\lambda - \rho)I)$$

at N reduces to a soliton type equation

$$m\nabla\nabla w = P - (\lambda - \rho)I.$$

In addition we also have from Proposition 3.4 that at N

$$0 = -Q(X, Y, Z, \nabla w) = \frac{w}{m} ((\nabla_X P)(Y, Z) - (\nabla_Y P)(X, Z)).$$

First we show that N is totally geodesic. Let Y be a normal vector field to N and X be a tangent vector field. Thus

$$\begin{aligned} P(Y) &= 0 & \nabla_Y \nabla w &= \bar{k}Y \\ P(X) &= (\lambda - \rho)X & \nabla_X \nabla w &= 0. \end{aligned}$$

Using that the only eigenvalues for P are 0 and $\lambda - \rho$ we can extend X, Y such that they remain eigenfields for P . In particular

$$-P(\nabla_X Y) = (\nabla_X P)(Y) = (\nabla_Y P)(X) = (\lambda - \rho) \nabla_Y X - P(\nabla_Y X).$$

Then from the soliton equation we see that

$$\begin{aligned} (\lambda - \rho)g(\nabla_X Y, X) &= P(\nabla_X Y, X) - m\text{Hess}w(\nabla_X Y, X) \\ &= P(\nabla_X Y, X) \\ &= P(\nabla_Y X, X) - \lambda g(\nabla_Y X, X) + \rho g(\nabla_Y X, X) \\ &= m\text{Hess}w(\nabla_Y X, X) \\ &= 0. \end{aligned}$$

So the second fundamental form vanishes and N is totally geodesic.

Next we show that P is parallel at N . We show that

$$\begin{aligned} \nabla_X P &= 0, \\ \nabla_Y P &= 0, \end{aligned}$$

where X is tangent to N , and Y is normal to N .

To show the first we evaluate on $X' \in TN$ and Y' normal to N . Since

$$\begin{aligned} P(X') &= (\lambda - \rho)X' \\ P(Y') &= 0 \end{aligned}$$

we obtain

$$\begin{aligned} (\nabla_X P)(X') &= (\lambda - \rho) \nabla_X X' - P(\nabla_X X'), \\ (\nabla_X P)(Y') &= -P(\nabla_X Y'). \end{aligned}$$

Both of these expressions vanish as N is totally geodesic.

For the second case use Proposition 3.4 again to obtain:

$$\begin{aligned} \frac{w}{m}(\nabla_{\nabla w} P)(Z) &= -\left(\frac{w}{m}\right)^2 (((\lambda - \rho)I - P)P)(Z) + Q(Z, \nabla w) \nabla w \\ &= Q(Z, \nabla w) \nabla w. \end{aligned}$$

Dividing by $|\nabla w|$ then yields

$$\frac{w}{m}(\nabla_{\frac{\nabla w}{|\nabla w|}} P)(Z) = Q(Z, \nabla w) \frac{\nabla w}{|\nabla w|}$$

and supposing that

$$\frac{\nabla w}{|\nabla w|} \rightarrow Y$$

as we approach N we obtain

$$-\frac{w}{m} \nabla_Y P = 0$$

as desired. \square

Remark 3.10. If a $(\lambda, n+m)$ -Einstein manifold satisfies the radial Q flatness condition (1.2), then we will show that the P tensor satisfies the equation in Proposition 3.9.

4. PROOF OF THEOREM 1.2, 1.7, AND 1.8

In this section we will discuss the proofs of theorems 1.2, 1.7, and 1.8. The easiest to prove is Theorem 1.7 which is the classification in the $\lambda = 0$ case. It already follows directly from the formulas in the past section, and is the same argument as in [CSW].

Proof of Theorem 1.7. Since the scalar curvature is constant and $\lambda = 0$, from Proposition 3.1 we have

$$\begin{aligned}\rho &= -\frac{1}{n+m-1}\mathrm{tr}(P), \\ |P|^2 &= -\rho\mathrm{tr}(P) = \frac{1}{n+m-1}(\mathrm{tr}P)^2.\end{aligned}$$

The second identity above and the Cauchy-Schwarz inequality $n|P|^2 \geq (\mathrm{tr}(P))^2$ imply that

$$\frac{n}{n+m-1}(\mathrm{tr}(P))^2 \geq (\mathrm{tr}(P))^2.$$

So we have $\mathrm{tr}(P) = 0$ as $m > 1$. It follows that $\rho = 0$ and then $|P|^2 = 0$. Hence $\mathrm{Ric} = \rho g$ and the result follows from the classification in Proposition 2.3. \square

Next we turn our attention to Theorem 1.8. Before the proof we give two corollaries that follow from combining the theorem with some of the other results we have already discussed.

Corollary 4.1. *Let (M, g, w) be a $(\lambda, n+m)$ -Einstein manifold with constant scalar curvature and $m > 1$. If $\lambda > 0$, then*

$$0 \leq \rho \leq \lambda,$$

and if $\lambda < 0$, then

$$0 \geq \rho \geq \lambda.$$

Moreover, in either case, $\rho = \lambda$ if and only if $\lambda = 0$ and the metric is rigid, and $\rho = 0$ if and only if $M = N \times \mathbb{R}$ where N is a λ -Einstein metric.

Proof. In Corollary 3.3 we already saw that $\lambda \geq \rho$ when $\lambda > 0$ and that $\lambda \leq \rho$ when $\lambda < 0$. To see the rigidity statement for this side of inequality, note that if $\lambda = \rho$, then $\bar{k} = 0$ and then we get rigidity by combining Proposition 3.6 and Theorem 1.7.

The other inequality is equivalent to Theorem 1.8. To see this note that, since $\mathrm{Ric}(\nabla w) = \rho \nabla w$, the hypothesis on $\mathrm{Ric}(\nabla w, \nabla w)$ is equivalent to assuming ρ is zero or has the opposite sign of λ . \square

As mentioned in the introduction, Theorem 1.8 can also be interpreted as a gap theorem about the scalar curvature.

Corollary 4.2. *Let (M, g, w) be a $(\lambda, n+m)$ -Einstein manifold with constant scalar curvature and $m > 1$. If the scalar curvature is between $(n-1)\lambda$ and $n\lambda$ then either the metric is a λ -Einstein metric, or it is rigid and splits as $N \times \mathbb{R}$ where N is a λ -Einstein metric.*

Proof. As we have seen in Proposition 3.2, the scalar curvature must be between $n\rho$ and $n\lambda$, and can only be equal to $n\lambda$ if the metric is λ -Einstein. On the other hand, by the definition of ρ , the scalar curvature being bounded away from zero by $(n-1)\lambda$ is equivalent to ρ having the opposite sign as λ , so the other half of the result is equivalent to Theorem 1.8. \square

Now we prove the theorem.

Proof of Theorem 1.8. For the discussion above we see that the hypothesis is equivalent to $\rho\bar{k} \leq 0$.

By Proposition 3.6 we see that if w is not constant, then $w = w(r)$ where r is a distance function and $w'' = -\bar{k}w$. This implies

$$\begin{aligned}\nabla w &= w'\nabla r \\ \Delta w &= -\bar{k}w + w'\Delta r\end{aligned}$$

and

$$\Delta w = \frac{w}{m}(\text{scal} - n\lambda)$$

so

$$-\bar{k}w + w'\Delta r = \frac{w}{m}(\text{scal} - n\lambda)$$

which implies that

$$w'\Delta r = w\left(\frac{\text{scal} - n\lambda}{m} + \frac{\lambda - \rho}{m}\right) = -\rho w.$$

The Bochner formula for r is

$$\begin{aligned}0 &= |\text{Hess}r|^2 + g(\nabla r, \nabla \Delta r) + \text{Ric}(\nabla r, \nabla r) \\ &= |\text{Hess}r|^2 + g(\nabla r, \nabla \Delta r) + \rho.\end{aligned}$$

The middle term can be calculated by using

$$\begin{aligned}-\rho w'\nabla r &= -\bar{k}w\Delta r\nabla r + w'\nabla \Delta r \\ &= \frac{-\bar{k}w}{w'}(-\rho w)\nabla r + w'\nabla \Delta r \\ &= \rho\frac{\bar{k}w^2}{w'}\nabla r + w'\nabla \Delta r\end{aligned}$$

showing that

$$g(\nabla r, \nabla \Delta r) = -\rho + \rho\bar{k}\frac{w^2}{(w')^2}.$$

Thus we obtain

$$\begin{aligned}0 &= |\text{Hess}r|^2 + g(\nabla r, \nabla \Delta r) + \rho \\ &= |\text{Hess}r|^2 - \rho\bar{k}\frac{w^2}{(w')^2}\end{aligned}$$

showing that $\rho\bar{k} \geq 0$, and can only vanish when $\text{Hess}r = 0$, which implies the splitting along the gradient of w . \square

We finish this section by showing Theorem 1.2, i.e., a three dimensional $(\lambda, 3+m)$ -Einstein manifold is rigid if it has constant scalar curvature.

Proof of Theorem 1.2. We start by showing that P has constant eigenvalues. Proposition 3.1 says that

$$\begin{aligned} P(\nabla w) &= 0, \\ \operatorname{tr} P &= 2\lambda - (m+2)\rho, \\ |P|^2 &= (\lambda - \rho)\operatorname{tr} P. \end{aligned}$$

Thus one eigenvalue is 0 and if the other two are p_1 and p_2 then

$$p_{1,2} = \frac{1}{2} \left(2\lambda - (m+2)\rho \pm \sqrt{2m\rho\lambda - 2m\rho^2 - m^2\rho^2} \right) = \frac{1}{2} \left(\operatorname{tr} P \pm \sqrt{m\rho(\operatorname{tr} P)} \right).$$

This shows in particular that $(\nabla P)(E, E) = 0$ if E is a unit eigenvector for P .

The goal now is to prove that either $\lambda = 0$ in which case we can use Theorem 1.7 or $p_1 = p_2$. In the latter case the metric is either ρ -Einstein or $\rho = 0$ reducing us respectively to Proposition 2.3 or Corollary 4.1.

Recall

$$Q = R + \frac{2}{m}P \odot g + \frac{\rho - \lambda}{m}g \odot g.$$

If $n = 3$, then we have $R = \frac{\operatorname{scal}}{6}g \odot g + (\operatorname{Ric} - \frac{2\operatorname{scal}}{3}g) \odot g$. Since $\operatorname{scal} = \operatorname{tr} P + 3\rho$, the tensor Q can be written as

$$Q = \frac{m+1}{m} \left(2P \odot g - \frac{1}{2}(\operatorname{tr} P)g \odot g \right).$$

If E_i is a unit eigenfield for p_i , then

$$\begin{aligned} Q(\nabla w, E_i, E_i, \nabla w) &= \frac{m+1}{m} \left(|\nabla w|^2 P(E_i, E_i) - \frac{\operatorname{tr} P}{2} |\nabla w|^2 g(E_i, E_i) \right) \\ &= \frac{m+1}{2m} (p_i - p_j) |\nabla w|^2 \end{aligned}$$

where $j \neq i$.

Using Proposition 3.4 we have

$$\begin{aligned} \frac{w}{m} (\nabla_{\nabla w} P)(X, Y) &= - \left(\frac{w}{m} \right)^2 (\lambda - \rho) P(X, Y) + \left(\frac{w}{m} \right)^2 g(P(X), P(Y)) \\ &\quad + Q(\nabla w, X, Y, \nabla w). \end{aligned}$$

Evaluating on E_i yields

$$0 = - \left(\frac{w}{m} \right)^2 (\lambda - \rho) p_i + \left(\frac{w}{m} \right)^2 p_i^2 + \frac{m+1}{2m} (p_i - p_j) |\nabla w|^2.$$

When M has boundary we know that $w = 0$ somewhere so this formula immediately shows that $p_1 = p_2$. In general we can subtract the two equations to obtain

$$0 = (p_1 - p_2) \left(- \frac{w^2}{m} (\lambda - \rho) + \frac{w^2}{m} (p_1 + p_2) + (m+1) |\nabla w|^2 \right).$$

This shows that either $p_1 = p_2$ or

$$2\lambda - (m+2)\rho = \operatorname{tr} P = (\lambda - \rho) - m(m+1) \frac{|\nabla w|^2}{w^2}.$$

As

$$\frac{\lambda - \rho}{m} w^2 + |\nabla w|^2 = \bar{\mu}$$

the latter case can only happen when $\bar{\mu} = 0$ and

$$\frac{-(\lambda - \rho) + m\rho}{m + 1} = m \frac{|\nabla w|^2}{w^2} = -(\lambda - \rho).$$

Thus $\lambda = 0$. □

Remark 4.3. In the proof above, we showed that Ric has constant eigenvalues. Such a metric is called *Ricci curvature homogeneous*. This condition is more general than being *curvature homogeneous*, i.e., for any two points p and q in M , there exists a linear isometry $\phi : T_p M \rightarrow T_q M$ such that $\phi^*(R_q) = R_p$, see [Si]. Both notions are equivalent in the two and three-dimensional cases but not for higher dimensions. In dimension three there are curvature homogeneous spaces that are not homogeneous, see [Bu, BV, Ko] and the references therein.

5. PROOF OF THEOREM 1.10 AND COROLLARY 1.12

In this section we prove Theorem 1.10, i.e., that radial flatness of Q implies the rigidity. There are a number of steps and the proof breaks down into different cases. The proof comes from studying the eigenvalues and eigen-distributions of P . First we show that 0 and $\lambda - \rho$ are the only possible eigenvalues.

Lemma 5.1. *Suppose a $(\lambda, n + m)$ -Einstein manifold (M, g, w) has constant scalar curvature and*

$$Q(\nabla w, \cdot, \cdot, \nabla w) = 0.$$

Then the eigenvalues of P are either 0 or $\lambda - \rho$, i.e., on M we have

$$P \circ (P - (\lambda - \rho)I) = 0.$$

Proof. Since the scalar curvature is constant, from Proposition 3.4 we have

$$\begin{aligned} (\nabla_{\nabla w} P) &= \frac{w}{m} P \circ (P - (\lambda - \rho)I) \\ &= \frac{w}{m} P \circ (P - m\bar{k}I), \end{aligned}$$

where $\bar{k} = \frac{\lambda - \rho}{m} \neq 0$ by Corollary 3.3.

In addition from Proposition 3.6 if $\bar{k} > 0$ then

$$w = \cos(\sqrt{\bar{k}r}),$$

and if $\bar{k} < 0$

$$\begin{aligned} w &= \exp(\sqrt{-\bar{k}r}), \\ w &= \cosh(\sqrt{-\bar{k}r}), \quad \text{or} \\ w &= \sinh(\sqrt{-\bar{k}r}). \end{aligned}$$

At points where $\nabla w \neq 0$ the radial curvature equation can be rewritten as

$$(\nabla_{\nabla r} P) = \frac{w}{mw'} (P \circ (P - m\bar{k}I)).$$

Let Y be a unit parallel vector field along an integral curve of r and let $y(r) = P(Y, Y)$. So the above equation becomes

$$y'(r) = \frac{w}{mw'}y(y - m\bar{k}).$$

Using that $w'' = -\bar{k}w$ we see that this equation has the general nonzero solution

$$\begin{aligned} y(r) &= m\bar{k} \frac{w'}{w' + A} \\ &= \frac{m\bar{k}}{1 + A(w')^{-1}}. \end{aligned}$$

Choose an orthonormal frame $\{Y_i\}_1^{n-1}$ in the normal space of ∇r and assume that they are parallel vector fields along ∇r . Suppose $y_i = P(Y_i, Y_i)$, $i = 1, 2, \dots, p$, are the non-constant values, i.e.,

$$y_i = \frac{m\bar{k}}{1 + A_i z(r)},$$

where $z(r) = (w')^{-1}$ and A_i 's are nonzero constants. As $\text{tr}(P)$ is constant, it follows that

$$\sum_{i=1}^p y_i = 0.$$

To see the constant on the right hand side is zero, let $r \rightarrow 0$. Now plugging in the specific form for y we obtain

$$\frac{p + b_1 z + b_2 z^2 + \dots + b_p z^p}{\prod_{i=1}^p (1 + A_i z)} = 0,$$

where the coefficients b_i can be derived from the binomial formula. However, unless $p = 0$ this is a contradiction since $z(r)$ has the property that $\{1, z, z^2, z^3, \dots\}$ is a linearly independent set of functions on their intervals of definition.

Therefore, in all cases, $P(Y_i, Y_i)$ is constant and thus either 0 or $\lambda - \rho$. It follows that the eigenvalues of P are bounded by 0 and $\lambda - \rho$. Using the identity $|P|^2 = (\lambda - \rho)\text{tr}P$ in Proposition 3.1, it is either 0 or $\lambda - \rho$. \square

From Proposition 3.6, there are four different cases:

- (1) $w = \cos(\sqrt{\bar{k}r})$, M has boundary and $\bar{\mu} > 0$,
- (2) $w = \cosh(\sqrt{-\bar{k}r})$, M has no boundary and $\bar{\mu} < 0$,
- (3) $w = \sinh(\sqrt{-\bar{k}r})$, M has boundary and $\bar{\mu} > 0$,
- (4) $w = \exp(\sqrt{-\bar{k}r})$, M has no boundary and $\bar{\mu} = 0$.

In the first two cases, the critical point set of w is non-empty and the proof follows from a similar argument as in Ricci soliton case, see [PW3].

Proof of Theorem 1.10 in Case (1) and (2). This comes from considering the critical point set of w , $N = \{x : \nabla w(x) = 0\}$. From Proposition 3.9 we see that N is a totally geodesic λ -Einstein manifold.

The normal exponential map $\nu(N) \rightarrow M$ follows the integral curves for ∇w (or ∇r) and therefore is a diffeomorphism. Using the fundamental equations, see for example [Pe, Chapter 2.5], the metric on M is completely determined by the radial

sectional curvatures and the metric on N as $N \subset M$ is totally geodesic. Since these match exactly with the values in the corresponding rigid cases, the metric must be rigid. \square

Now we are left with Case (3) and (4). The proof of the theorem in these cases is much more involved since $N = \emptyset$.

From the explicit formula of the function w the maximal interval I of r is either $(-\infty, \infty)$ or $[0, \infty)$. On this interval we can write the metric as

$$\begin{aligned} M &= I \times \Sigma \\ g &= dr^2 + g_r. \end{aligned}$$

From Lemma 5.1, the tangent space of Σ has an orthogonal splitting of eigen-distributions of P :

$$\begin{aligned} T\Sigma &= \mathcal{N} \oplus \mathcal{P} \\ P|_{\mathcal{N}} &= (\lambda - \rho)\text{id}_{\mathcal{N}} \\ P|_{\mathcal{P}} &= 0, \end{aligned}$$

and they are parallel along ∇r .

Theorem 1.10 will follow by showing that the distributions are parallel, see Lemma 5.3 for Case (3) and Lemma 5.4 for Case (4). In Lemma 5.4 we assume that the curvature does not grow as fast as exponentially. In fact we can show that the eigen-distribution of P is integrable without this curvature growth assumption, see Proposition A.2 and Theorem A.1 in Appendix A.

Before proceeding to the general case we note that these steps are considerably simpler in the harmonic curvature case. In fact, the argument in [Be, Proposition 16.11] shows that the eigen-distributions of P are always integrable in this case. Moreover, if the curvature is harmonic, then

$$(\nabla_{E_1} \text{Ric})(E_2, E_3) - (\nabla_{E_2} \text{Ric})(E_1, E_3) = \text{div}R(E_1, E_2, E_3) = 0$$

for any vector fields E_1, E_2 and E_3 . The scalar curvature is also constant and thus Proposition 3.4 shows that $Q(E_1, E_2, E_3, \nabla w) = 0$. This explains why harmonic curvature is a stronger assumption than $Q(\nabla w, E, F, \nabla w) = 0$. For any two eigen-vector fields X with $P(X) = (\lambda - \rho)(X)$ and U with $P(U) = 0$ the vanishing of Q also implies that

$$R(X, U, X, \nabla r) = R(U, X, U, \nabla r) = 0.$$

On the other hand, we also have the following calculation. We will only use the first and third equations for the harmonic curvature case, but will find the other two equations useful later.

Proposition 5.2. *Suppose X, Y and U, V are $\lambda - \rho$ and 0 eigenvector fields of P respectively. Then we have*

$$(5.1) \quad R(X, U, Y, \nabla r) = -\bar{k} \frac{w}{w'} g(\nabla_X Y, U)$$

$$(5.2) \quad (\nabla_{\nabla r} R)(X, U, Y, \nabla r) = \frac{w'}{w} R(X, U, Y, \nabla r) + \bar{k} \frac{w}{w'} (R(X, U, Y, \nabla r) + R(Y, U, X, \nabla r))$$

$$(5.3) \quad R(U, X, V, \nabla r) = \bar{k} \frac{w}{w'} g(\nabla_U V, X)$$

$$(5.4) \quad (\nabla_{\nabla r} R)(U, X, V, \nabla r) = \left(\frac{w'}{w} + \bar{k} \frac{w}{w'} \right) R(U, X, V, \nabla r) + \bar{k} \frac{w}{w'} R(U, V, X, \nabla r).$$

Proof. First note that

$$\text{Hess}w = -\bar{k}wdr \otimes dr + w'\text{Hess}r.$$

Therefore the second fundamental form for the hypersurface $w^{-1}(r)$ is

$$\text{II} = \text{Hess}r = (w')^{-1} \text{Hess}w = \frac{w}{mw'} (\text{Ric} - \lambda g),$$

i.e.,

$$\begin{aligned} \text{II}(X, X) &= 0 \\ \text{II}(X, U) &= 0 \\ \text{II}(U, U) &= -\bar{k} \frac{w}{w'} |U|^2. \end{aligned}$$

This implies

$$\begin{aligned} R(X, \nabla r) \nabla r &= 0 \\ R(U, \nabla r) \nabla r &= \bar{k}U, \end{aligned}$$

and

$$\begin{aligned} R(X, U, Y, \nabla r) &= (\nabla_U \text{II})(X, Y) - (\nabla_X \text{II})(U, Y) \\ &= \text{II}(U, \nabla_X Y) \\ &= -\bar{k} \frac{w}{w'} g(\nabla_X Y, U). \end{aligned}$$

From the second Bianchi identity

$$(\nabla_{\nabla r} R)(X, U, Y, \nabla r) + (\nabla_X R)(U, \nabla r, Y, \nabla r) + (\nabla_U R)(\nabla r, X, Y, \nabla r) = 0,$$

it follows then that

$$\begin{aligned} (\nabla_{\nabla r} R)(X, U, Y, \nabla r) &= (\nabla_X R)(\nabla r, U, Y, \nabla r) - (\nabla_U R)(\nabla r, X, Y, \nabla r) \\ &= -R(\nabla r, U, \nabla_X Y, \nabla r) + R(\nabla_U \nabla r, X, Y, \nabla r) + R(\nabla r, X, Y, \nabla_U \nabla r) \\ &= \frac{1}{m} (\text{Ric}(U, \nabla_X Y) - \lambda g(U, \nabla_X Y)) - \bar{k} \frac{w}{w'} R(U, X, Y, \nabla r) - \bar{k} \frac{w}{w'} R(\nabla r, X, Y, U) \\ &= -\bar{k} g(U, \nabla_X Y) + \bar{k} \frac{w}{w'} (R(X, U, Y, \nabla r) + R(Y, U, X, \nabla r)) \\ &= \frac{w'}{w} R(X, U, Y, \nabla r) + \bar{k} \frac{w}{w'} (R(X, U, Y, \nabla r) + R(Y, U, X, \nabla r)). \end{aligned}$$

An analogous argument gives us the other equations:

$$\begin{aligned}
R(U, X, V, \nabla r) &= (\nabla_X \text{II})(U, V) - (\nabla_U \text{II})(X, V) \\
&= \text{II}(\nabla_U X, V) \\
&= -\bar{k} \frac{w}{w'} g(\nabla_U X, V) \\
&= \bar{k} \frac{w}{w'} g(X, \nabla_U V)
\end{aligned}$$

and

$$\begin{aligned}
(\nabla_{\nabla r} R)(U, X, V, \nabla r) &= (\nabla_U R)(\nabla r, X, V, \nabla r) - (\nabla_X R)(\nabla r, U, V, \nabla r) \\
&= -R(\nabla r, \nabla_U X, V, \nabla r) - R(\nabla_U \nabla r, X, V, \nabla r) - R(\nabla r, X, V, \nabla_U \nabla r) \\
&= \frac{1}{m} (\text{Ric}(V, \nabla_U X) - \lambda g(V, \nabla_U X)) + \bar{k} \frac{w}{w'} R(U, X, V, \nabla r) + \bar{k} \frac{w}{w'} R(\nabla r, X, V, U) \\
&= -\bar{k} g(V, \nabla_U X) + \bar{k} \frac{w}{w'} R(U, X, V, \nabla r) + \bar{k} \frac{w}{w'} R(U, V, X, \nabla r) \\
&= \bar{k} g(\nabla_U V, X) + \bar{k} \frac{w}{w'} R(U, X, V, \nabla r) + \bar{k} \frac{w}{w'} R(U, V, X, \nabla r) \\
&= \left(\frac{w'}{w} + \bar{k} \frac{w}{w'} \right) R(U, X, V, \nabla r) + \bar{k} \frac{w}{w'} R(U, V, X, \nabla r) \\
&= \frac{w'}{w} R(U, X, V, \nabla r) + \bar{k} \frac{w}{w'} (R(U, X, V, \nabla r) - R(V, X, U, \nabla r)).
\end{aligned}$$

These give us the desired equations. \square

We can now finish the proof when the curvature is harmonic.

Proof of Corollary 1.12. As the curvature is harmonic we have seen above that $R(X, U, X, \nabla r) = 0$ and $R(U, X, U, \nabla r) = 0$. When combined with the previous proposition, this tells us that the eigen-distributions are totally geodesic. This gives a splitting of the universal cover along the eigen-distributions of P . The results from section 2 tell us the metric is rigid. \square

Lemma 5.3. *If M has non-empty boundary then the two eigen-distributions of P are parallel.*

Proof. As $\nabla_{\nabla r} P = 0$, we can assume that the eigenvector fields $X, Y, \dots \in \mathcal{N}$ and $U, V, \dots \in \mathcal{P}$ are parallel along ∇r . The fact that the boundary is nonempty is used as an initial value for the curvatures. We assume that the boundary corresponds to the level set $r = 0$. Specifically we see that Proposition 5.2 implies that $R(E_1, E_2, E_3, \nabla r) = 0$ on the boundary. If we set $X = Y$ in (5.1) and (5.2), then we obtain

$$\begin{aligned}
(\nabla_{\nabla r} R)(X, U, X, \nabla r) &= \left(\frac{w'}{w} + 2\bar{k} \frac{w}{w'} \right) R(X, U, X, \nabla r) \\
R(X, U, X, \nabla r) &= 0 \quad \text{at } r = 0.
\end{aligned}$$

So it follows that $R(X, U, X, \nabla r) = 0$ on M . This shows that $R(X, U, Y, \nabla r)$ is skew-symmetric in X, Y and hence that

$$(\nabla_{\nabla r} R)(X, U, Y, \nabla r) = \frac{w'}{w} R(X, U, Y, \nabla r).$$

We can then similarly conclude that $R(X, U, Y, \nabla r)$ vanishes as long as it vanishes on the boundary. Using (5.1) again this shows that \mathcal{N} is integrable as well as totally geodesic.

A similar argument works for \mathcal{P} . Setting $U = V$ in (5.4) we have

$$(\nabla_{\nabla r} R)(U, X, U, \nabla r) = \left(\frac{w'}{w} + \bar{k} \frac{w}{w'} \right) R(U, X, U, \nabla r).$$

So $R(U, X, U, \nabla r)$ vanishes as it vanishes on the boundary. This implies that $R(U, X, V, \nabla r)$ is skew-symmetric in U and V , which in turn shows that

$$\begin{aligned} R(U, V, X, \nabla r) &= R(U, X, V, \nabla r) - R(V, X, U, \nabla r) \\ &= 2R(U, X, V, \nabla r) \end{aligned}$$

and consequently

$$(\nabla_{\nabla r} R)(U, X, V, \nabla r) = \left(\frac{w'}{w} + 3\bar{k} \frac{w}{w'} \right) R(U, X, V, \nabla r).$$

As $R(U, X, V, \nabla r)$ vanishes on the boundary, it vanishes everywhere showing that \mathcal{P} is totally geodesic. \square

Lemma 5.4. *If M has no boundary, i.e., $\bar{\mu} = 0$, and we further assume that $|R| = o\left(\exp\left(\text{dist}(x, p)\sqrt{-\bar{k}}\right)\right)$, then the two eigenvalue distributions for P are parallel.*

Proof. We now turn to Case (4) where $w = \exp\left(\sqrt{-\bar{k}r}\right)$. This means that equation (5.2) reduces to

$$D_{\nabla r} R(X, U, Y, \nabla r) = \sqrt{-\bar{k}} R(X, U, Y, \nabla r) - \sqrt{-\bar{k}} (R(X, U, Y, \nabla r) + R(Y, U, X, \nabla r)),$$

i.e.,

$$R(X, U, X, \nabla r) = R(X, U, X, \nabla r)|_{r=0} \exp\left(-\sqrt{-\bar{k}r}\right).$$

In particular we see again that $R(X, U, X, \nabla r)$ must vanish if we assume that it cannot grow as fast as $\exp\left(\sqrt{-\bar{k}r}\right)$. This again shows that $R(X, U, Y, \nabla r)$ is skew-symmetric in X, Y and thus

$$\begin{aligned} D_{\nabla r} R(X, U, Y, \nabla r) &= \sqrt{-\bar{k}} R(X, U, Y, \nabla r) - \sqrt{-\bar{k}} (R(X, U, Y, \nabla r) + R(Y, U, X, \nabla r)) \\ &= \sqrt{-\bar{k}} R(X, U, Y, \nabla r). \end{aligned}$$

Using the growth assumption for $R(X, U, Y, \nabla r)$ this in turn shows that $R(X, U, Y, \nabla r)$ vanishes and then that \mathcal{N} is totally geodesic.

Next we note that $R(U, X, V, \nabla r) = \bar{k} \frac{w}{w'} g(\nabla_U V, X)$. The right hand side can be calculated using Koszul's formula together with the metric decomposition

$$\begin{aligned} g &= dr^2 + g_r \\ &= dr^2 + h_0 + \frac{(w'(r))^2}{(w'(0))^2} h_1, \end{aligned}$$

where h_0 and h_1 are the restrictions of g_0 to \mathcal{N} and \mathcal{P} at $r = 0$ respectively.

We wish to calculate $g(\nabla_U V, X)$ and relate it to what happens at $r = 0$. This requires that the fields we use commute with ∇r . As our fields are chosen to be parallel along ∇r we simply have to switch to $X, w'U, w'V$ instead. After eliminating $(w')^2$ on both sides this yields the formula

$$\begin{aligned} g(\nabla_U V, X) &= g_r(\nabla_U V, X) \\ &= g_0(\nabla_U V, X) + \left(\frac{1}{2} - \frac{1}{2(w'(r))^2} \right) g_0([V, U], X). \end{aligned}$$

Thus

$$R(U, X, V, \nabla r) = \bar{k} \frac{w}{w'} \left(g_0(\nabla_U V, X) + \left(\frac{1}{2} - \frac{1}{2(w'(r))^2} \right) g_0([V, U], X) \right)$$

is forced to grow like $(w')^{-2} = |\bar{k}| \exp(-2r\sqrt{-\bar{k}})$ unless $[U, V]$ is perpendicular to X . As we have assumed that the curvature grows slower than $\exp(\sqrt{-\bar{k}}r)$ this shows that \mathcal{P} is an integrable distribution.

Having shown that \mathcal{N} is totally geodesic and its orthogonal distribution $\{\nabla r\} \oplus \mathcal{P}$ is integrable it follows that the foliation with vertical space given by $\{\nabla r\} \oplus \mathcal{P}$ is Riemannian. This means that we can use [BH] to conclude that there is a map $F : B \times H \rightarrow M$ such that $F(B \times \{q\})$ is an integral manifold for \mathcal{N} for all $q \in H$ and similarly $F(\{p\} \times H)$ is an integral manifold for $\{\nabla r\} \oplus \mathcal{P}$ for all $p \in B$. Below we shall show that the fibers are all isometric to each other and are in fact a simply connected hyperbolic space of constant curvature \bar{k} .

Note that if X is chosen to be basic along this Riemannian foliation then $[X, U] \in \mathcal{P}$ for any $U \in \mathcal{P}$. As \mathcal{N} is totally geodesic we have $\nabla_X U \in \mathcal{P}$, consequently also $\nabla_U X \in \mathcal{P}$. Let $U, V \in \mathcal{P}$ with $g(U, V) = 0$, then

$$\begin{aligned} D_{\nabla r} R(V, U, U, V) &= (\nabla_V R)(V, U, U, \nabla r) - (\nabla_U R)(V, U, V, \nabla r) \\ &= -R(\nabla_V V, U, U, \nabla r) - R(V, \nabla_V U, U, \nabla r) \\ &\quad -R(V, U, \nabla_V U, \nabla r) - R(V, U, U, \nabla_V \nabla r) \\ &\quad +R(\nabla_U V, U, V, \nabla r) + R(V, \nabla_U U, V, \nabla r) \\ &\quad +R(V, U, \nabla_U V, \nabla r) + R(V, U, V, \nabla_U \nabla r) \\ &= -\sqrt{-\bar{k}}g(\mathcal{N}(\nabla_V V), \nabla_U U) + R(\nabla r, U, U, \nabla r)\text{II}(V, V) \\ &\quad +\sqrt{-\bar{k}}g(\mathcal{N}(\nabla_V U), \nabla_V U) + \sqrt{-\bar{k}}g(\mathcal{N}(\nabla_U V), \nabla_U V) \\ &\quad -\sqrt{-\bar{k}}g(\mathcal{N}(\nabla_U U), \nabla_V V) - R(V, \nabla r, V, \nabla r)\text{II}(U, U) \\ &\quad -2\sqrt{-\bar{k}}R(V, U, U, V) \\ &= -2\sqrt{-\bar{k}}g(\mathcal{N}(\nabla_V V), \mathcal{N}(\nabla_U U)) + 2\sqrt{-\bar{k}}|\mathcal{N}(\nabla_V U)|^2 \\ &\quad -2\sqrt{-\bar{k}}R(V, U, U, V) + 2\bar{k}\sqrt{-\bar{k}}g(U, U)g(V, V). \end{aligned}$$

Here the term from the second fundamental form

$$(5.5) \quad g(\mathcal{N}(\nabla_V V), \mathcal{N}(\nabla_U U)) - |\mathcal{N}(\nabla_V U)|^2 - \bar{k}g(U, U)g(V, V)$$

is constant in the ∇r direction. Thus $R(V, U, U, V)$ will grow exponentially unless it is constant and precisely cancels the term in (5.5), i.e.,

$$R(V, U, U, V) = |\mathcal{N}(\nabla_V U)|^2 - g(\mathcal{N}(\nabla_V V), \mathcal{N}(\nabla_U U)) + \bar{k}g(U, U)g(V, V).$$

This indicates that \mathcal{P} is flat in M and more importantly that the fibers are isometric to the simply-connected hyperbolic space with curvature \bar{k} which we denote by (H, h) .

We then conclude that the structure $F : B \times H \rightarrow M$ gives us a metric decomposition of the form

$$F^*(g) = h_0 + f_b^*(h)$$

where $f_b : H \rightarrow H$ is a family of isometries parametrized by the base space B . From this it follows that any isometry of (H, h) extends to an isometry of M that fixes the horizontal space B and maps a fiber to itself. Recall that we have a distance function r such that ∇r is tangent to the fibers. This means that a geodesic γ with velocity vector ∇r stays in the fiber. Using an isometry $f \in \text{Iso}(H, h)$ then yields a new geodesic $f \circ \gamma$ with velocity $Df(\nabla r)$ that also stays in the fiber. As the isometry group of hyperbolic space is isotropic, i.e., the isotropy action is transitive on the unit sphere in the tangent space, it follows that any geodesic tangent to a fiber will stay in that fiber, i.e., the fiber is totally geodesic. \square

Now we finish the proof of Theorem 1.10.

Proof of Theorem 1.10. Suppose (M, g, w) is a non-trivial $(\lambda, n+m)$ -Einstein manifold. By the definition of Q , the radial section curvature condition (1.2) is equivalent to $Q(\nabla w, \cdot, \cdot, \nabla w) = 0$. From Lemma 5.1, the eigenvalues of P are either 0 or $\lambda - \rho$. From Lemma 5.3 and 5.4, the distribution for each eigenvalue is parallel. Hence the manifold splits as $(M, g) = (M_1 \times M_2, g_1 + g_2)$ such that $\text{Ric}_{g_1} = \lambda g_1$ and $\text{Ric}_{g_2} = \rho g_2$. So the rigidity of M follows from Proposition 2.5. \square

Remark 5.5. We have only used the assumption about exponential growth of curvature in the special case when $\bar{\mu} = 0$ and the potential function is $w = \exp(\sqrt{-k}r)$. Moreover, from the proof of Lemma 5.4, we can see that in this case we get a stronger conclusion than rigidity. Namely that the manifold M splits isometrically as $L \times H$ where L is λ -Einstein and H is the simply-connected hyperbolic space with Ricci curvature ρ .

6. NON-RIGID $(\lambda, n+m)$ -EINSTEIN METRICS ON SOLVABLE LIE GROUPS

In this section, we consider $(\lambda, n+m)$ -Einstein metrics on homogeneous spaces. The general case of the $(\lambda, n+m)$ -Einstein equation is studied in [HPW2] where the function w is also allowed to take negative values. Using this generalized equation, we obtain

Theorem 6.1. *If (M, g) be a simply connected homogeneous non-trivial $(\lambda, m+n)$ -Einstein manifold with $m > 1$ and $\lambda < 0$, then one of the following cases holds.*

- (1) $\mu < 0$ and $M = S \times L$ as a Riemannian product where S is a space form that is ρ -Einstein and L is λ -Einstein.
- (2) $\mu = 0$ and there are no other solutions to the λ -Einstein equations with $\mu < 0$, then the isometry group $\text{Iso}(M)$ contains a codimension one normal subgroup H that acts transitively on the connected components of the level set of w . Moreover if F is an isometry of M , then $w \circ F = Cw$ for some constant C .

Remark 6.2. In Case (1) above when $\mu < 0$, one can show that M is rigid by Proposition 3.6, Proposition 3.8 and Proposition 3.9. In fact, $\mu < 0$ implies that w has critical points. At a critical point $\nabla \text{Ric} = 0$ and the eigenvalue of Ric is either λ or ρ . Since the metric is homogeneous, this holds everywhere on M . So the metric splits along the eigen-distributions of Ric , i.e., it is rigid.

Theorem 6.1 suggests that we are most likely to find non-rigid examples on solvable Lie groups. Using left invariant vector fields, we can rewrite the $(\lambda, n+m)$ -Einstein equation.

Lemma 6.3. *Let (G, g) be a simply-connected Lie group with left invariant metric. Suppose it admits a non-trivial $(\lambda, n+m)$ -Einstein structure with $m > 1$ and $\mu = 0$. Then there exists a codimension one normal subgroup $H \subset G$ and left invariant vector field $X_0 = \nabla r$, where r is the signed distance function from H . Furthermore the $(\lambda, n+m)$ -Einstein equation can be written as*

$$(6.1) \quad \text{Ric}(X, Y) - \frac{1}{2} \sqrt{m(\rho - \lambda)} (g([X_0, X], Y) + g([X_0, Y], X)) - (\rho - \lambda)g(X_0, X)g(X_0, Y) = \lambda g(X, Y),$$

for any two left invariant vector fields X and Y .

Proof. From Proposition 3.6 and Theorem 6.1 we may assume that $w = e^{\sqrt{-k}r}$ and the level hypersurface of $w = 1$ is a codimension one normal subgroup H in G . Thus we obtain a Riemannian submersion $G \rightarrow G/H$ which is also a Lie algebra homomorphism. This shows that left invariant vector fields on G/H lift to left invariant vector fields on G that are perpendicular to H . As $G/H = \mathbb{R}$ it follows that ∇r is a left invariant vector field on G .

Using Koszul's formula we note that

$$\text{Hess}r(X, Y) = g(\nabla_X X_0, Y) = \frac{1}{2} (g([X_0, X], Y) + g([X_0, Y], X))$$

when X, Y are left invariant. As

$$\text{Hess}w = \sqrt{-k}w \text{Hess}r - \bar{k}w dr^2$$

we obtain the desired form for the $(\lambda, n+m)$ -Einstein equation (1.1). \square

Proposition 6.4. *Let G be a unimodular Lie group with left invariant metric g . If (G, g) is a nontrivial $(\lambda, n+m)$ -Einstein manifold with $m > 1$, $\lambda < 0$ and $\mu = 0$, then $G = H \times \mathbb{R}$ where H is λ -Einstein.*

Proof. Choose an orthonormal basis $\{X_i\}_{i=0}^{n-1}$ of left invariant vector fields. Then from the equation (6.1) we have

$$\text{Ric}(X_i, X_i) = \sqrt{m(\rho - \lambda)}g([X_0, X_i], X_i) + (\rho - \lambda)\delta_{0i} + \lambda$$

which implies that

$$\text{scal} = \sqrt{m(\rho - \lambda)}\text{tr}(\text{ad}_{X_0}) + (n-1)\lambda + \rho$$

The definition of being unimodular is that $\text{tr}(\text{ad}_Z) = 0$ for all Z , so it follows that

$$(n-1)\lambda - (m-1)\rho = (n-1)\lambda + \rho.$$

In particular $\rho = 0$ and $\text{scal} = (n-1)\lambda$. The manifold splits by Theorem 1.8. \square

In the following we give an explicit construction of non-rigid $(\lambda, 4 + m)$ -Einstein metrics on solvable Lie groups with $\lambda < 0$ for all $m > 0$.

Let \mathfrak{g} be a 4-dimensional non-unimodular Lie algebra of the type $\mathfrak{g} = \mathfrak{a} + [\mathfrak{g}, \mathfrak{g}]$, where \mathfrak{a} is a 2-dimensional abelian subalgebra and $[\mathfrak{g}, \mathfrak{g}]$ the derived subalgebra, which we also assume to be abelian. Let g denote a left invariant metric on the corresponding simply connected Lie group G with X_0, X_1, X_2, X_3 an orthonormal frame of left invariant vector fields such that $\mathfrak{a} = \text{span}\{X_0, X_1\}$ and $[\mathfrak{g}, \mathfrak{g}] = \text{span}\{X_2, X_3\}$. The distribution \mathfrak{a} defines a totally geodesic foliation whose leaves are \mathbb{R}^2 . The distribution $[\mathfrak{g}, \mathfrak{g}]$ is Riemannian with vanishing A tensor and the intrinsic geometry of the leaves are also \mathbb{R}^2 . In other words we have a Riemannian submersion

$$\begin{array}{ccc} \mathbb{R}^2 & \longrightarrow & G \\ & & \downarrow \\ & & \mathbb{R}^2 \end{array}$$

where the base and intrinsic geometry of the fibers are flat and the A tensor for the submersion vanishes. We shall make further simplifying assumptions about the Lie algebra, but in general we see that the T tensor controls everything. The Ricci curvatures are computed using the follows properties.

Proposition 6.5. *The Ricci tensor preserves the distributions $\mathfrak{g} = \mathfrak{a} + [\mathfrak{g}, \mathfrak{g}]$. And for $i, j = 0, 1$*

$$\text{Ric}(X_i, X_j) = \sum_{k=2}^3 2g([X_i, X_k], \nabla_{X_k} X_j) + g(\nabla_{X_k} X_i, \nabla_{X_k} X_j)$$

and $k, l = 2, 3$

$$\text{Ric}(X_k, X_l) = \sum_{i \neq k, l} -g(\nabla_{X_i} X_i, \nabla_{X_k} X_l) + g(\nabla_{X_k} X_i, [X_i, X_l]) + g(\nabla_{X_l} X_i, [X_i, X_k]) + g(\nabla_{X_k} X_i, \nabla_{X_l} X_i)$$

Proof. This relies on our knowledge of the covariant derivatives, specifically that the foliation defined by \mathfrak{a} is totally geodesic. The only possible nonvanishing terms up to permutations of indices are terms of the form

$$g(\nabla_{X_i} X_j, X_k), \text{ where } k = 0, 1 \text{ and } i, j = 2, 3.$$

These can be computed by observing

$$\begin{aligned} g(\nabla_{X_i} X_j, X_k) &= \frac{1}{2}(g([X_k, X_i], X_j) + g([X_k, X_j], X_i)) \\ &= -g(X_j, \nabla_{X_i} X_k). \end{aligned}$$

Note also that these quantities are all constant since the metric is left invariant.

With this in mind we can do the calculations. For $i = 0, 1$

$$\begin{aligned} \text{Ric}(X_i) &= \sum_{k=1}^4 R(X_i, X_k) X_k \\ &= \sum_{k=3}^4 R(X_i, X_k) X_k \end{aligned}$$

and

$$\begin{aligned}
R(X_i, X_k)X_k &= \nabla_{X_i}\nabla_{X_k}X_k - \nabla_{X_k}\nabla_{X_i}X_k - \nabla_{[X_i, X_k]}X_k \\
&= -\nabla_{X_k}[X_i, X_k] - \nabla_{X_k}\nabla_{X_k}X_i - \nabla_{[X_i, X_k]}X_k \\
&= -2\nabla_{X_k}[X_i, X_k] - \nabla_{X_k}\nabla_{X_k}X_i \in \text{span}\{X_0, X_1\}.
\end{aligned}$$

It shows that the Ricci tensor preserves the distributions.

This also helps us calculate the Ricci curvatures when $i, j = 0, 1$

$$\begin{aligned}
\text{Ric}(X_i, X_j) &= \sum_{k=2}^3 R(X_i, X_k, X_k, X_j) \\
&= \sum_{k=2}^3 g(-2\nabla_{X_k}[X_i, X_k] - \nabla_{X_k}\nabla_{X_k}X_i, X_j) \\
&= \sum_{k=2}^3 2g([X_i, X_k], \nabla_{X_k}X_j) + g(\nabla_{X_k}X_i, \nabla_{X_k}X_j).
\end{aligned}$$

When $k, l = 2, 3$ we obtain similar formulas

$$\begin{aligned}
\text{Ric}(X_k, X_l) &= \sum_{i \neq k, l} R(X_k, X_i, X_i, X_l) \\
&= \sum_{i \neq k, l} g(\nabla_{X_k}\nabla_{X_i}X_i, X_l) - g(\nabla_{X_i}\nabla_{X_k}X_i, X_l) - g(\nabla_{[X_k, X_i]}X_i, X_l) \\
&= \sum_{i \neq k, l} -g(\nabla_{X_i}X_i, \nabla_{X_k}X_l) - g(\nabla_{X_i}\nabla_{X_k}X_i, X_l) - g(\nabla_{[X_k, X_i]}X_i, X_l) \\
&= \sum_{i \neq k, l} -g(\nabla_{X_i}X_i, \nabla_{X_k}X_l) + g(\nabla_{X_k}X_i, \nabla_{X_i}X_l) - g(\nabla_{X_l}X_i, [X_k, X_i]) \\
&= \sum_{i \neq k, l} -g(\nabla_{X_i}X_i, \nabla_{X_k}X_l) + g(\nabla_{X_k}X_i, [X_i, X_l]) + g(\nabla_{X_l}X_i, [X_i, X_k]) \\
&\quad + g(\nabla_{X_k}X_i, \nabla_{X_l}X_i)
\end{aligned}$$

that finishes the proof. \square

Next we need to determine the structure constants. We proceed as in the excellent reference [Je] also using [Mi] to find an appropriate 3 dimensional non-unimodular solvable Lie algebra satisfying

$$\begin{aligned}
[X_1, X_2] &= \alpha X_2 + \beta X_3 \\
[X_1, X_3] &= \gamma X_2 + (2 - \alpha)X_3,
\end{aligned}$$

where α, β and γ are constants. We will make the further simplifying assumption that $\gamma = \beta \neq 0$. Thus

$$\begin{aligned}
\nabla_{X_2}X_1 &= -\alpha X_2 - \frac{1}{2}\beta X_3 \\
\nabla_{X_3}X_1 &= -\beta X_2 - (2 - \alpha)X_3.
\end{aligned}$$

There are similar constraints for the structure constants

$$[X_0, X_i] = F_{ij}X_j, \quad \text{for } i, j = 2, 3,$$

that yield the other relevant covariant derivatives:

$$\begin{aligned}\nabla_{X_2}X_0 &= -F_{22}X_2 - \frac{1}{2}(F_{23} + F_{32})X_3 \\ \nabla_{X_3}X_0 &= -\frac{1}{2}(F_{23} + F_{32})X_2 - F_{33}X_3.\end{aligned}$$

However, we also need to see how they interact with the other structure constants. Checking the Jacobi identity shows

Proposition 6.6. *The constants α , β and F_{ij} satisfy the following identities:*

$$\begin{aligned}\beta F_{22} + 2(1 - \alpha)F_{23} - \beta F_{33} &= 0 \\ F_{23} &= F_{32}.\end{aligned}$$

Proof. To verify the Jacobi identity we only have to consider three vectors X_0 , X_i and X_j for $1 \leq i < j \leq 3$, i.e., the equations

$$\begin{aligned}[[X_0, X_1], X_2] + [[X_2, X_0], X_1] + [[X_1, X_2], X_0] &= 0 \\ [[X_0, X_1], X_3] + [[X_3, X_0], X_1] + [[X_1, X_3], X_0] &= 0 \\ [[X_0, X_2], X_3] + [[X_3, X_0], X_2] + [[X_2, X_3], X_0] &= 0\end{aligned}$$

which can be reduced to

$$\begin{aligned}-[F_{2j}X_j, X_1] + [\alpha X_2 + \beta X_3, X_0] &= 0 \\ -[F_{3j}X_j, X_1] + [\beta X_2 + (2 - \alpha)X_3, X_0] &= 0\end{aligned}$$

or more explicitly

$$\begin{aligned}F_{22}(\alpha X_2 + \beta X_3) + F_{23}(\beta X_2 + (2 - \alpha)X_3) - \alpha(F_{22}X_2 + F_{23}X_3) - \beta(F_{32}X_2 + F_{33}X_3) &= 0 \\ F_{32}(\alpha X_2 + \beta X_3) + F_{33}(\beta X_2 + (2 - \alpha)X_3) - \beta(F_{22}X_2 + F_{23}X_3) - (2 - \alpha)(F_{32}X_2 + F_{33}X_3) &= 0.\end{aligned}$$

This finally yields

$$\begin{aligned}\beta F_{23} - \beta F_{32} &= 0 \\ \beta F_{22} + (2 - 2\alpha)F_{23} - \beta F_{33} &= 0 \\ -\beta F_{22} - (2 - 2\alpha)F_{32} + \beta F_{33} &= 0 \\ -\beta F_{23} + \beta F_{32} &= 0.\end{aligned}$$

This immediately gives the relationships. \square

We now wish to solve the $(\lambda, 4 + m)$ -Einstein equations on the Lie group G . Let H be the Lie subgroup corresponding to the Lie subalgebra spanned by $\{X_1, X_2, X_3\}$ and r is the distance function from H . Since $\nabla_{X_0}X_0 = 0$ and X_0 is the unit normal vector field of H , we have $X_0 = \nabla r$. From the $(\lambda, n + m)$ -Einstein equation (6.1) in terms of left invariant vector fields, we have

Lemma 6.7. *The $(\lambda, 4 + m)$ -Einstein equations on G can be written as*

$$\begin{aligned}\text{Ric}(X_0, X_0) &= \rho \\ \text{Ric}(X_1, X_1) &= \lambda \\ \text{Ric}(X_1, X_0) &= 0 \\ \text{Ric}(X_2, X_2) - \sqrt{m(\rho - \lambda)}F_{22} &= \lambda \\ \text{Ric}(X_3, X_3) - \sqrt{m(\rho - \lambda)}F_{33} &= \lambda \\ \text{Ric}(X_2, X_3) - \sqrt{m(\rho - \lambda)}F_{32} &= 0.\end{aligned}$$

These equations impose further constraints on the structure constants and as we shall see the last three of them are equivalent when we know that the first three hold.

Theorem 6.8. *Given $m > 0$ and $\alpha, \beta \in \mathbb{R}$ with $\beta \neq 0$ and $(\alpha - 1)^2 + \beta^2 \neq 1$, we obtain a left invariant $(\lambda, 4 + m)$ -Einstein metric on G with $\mu = 0$. Moreover, this structure is not rigid.*

Proof. We start by observing that if we have such a solution then

$$\begin{aligned}
\text{Ric}(X_1, X_0) &= -\frac{2(-\alpha + \alpha^2 + \beta^2)F_{32} + 2\beta F_{33}}{\beta} = 0, \\
\text{Ric}(X_0, X_0) &= -\frac{2(F_{32})^2(2 - 4\alpha^3 + \alpha^4 + 3\beta^2 + \beta^4 - 2\alpha(3 + 2\beta^2) + \alpha^2(7 + 2\beta^2))}{\beta^2} = \rho, \\
\text{Ric}(X_1, X_1) &= -2(2 - 2\alpha + \alpha^2 + \beta^2) = \lambda, \\
\text{Ric}(X_2, X_2) &= -\frac{2(\alpha\beta^2 + (F_{32})^2(2 - 5\alpha^3 + \alpha^4 + 3\beta^2 + \beta^4 + \alpha^2(9 + 2\beta^2) - \alpha(7 + 5\beta^2)))}{\beta^2} \\
&= \lambda - \frac{(F_{32})(2 - 3\alpha + \alpha^2 + \beta^2)}{\beta} \sqrt{m(\rho - \lambda)}, \\
\text{Ric}(X_2, X_3) &= \frac{2(-\beta^2 + (F_{32})^2(1 - 2\alpha + \alpha^2 + \beta^2))}{\beta} = (F_{32}) \sqrt{m(\rho - \lambda)}, \\
\text{Ric}(X_3, X_3) &= -\frac{2((2 - \alpha)\beta^2 + (F_{32})^2(-3\alpha^3 + \alpha^4 + \beta^2 + \beta^4 + \alpha^2(3 + 2\beta^2) - \alpha(1 + 3\beta^2)))}{\beta^2} \\
&= \lambda - \frac{(F_{32})(-\alpha + \alpha^2 + \beta^2)}{\beta} \sqrt{m(\rho - \lambda)}.
\end{aligned}$$

The first equation forces the relationship

$$F_{33} = -\frac{-\alpha + \alpha^2 + \beta^2}{\beta} F_{32}.$$

If we introduce the notation

$$z = \frac{F_{32}}{\beta}$$

then we have to solve the equations

$$\begin{aligned}
\text{Ric}(X_0, X_0) &= -2z^2(2 - 4\alpha^3 + \alpha^4 + 3\beta^2 + \beta^4 - 2\alpha(3 + 2\beta^2) + \alpha^2(7 + 2\beta^2)) = \rho, \\
\text{Ric}(X_1, X_1) &= -2(2 - 2\alpha + \alpha^2 + \beta^2) = \lambda, \\
\text{Ric}(X_2, X_2) &= -2\alpha - 2z^2(2 - 5\alpha^3 + \alpha^4 + 3\beta^2 + \beta^4 + \alpha^2(9 + 2\beta^2) - \alpha(7 + 5\beta^2)) \\
&= \lambda - z(2 - 3\alpha + \alpha^2 + \beta^2) \sqrt{m(\rho - \lambda)}, \\
\frac{1}{\beta} \text{Ric}(X_2, X_3) &= -2 + 2z^2(1 - 2\alpha + \alpha^2 + \beta^2) = z \sqrt{m(\rho - \lambda)}, \\
\text{Ric}(X_3, X_3) &= -2(2 - \alpha)\beta^2 - 2z^2(-3\alpha^3 + \alpha^4 + \beta^2 + \beta^4 + \alpha^2(3 + 2\beta^2) - \alpha(1 + 3\beta^2)) \\
&= \lambda - z(-\alpha + \alpha^2 + \beta^2) \sqrt{m(\rho - \lambda)}.
\end{aligned}$$

Note that the first equation combined with the second yields

$$\begin{aligned}\rho &= -2z^2(2 - 4\alpha^3 + \alpha^4 + 3\beta^2 + \beta^4 - 2\alpha(3 + 2\beta^2) + \alpha^2(7 + 2\beta^2)) \\ &= -2z^2(2 - 2\alpha + \alpha^2 + \beta^2)(1 - 2\alpha + \alpha^2 + \beta^2) \\ &= -z^2 \frac{\lambda(\lambda + 2)}{2}\end{aligned}$$

showing

$$\begin{aligned}\rho - \lambda &= -z^2 \frac{\lambda(\lambda + 2)}{2} - \lambda \\ &= -\lambda \left(\frac{\lambda + 2}{2} z^2 + 1 \right).\end{aligned}$$

The last three equations turn out to be identical. Specifically, the fourth equation is the same as the third if we factor by $(2 - 3\alpha + \alpha^2 + \beta^2)$ and the same as the fifth if we factor by $(-\alpha + \alpha^2 + \beta^2)$. Thus these three equations reduce to

$$-2 + 2z^2(1 - 2\alpha + \alpha^2 + \beta^2) = z\sqrt{m(\rho - \lambda)}.$$

or

$$-2 - (\lambda + 2)z^2 = z\sqrt{-\lambda m \left(\frac{\lambda + 2}{2} z^2 + 1 \right)}.$$

This calculates z as a function of λ . Either

$$z^2 = \frac{-2}{\lambda + 2}$$

forcing $\rho = \lambda$, or

$$(6.2) \quad z^2 = \frac{2}{-\frac{\lambda}{2}m - (\lambda + 2)} = \frac{-4}{\lambda(m + 2) + 4}$$

which is equivalent to the first when $m = 0$. Only the latter case with $m > 0$ gives us a nontrivial solution. If we substitute it into the equation for ρ we are left with the two equations

$$(6.3) \quad \begin{aligned}-2(2 - 2\alpha + \alpha^2 + \beta^2) &= \lambda, \\ \frac{2\lambda(\lambda + 2)}{\lambda(m + 2) + 4} &= \rho.\end{aligned}$$

Note that we always have

$$2 - 2\alpha + \alpha^2 + \beta^2 \geq 1$$

so $\lambda < -2$.

Finally we do a few calculations to show that this is not rigid. Since $\lambda \neq \rho$ it can only be rigid when the other two eigenvalues are also λ, ρ .

We have

$$\begin{aligned}z\sqrt{m(\rho - \lambda)} &= -2 - (\lambda + 2) \frac{-4}{\lambda(m + 2) + 4} \\ &= \frac{-2(\lambda(m + 2) + 4) + 4(\lambda + 2)}{\lambda(m + 2) + 4} = \frac{-2\lambda m}{\lambda(m + 2) + 4} \\ &= -\frac{m\rho}{\lambda + 2}.\end{aligned}$$

Thus

$$\begin{aligned} \text{scal} &= \lambda + \rho + \text{Ric}(X_2, X_2) + \text{Ric}(X_3, X_3) \\ &= \lambda + \rho + 2\lambda + (\lambda + 2)z\sqrt{m(\rho - \lambda)} \\ &= 3\lambda - (m - 1)\rho. \end{aligned}$$

So the only possibility for having a rigid solution is when

$$3\lambda - (m - 1)\rho = 2\rho + 2\lambda$$

which implies

$$\lambda = (m + 1)\rho = \frac{2\lambda(\lambda + 2)(m + 1)}{\lambda(m + 2) + 4}$$

or

$$\frac{2(\lambda + 2)(m + 1)}{\lambda(m + 2) + 4} = 1, \quad \text{i.e.,} \quad \lambda = -4.$$

It follows that

$$\rho = -\frac{4}{m + 1}$$

and

$$\text{Ric}(X_2, X_2) = -4 + \frac{2m}{m + 1}(2 - \alpha), \quad \text{Ric}(X_2, X_3) = -\frac{2m}{m + 1}\beta, \quad \text{Ric}(X_3, X_3) = -4 + \frac{2m}{m + 1}\alpha.$$

The principal Ricci curvatures are λ and ρ if $(\alpha - 1)^2 + \beta^2 = 1$. In this case, the metric splits along the eigendistributions of Ric . \square

Finally we consider the convergence of the metrics in Theorem 6.8 as $m \rightarrow \infty$. We will see that the Ricci soliton structure naturally appears in the limiting metric. First we recall

Definition 6.9. A left invariant metric g on a simply-connected nilpotent (or solvable) Lie group is called a *nilsoliton* (or *solvsoliton*) if the Ricci tensor satisfies the following equation

$$(6.4) \quad \text{Ric}(g) = cI + D$$

where c is a constant and $D \in \text{Der}(\mathfrak{g})$ is a derivation.

These manifolds are closely related to Ricci flows and Einstein metrics on non-compact homogeneous spaces, see [La1, La2, La3] and references therein.

Theorem 6.10. *Let (G, g_m) be the $(\lambda, 4 + m)$ -Einstein metrics in Theorem 6.8. As $m \rightarrow \infty$, they converge in C^∞ to the Riemannian product $(\mathbb{R} \times H, dt^2 + g_0)$ where (H, g_0) is a three dimensional solvsoliton.*

Proof. Recall that $\{X_0, X_1, X_2, X_3\}$ is an orthonormal frame of left invariant vector fields on G and then any left invariant metric g is determined by the structure constants $C_{ijk} = g([X_i, X_j], X_k)$ for $i, j, k = 0, \dots, 3$. That the metrics g_m converge to g_∞ in C^∞ is equivalent to that the structure constants C_{ijk}^m of g_m converge to those of g_∞ , see for example [La3, Proposition 2.8] when G is a nilpotent group. Alternatively, the C^∞ convergence of the structure constants implies C^∞ convergence of the Levi-Civita connections and then the convergence of the metrics follows, see for example [GP, Section 6].

From the equation (6.3) of λ in terms of α and β , since λ is unchanged for the metrics g_m , α and β are also unchanged. From the equation (6.2)

$$\lim_{m \rightarrow \infty} z^2 = \lim_{m \rightarrow \infty} \frac{-4}{\lambda(m+2)+4} = 0$$

It follows that F_{32} and F_{33} converge to zero, i.e., $[X_0, X_i] = 0$ for all $i = 0, \dots, 3$ as $m \rightarrow \infty$. So the Lie algebra \mathfrak{g} corresponding to g_∞ on G splits as direct sum of $\mathbb{R}X_0 \oplus \mathfrak{h}$ where \mathfrak{h} is spanned by X_1, X_2 and X_3 , and the limiting metric $g_\infty = dt^2 + g_0$ where g_0 is the restriction on the Lie group H whose Lie algebra is \mathfrak{h} . On H , the nonzero Lie brackets are given by

$$\text{ad}_{X_1} \begin{pmatrix} X_2 \\ X_3 \end{pmatrix} = \begin{pmatrix} \alpha & \beta \\ \beta & 2 - \alpha \end{pmatrix} \begin{pmatrix} X_2 \\ X_3 \end{pmatrix}.$$

A computation shows that the nonzero Ricci curvatures are

$$\begin{aligned} \text{Ric}(X_1, X_1) &= \lambda \\ \text{Ric}(X_2, X_2) &= -2\alpha \\ \text{Ric}(X_2, X_3) &= -2\beta \\ \text{Ric}(X_3, X_3) &= -2(2 - \alpha). \end{aligned}$$

Let D be the derivation on \mathfrak{h} as $D(X_1) = 0$ and $D(X_i) = -\lambda X_i - 2\text{ad}_{X_1}(X_i)$ for $i = 2, 3$, then we have $\text{Ric} = \lambda I + D$ which shows (H, g_0) is a solvsoliton. The solvsoliton structure can also be obtained by viewing \mathfrak{h} as an extension of the abelian Lie algebra $\text{span}_{\mathbb{R}}\{X_2, X_3\}$ by $\mathbb{R}X_1$, see the construction in [La2, Proposition 4.3]. Note the solvsoliton metric is not Einstein since $\beta \neq 0$.

This derivation can be constructed directly as a limit in the following way: we know that

$$\frac{m}{w} \nabla \nabla w = \sqrt{m(\rho - \lambda)} \nabla X_0 + (\rho - \lambda) X_0^\flat \otimes X_0^\flat,$$

where X_0^\flat is a 1-form dual to X_0 . And the only nontrivial covariant derivatives for X_0 are

$$\begin{aligned} \nabla_{X_2} X_0 &= (2 - 3\alpha + \alpha^2 + \beta^2) - \beta z X_3 \\ &= \left(-\frac{\lambda}{2} - \alpha\right) z X_2 - \beta z X_3 \\ \nabla_{X_3} X_0 &= -\beta z X_2 + (-\alpha + \alpha^2 + \beta^2) z X_3 \\ &= -\beta z X_2 + \left(-\frac{\lambda}{2} + 2 - \alpha\right) z X_3. \end{aligned}$$

Next note that

$$\lim_{m \rightarrow \infty} z \sqrt{m(\rho - \lambda)} = \lim_{m \rightarrow \infty} z \sqrt{\frac{-4m(\rho - \lambda)}{\lambda(m+2)+4}} = -2$$

so it follows that $\frac{m}{w} \nabla \nabla w$ converges to the derivation D as $m \rightarrow \infty$.

A similar limit argument also shows that the Ricci curvatures of g_m converge to those of the limiting metric g_∞ . \square

APPENDIX A. AN INTEGRABILITY CRITERION OF EIGEN-DISTRIBUTION OF Ric

In this appendix we will show the eigen-distributions of P (or Ric) in Theorem 1.10 are integrable without using the curvature growth assumption. It follows from the following general integrability criterion of eigen-distribution of Ricci curvature.

Theorem A.1. *Suppose $g_t = h_0 + th_1$ ($t > 0$) is a family of Riemannian metric on Σ such that*

$$\begin{aligned} T\Sigma &= \mathcal{N} \oplus \mathcal{P}, \\ g_1|_{\mathcal{N}} &= h_0, \\ g_1|_{\mathcal{P}} &= h_1, \end{aligned}$$

where \mathcal{N} and \mathcal{P} are two orthogonal distributions. If the Ricci curvature of g_t satisfies the following condition

$$\text{Ric}^t = \lambda h_0 + \sigma h_1$$

where $\lambda \neq \sigma$ are two constants, then the two distributions \mathcal{N} and \mathcal{P} are integrable.

Before proving Theorem A.1, we show that in Theorem 1.10 the restriction of the metric g on the level sets of ∇r satisfies the assumptions in Theorem A.1. Then the integrability of the eigen-distributions of P follows directly.

We denote orthonormal frames of \mathcal{N} and \mathcal{P} by $\{E_i\}$ and $\{E_a\}$ respectively and, use i, j, k, \dots for the indices of \mathcal{N} , a, b, c, \dots for \mathcal{P} and $\alpha, \beta, \gamma, \dots$. The vector fields in \mathcal{N} and \mathcal{P} are denoted by X, Y, Z, \dots and U, V, W, \dots respectively. If A is a vector field in TM , then $\mathcal{N}A$ and $\mathcal{P}A$ denote its projections.

Recall from Lemma 5.1 the eigenvalue of P is either 0 or $\lambda - \rho$. Thus we have the orthogonal decomposition of the tangent space of the level sets Σ of r as

$$\begin{aligned} M &= I \times \Sigma, \\ g &= dr^2 + g_r, \\ T\Sigma &= \mathcal{N} \oplus \mathcal{P}. \end{aligned}$$

Instead of parametrizing by r , it will turn out to be more convenient to parametrize the metrics on the level sets by $t = (w')^2(r)$. We consider the one parameter family of metrics g_t on Σ . In the next proposition we compute the Ricci curvature of this family.

Proposition A.2. *The metrics (Σ, g_t) satisfy*

$$\begin{aligned} g_t &= h_0 + th_1 \\ \text{Ric}^t &= \lambda h_0 + \sigma h_1 \\ \text{scal}^t &= \lambda \dim \mathcal{N} + \frac{\sigma}{t} \dim \mathcal{P}, \end{aligned}$$

where Ric^t and scal^t denote the curvatures of g_t , h_0, h_1 are metrics on \mathcal{N} and \mathcal{P} respectively, and $\sigma = (\dim \mathcal{P} - 1) \bar{k} \bar{\mu}$.

Remark A.3. By multiplying w by a constant, we can always assume that $\sigma \neq \lambda$.

Remark A.4. In Proposition A.2, instead of considering the metrics on a family of hypersurfaces in M we are fixing just one hypersurface Σ and then pulling the metrics on the other hypersurfaces back via the flow generated by ∇r . If $\{E_i\}, \{E_a\}$ are orthonormal frames of \mathcal{N}, \mathcal{P} respectively with respect to g_t , then the formula tells us that $\{e_i = E_i\}$ and $\{e_a = \sqrt{t}E_a\}$ are orthonormal frames with respect to g_1 .

Proof. Write the metric splitting on Σ corresponding to some regular value w_0 as

$$g_0 = g_{\mathcal{N}} + g_{\mathcal{P}}.$$

From the $(\lambda, n + m)$ -Einstein equation we know that

$$\begin{aligned}\mathcal{L}_{\nabla w} g_{\mathcal{N}} &= 0, \\ \mathcal{L}_{\nabla w} g_{\mathcal{P}} &= -2w\bar{k}g_{\mathcal{P}},\end{aligned}$$

and using $w = w(r)$ where $w'' = -\bar{k}w$ we obtain

$$\begin{aligned}\mathcal{L}_{\nabla r} g_{\mathcal{N}} &= 0, \\ \mathcal{L}_{\nabla r} g_{\mathcal{P}} &= -2\bar{k}\frac{w}{w'}g_{\mathcal{P}},\end{aligned}$$

so the metric has the form

$$g = dr^2 + h_0 + (w')^2 h_1$$

where h_0 and h_1 are restrictions of g on \mathcal{N} and \mathcal{P} at the regular value w_0 respectively.

As in the proof of Proposition 5.2, we have the following formulas for the second fundamental form:

$$\begin{aligned}\text{II}(X, X) &= 0 \\ \text{II}(X, U) &= 0 \\ \text{II}(U, U) &= -\bar{k}\frac{w}{w'}|U|^2.\end{aligned}$$

The radial flatness assumption

$$0 = Q(\nabla w, E, E, \nabla w) = R(E, \nabla w, \nabla w, E) + \frac{|\nabla w|^2}{m}(\text{Ric}(E, E) - \lambda|E|^2).$$

then tells us that the radial curvatures satisfy

$$\begin{aligned}R(X, \nabla r) \nabla r &= 0, \\ R(U, \nabla r) \nabla r &= \bar{k}U,\end{aligned}$$

and $\text{Ric}(\nabla r, \nabla r) = \rho$ implies that

$$\rho = \bar{k} \dim \mathcal{P}.$$

This implies that for any two unit vectors $X \in \mathcal{N}$ and $U \in \mathcal{P}$, we have

$$\begin{aligned}\text{Ric}_{g_r}(X, X) &= \lambda g_r(X, X) = \lambda \\ \text{Ric}_{g_r}(X, U) &= 0 \\ \text{Ric}_{g_r}(U, U) &= \rho - \bar{k} + (\dim \mathcal{P} - 1)\bar{k}^2 \frac{w^2}{(w')^2} = (\dim \mathcal{P} - 1)\bar{k} \left(\bar{k} \frac{w^2}{(w')^2} + 1 \right) \\ &= (\dim \mathcal{P} - 1)\bar{k}\bar{\mu}(w')^{-2},\end{aligned}$$

where

$$\begin{aligned}\bar{\mu} &= \bar{k}w^2 + |\nabla w|^2 \\ &= \bar{k}w^2 + (w')^2,\end{aligned}$$

which gives us

$$\begin{aligned}
\text{Ric}^t(E, F) &= \text{Ric}_{g_t}(E, F) \\
&= \lambda h_0(E, F) + (\dim \mathcal{P} - 1) \bar{k} \bar{\mu} h_1(E, F) \\
&= \lambda h_0(E, F) + \sigma h_1(E, F) \\
\sigma &= (\dim \mathcal{P} - 1) \bar{k} \bar{\mu}.
\end{aligned}$$

The formula for the scalar curvature then follows from tracing this equation. \square

As the scalar curvature of g_t is constant, we have the following

Corollary A.5. *For the metric g_t , we have*

$$\sum_i \nabla_{E_i} E_i \in \mathcal{N}, \quad \text{and} \quad \sum_a \nabla_{E_a} E_a \in \mathcal{P}.$$

Proof. The scalar curvature is constant on Σ . Thus, by the second Bianchi identity,

$$0 = \text{div}_{g_t} \text{Ric}^t = \lambda \text{div}_{g_t} h_0 + \sigma \text{div}_{g_t} h_1.$$

On the other hand

$$0 = \text{div}_{g_t} g_t = \text{div}_{g_t} h_0 + t \text{div}_{g_t} h_1.$$

So if $t \neq \sigma/\lambda$, then

$$\text{div}_{g_t} h_0 = \text{div}_{g_t} h_1 = 0.$$

For any vector $X \in \mathcal{N}$ we then have

$$\begin{aligned}
0 &= (\text{div} h_0)(X) = \sum_{\alpha} (\nabla_{E_{\alpha}} h_0)(E_{\alpha}, X) \\
&= \nabla_{E_{\alpha}} (h_0(E_{\alpha}, X)) - h_0(\nabla_{E_{\alpha}} E_{\alpha}, X) - h_0(E_{\alpha}, \nabla_{E_{\alpha}} X) \\
&= \nabla_{E_{\alpha}} (g_t(E_{\alpha}, X)) - g_t(\nabla_{E_{\alpha}} E_{\alpha}, X) - g_t(E_{\alpha}, \nabla_{E_{\alpha}} X) \\
&= g_t(\nabla_{E_{\alpha}} E_{\alpha}, X),
\end{aligned}$$

which shows that $\sum_a \nabla_{E_a} E_a \in \mathcal{P}$ when $t \neq \sigma/\lambda$. However, by continuity, this must also be true at $t = \sigma/\lambda$. Similarly, we have $\sum_i \nabla_{E_i} E_i \in \mathcal{N}$. \square

The integrability of the eigen-distributions follows from a more careful analysis of the formula for Ric^t .

Proof of Theorem A.1. We consider g_t as a variation along the h_1 direction as $g_t(s) = g_{t+s}$ for $s > 0$. From the assumption of the Ricci curvature, we have

$$\text{Ric}^{t+s}(E_i, E_j) = \lambda \delta_{ij} \quad \text{and} \quad \text{Ric}^{t+s}(E_a, E_b) = \sigma h_1(E_a, E_b) = \frac{\sigma}{t} \delta_{ab}.$$

As $\text{tr}_{g_t} h_1$ is constant and $\text{div}_{g_t} h_1 = 0$, it follows from the first variation formula of the Ricci tensor, see [Be, 1.174(d)] that

$$0 = \frac{1}{2} \Delta_L^t h_1,$$

where Δ_L^t is the Lichnerowicz Laplacian with respect to the metric g_t , see [Be, 1.180b]. In the following we compute the expansion of Δ_L^t at $t = 1$. The vanishing of the coefficients will show that both distributions are integrable.

Using Koszul's formula, the connection of g_t is determined by

$$\begin{aligned} g_t(\nabla_X Y, Z) &= g_1(\nabla_X Y, Z) \\ g_t(\nabla_X Y, U) &= g_1(\nabla_X Y, U) + \frac{t-1}{2}g_1([X, Y], U) \\ g_t(\nabla_X U, Y) &= g_1(\nabla_X U, Y) + \frac{t-1}{2}g_1([Y, X], U) \\ g_t(\nabla_X U, V) &= tg_1(\nabla_X U, V) + \frac{t-1}{2}g_1([U, V], X); \end{aligned}$$

and

$$\begin{aligned} g_t(\nabla_U X, Y) &= g_1(\nabla_U X, Y) + \frac{t-1}{2}g_1([Y, X], U) \\ g_t(\nabla_U X, V) &= tg_1(\nabla_U X, V) + \frac{t-1}{2}g_1([U, V], X) \\ g_t(\nabla_U V, W) &= tg_1(\nabla_U V, W) \\ g_t(\nabla_U V, X) &= tg_1(\nabla_U V, X) + \frac{t-1}{2}g_1([V, U], X). \end{aligned}$$

Since $\text{Ric} = \lambda h_0 + \sigma h_1$, we have $\text{Ric} \circ h_1 = h_1 \circ \text{Ric} = \frac{\sigma}{t} h_1$ as symmetric 2-tensors which implies that

$$(A.1) \quad \Delta_L^t h_1 = \nabla^* \nabla h_1 + \frac{2\sigma}{t} h_1 - 2\dot{R}_{g_t} h_1,$$

where $\dot{R}_g h$ is defined by

$$(\dot{R}_{g_t} h)(A, B) = \sum_{\alpha} h(R^t(E_{\alpha}, A)B, E_{\alpha}) = \sum_{\alpha, \beta} R^t(E_{\alpha}, A, B, E_{\beta})h(E_{\alpha}, E_{\beta})$$

for any symmetric 2-tensor h and R^t is the (3,1) or (4,0) curvature tensor of g_t . It follows that

$$\dot{R}_{g_t} h_1(A, B) = \frac{1}{t} \sum_a R^t(E_a, A, B, E_a).$$

Let $H = \sum_{\alpha} \nabla_{E_{\alpha}} E_{\alpha}$ and then the term $\nabla^* \nabla h_1$ can be computed as

$$\begin{aligned} &(\nabla^* \nabla h_1)(A, B) \\ &= - \sum_{\alpha} (\nabla_{E_{\alpha}, E_{\alpha}}^2 h_1)(A, B) \\ &= -\nabla_{E_{\alpha}} ((\nabla h_1)(E_{\alpha}, A, B)) + (\nabla h_1)(H, A, B) + (\nabla h_1)(E_{\alpha}, \nabla_{E_{\alpha}} A, B) + (\nabla h_1)(E_{\alpha}, A, \nabla_{E_{\alpha}} B) \\ &= -\nabla_{E_{\alpha}} \nabla_{E_{\alpha}} (h_1(A, B)) + 2\nabla_{E_{\alpha}} (h_1(\nabla_{E_{\alpha}} A, B)) + 2\nabla_{E_{\alpha}} (h_1(A, \nabla_{E_{\alpha}} B)) \\ &\quad + \nabla_H (h_1(A, B)) - h_1(\nabla_H A, B) - h_1(A, \nabla_H B) \\ &\quad - h_1(\nabla_{E_{\alpha}} \nabla_{E_{\alpha}} A, B) - 2h_1(\nabla_{E_{\alpha}} A, \nabla_{E_{\alpha}} B) - h_1(A, \nabla_{E_{\alpha}} \nabla_{E_{\alpha}} B). \end{aligned}$$

It follows that

$$\begin{aligned}
(\nabla^* \nabla h_1)(X, Y) &= -2h_1(\nabla_{E_\alpha} X, \nabla_{E_\alpha} Y) \\
&= -\frac{2}{t} g_t(\nabla_{E_\alpha} X, E_a) g_t(\nabla_{E_\alpha} Y, E_a), \\
(\nabla^* \nabla h_1)(X, U) &= \frac{1}{t} g_t(\nabla_{E_\alpha} \nabla_{E_\alpha} X, U) + \frac{2}{t} g_t(\nabla_{E_\alpha} X, E_i) g_t(\nabla_{E_\alpha} U, E_i) - \frac{1}{t} g_t(\nabla_H X, U), \\
(\nabla^* \nabla h_1)(U, V) &= -\frac{1}{t} \nabla_{E_\alpha} \nabla_{E_\alpha} (g_t(U, V)) + \frac{2}{t} \nabla_{E_\alpha} (g_t(\nabla_{E_\alpha} U, V)) + \frac{2}{t} \nabla_{E_\alpha} (g_t(U, \nabla_{E_\alpha} V)) \\
&\quad - \frac{1}{t} g_t(\nabla_{E_\alpha} \nabla_{E_\alpha} U, V) - \frac{2}{t} g_t(\nabla_{E_\alpha} U, E_a) g_t(\nabla_{E_\alpha} V, E_a) - \frac{1}{t} g_t(U, \nabla_{E_\alpha} \nabla_{E_\alpha} V) \\
&= \frac{2}{t} g_t(\nabla_{E_\alpha} U, \nabla_{E_\alpha} V) - \frac{2}{t} g_t(\nabla_{E_\alpha} U, E_a) g_t(\nabla_{E_\alpha} V, E_a) \\
&= \frac{2}{t} g_t(\nabla_{E_\alpha} U, E_i) g_t(\nabla_{E_\alpha} V, E_i).
\end{aligned}$$

We choose an arbitrary vector field U in \mathcal{P} . Since $(\Delta_L^t h_1)(U, U) = 0$, we have

$$(A.2) \quad \frac{t}{2} (\nabla^* \nabla h_1)(U, U) + \sigma h_1(U, U) = \sum_a R^t(E_a, U, U, E_a).$$

The first term on the left hand side is

$$\begin{aligned}
\frac{t}{2} (\nabla^* \nabla h_1)(U, U) &= g_t(\nabla_{E_\alpha} U, E_i) g_t(\nabla_{E_\alpha} U, E_i) = g_t(\nabla_{E_j} U, E_i)^2 + g_t(\nabla_{E_a} U, E_i)^2 \\
&= \left(g_1(\nabla_{E_j} U, E_i) + \frac{t-1}{2} g_1([E_i, E_j], U) \right)^2 \\
&\quad + \left(t g_1(\nabla_{E_a} U, E_i) + \frac{t-1}{2} g_1([U, E_a], E_i) \right)^2 \\
&= \left(g_1(\nabla_{e_j} U, e_i) + \frac{t-1}{2} g_1([e_i, e_j], U) \right)^2 \\
&\quad + \left(\sqrt{t} g_1(\nabla_{e_a} U, e_i) + \frac{t-1}{2\sqrt{t}} g_1([U, e_a], e_i) \right)^2.
\end{aligned}$$

In particular, the highest power of t in the expansion above is t^2 and the lowest one is t^{-1} . We call the terms with t 's power 0 or 1 the middle order terms. Then we have

$$\frac{t}{2} (\nabla^* \nabla h_1)(U, U) = \frac{t^2}{4} \sum_{i,j} g_1([e_i, e_j], U)^2 + \frac{1}{4t} \sum_{a,i} g_1([U, e_a], e_i)^2 + \text{middle order terms}.$$

The second term on the left hand side is

$$\sigma h_1(U, U) = \sigma g_1(U, U).$$

Therefore the expansion of the left hand side is given by

$$(A.3) \quad \left(\frac{t}{2} (\nabla^* \nabla h_1) + \sigma h_1 \right) (U, U) = \frac{t^2}{4} \sum_{i,j} g_1([e_i, e_j], U)^2 + \frac{1}{4t} \sum_{a,i} g_1([U, e_a], e_i)^2 + \text{middle order terms}.$$

The right hand side is

$$\begin{aligned} \sum_a R^t(E_a, U, U, E_a) &= D_{E_a} g_t(\nabla_U U, E_a) - g_t(\nabla_U U, \sum_a \nabla_{E_a} E_a) - D_U g_t(\nabla_{E_a} U, E_a) \\ &\quad + g_t(\nabla_{E_a} U, \nabla_U E_a) - g_t(\nabla_{[E_a, U]} U, E_a). \end{aligned}$$

We calculate each term in $R^t(E_a, U, U, E_a)$ and only consider the term t^2 and t^{-1} . We have

$$\begin{aligned} D_{E_a} g_t(\nabla_U U, E_a) &= D_{E_a} (t g_1(\nabla_U U, E_a)) = D_{e_a} g_1(\nabla_U U, e_a) = \text{middle order terms,} \\ g_t(\nabla_U U, \nabla_{E_a} E_a) &= g_t(\nabla_U U, E_b) g_t(\nabla_{E_a} E_a, E_b) = t g_1(\nabla_U U, E_b) \cdot t g_1(\nabla_{E_a} E_a, E_b) \\ &= g_1(\nabla_U U, e_b) g_1(\nabla_{e_a} e_a, e_b) = g_1(\nabla_U U, \nabla_{e_a} e_a) = \text{middle order terms,} \\ D_U g_t(\nabla_{E_a} U, E_a) &= D_U (t g_1(\nabla_{E_a} U, E_a)) = D_U g_1(\nabla_{e_a} U, e_a) = \text{middle order terms;} \end{aligned}$$

and

$$\begin{aligned} g_t(\nabla_{E_a} U, \nabla_U E_a) &= g_t(\nabla_{E_a} U, E_i) g_t(\nabla_U E_a, E_i) + g_t(\nabla_{E_a} U, E_b) g_t(\nabla_U E_a, E_b) \\ &= \left(t g_1(\nabla_{E_a} U, E_i) + \frac{t-1}{2} g_1([U, E_a], E_i) \right) \cdot \\ &\quad \left(t g_1(\nabla_U E_a, E_i) + \frac{t-1}{2} g_1([E_a, U], E_i) \right) \\ &\quad + t g_1(\nabla_{E_a} U, E_b) \cdot t g_1(\nabla_U E_a, E_b) \\ &= \left(\sqrt{t} g_1(\nabla_{e_a} U, e_i) + \frac{t-1}{2\sqrt{t}} g_1([U, e_a], e_i) \right) \cdot \\ &\quad \left(\sqrt{t} g_1(\nabla_U e_a, e_i) + \frac{t-1}{2\sqrt{t}} g_1([e_a, U], e_i) \right) \\ &\quad + g_1(\nabla_{e_a} U, e_b) \cdot g_1(\nabla_U e_a, e_b) \\ &= -\frac{1}{4t} \sum_{a,i} g_1([U, e_a], e_i)^2 + \text{middle order terms,} \\ g_t(\nabla_{[E_a, U]} U, E_a) &= g_t(\nabla_{\mathcal{N}[E_a, U]} U, E_a) + g_t(\nabla_{\mathcal{P}[E_a, U]} U, E_a) \\ &= t g_1(\nabla_{\mathcal{N}[E_a, U]} U, E_a) + \frac{t-1}{2} g_1([U, E_a], \mathcal{N}[E_a, U]) + t g_1(\nabla_{\mathcal{P}[E_a, U]} U, E_a) \\ &= g_1(\nabla_{\mathcal{N}[e_a, U]} U, e_a) + \frac{t-1}{2t} g_1([U, e_a], \mathcal{N}[e_a, U]) + g_1(\nabla_{\mathcal{P}[e_a, U]} U, e_a) \\ &= \frac{1}{2t} \sum_{a,i} g_1([U, e_a], e_i)^2 + \text{middle order terms.} \end{aligned}$$

It follows that

$$(A.4) \quad \sum_a R^t(E_a, U, U, E_a) = -\frac{3}{4t} \sum_{a,i} g_1([U, e_a], e_i)^2 + \text{middle order terms.}$$

Since the two expansions given by (A.3) and (A.4) in the variable t are equal and U is an arbitrary vector field in \mathcal{P} , both distributions \mathcal{N} and \mathcal{P} are integrable. \square

REFERENCES

- [An] M. T. Anderson, *Scalar curvature, metric degenerations and the static vacuum Einstein equations on 3-manifolds. I*, *Geom. Funct. Anal.* **9**(1999), no. 5, 855–967.
- [Be] A. L. Besse, *Einstein manifolds*, *Ergebnisse der Mathematik und ihrer Grenzgebiete(3)[Results in Mathematics and Related Areas(3)]*, **10**. Springer-Verlag, Berlin, 1987.
- [BH] R. A. Blumenthal and J. J. Hebda, *De Rham decomposition theorems for foliated manifolds*, *Ann. Inst. Fourier(Grenoble)* **33**(1983), 183–198.
- [Bö1] C. Böhm, *Inhomogeneous Einstein metrics on low-dimensional spheres and other low-dimensional spaces*, *Invent. Math.* (1998) vol. **134**, no 1., 145–176.
- [Bö2] C. Böhm, *Non-compact cohomogeneity one Einstein manifolds*, *Bull. Soc. math. France* (1999) vol. **127**, 135–177.
- [Bu] P. Bueken, *Three-dimensional Riemannian manifolds with constant principal Ricci curvatures $\rho_1 = \rho_2 \neq \rho_3$* , *J. Math. Phys.* **37**(1996), 4062–4075.
- [BV] P. Bueken and L. Vanhecke, *Three- and four-dimensional Einstein-like manifolds and homogeneity*, *Geom. Dedicata*, **75**(1999), 123–136.
- [CSW] J. Case, Y.-J. Shu, and G. Wei, *Rigidity of Quasi-Einstein Metrics*, arXiv: 0805.3132v1, 2008.
- [CWZ] X. Cao, B. Wang and Z. Zhang, *On locally conformally flat gradient shrinking Ricci solitons*, arXiv: 0807.0588.
- [Co] J. Corvino, *Scalar curvature deformation and a gluing construction for the Einstein constraint equations*, *Comm. Math. Phys.* **214** (2000), no. 1, 137–189.
- [ENM] M. Eminent, G. La Nave and C. Mantegazza, *Ricci solitons: the equation point of view*, *Manuscripta Math.* **127**, (2008) 345–367.
- [GP] D. Glickenstein and T. L. Payne, *Ricci flow on three-dimensional, unimodular metric Lie algebras*, arXiv: 0909.0938.
- [GW] D. Gromoll and G. Walschap, *Metric foliations and curvature*. *Progress in Mathematics*, **268**. Birkhäuser Verlag, Basel, 2009.
- [Ha] R. Hamilton. *Three-manifolds with positive Ricci curvature*, *J. Diff. Geom.* **17**(1982), 255–306.
- [HPW1] C. He, P. Petersen and W. Wylie, *On the classification of warped product Einstein metrics*, arXiv:1010.5488.
- [HPW2] C. He, P. Petersen and W. Wylie, *Rigidity of virtual quasi-Einstein metrics*, Preprint, 2010.
- [Iv] T. Ivey. *Ricci solitons on compact three-manifolds*, *Differential Geom. Appl.* **3**(1993), no. 4, 301–307.
- [Je] G. R. Jensen, *Homogeneous Einstein spaces of dimension four*, *J. Differential Geom.* **3**(1969), 309–349.
- [JW] D. L. Johnson and L. B. Whitt, *Totally geodesic foliations*, *J. Differential Geom.* **15**(1980), no. 2, 225–235.
- [KK] D.-S. Kim and Y.-H. Kim, *Compact Einstein warped product spaces with nonpositive scalar curvature*, *Proc. Amer. Math. Soc.* **131**(2003), no. 8, 2573–2576.
- [Ko] O. Kowalski, *A classification of Riemannian 3-manifolds with constant principal Ricci curvatures $\rho_1 = \rho_2 \neq \rho_3$* , *Nagoya Math. J.* **132**(1993), 1–36.
- [La1] J. Lauret, *Ricci soliton homogeneous nilmanifolds*, *Math. Ann.* **319**(2001), 715–733.
- [La2] J. Lauret, *Ricci soliton solvmanifolds*, arXiv: 1002.0384, To appear in *J. Reine Angew. Math.*, 2010.
- [La3] J. Lauret, *The Ricci flow for simply connected nilmanifolds*, arXiv: 1004.0946.
- [LPP] H. Lü, Don N. Page, and C. N. Pope, *New inhomogeneous Einstein metrics on sphere bundles over Einstein-Kähler manifolds*, *Phys. Lett. B* **593** (2004), no. 1-4, 218–226.
- [Mi] J. Milnor, *Curvatures of left invariant metrics on Lie groups*, *Advances in Math.* **21**(1976), no. 3, 293–329.
- [MS] O. Munteanu and N. Sesum, *On gradient Ricci solitons*, arXiv: 0910.1105v1.
- [NW] L. Ni and N. Wallach, *On a classification of the gradient shrinking solitons*, *Math. Res. Lett.* **15**(2008), no. 5, 941–955.
- [Per] G. Perelman. *Ricci flow with surgery on three manifolds*, arXiv: math.DG/0303109.
- [Pe] P. Petersen, *Riemannian geometry*, Second edition. *Graduate Texts in Mathematics*, **171**. Springer, New York, 2006.

- [PW1] P. Petersen and W. Wylie, *On gradient Ricci solitons with symmetry*, Proc. Amer. Math. Soc. **137**(2009), no. 6, 2085–2092.
- [PW2] P. Petersen and W. Wylie, *Rigidity of gradient Ricci solitons*, Pacific J. Math. **241**(2009), no. 2, 329–345.
- [PW3] P. Petersen and W. Wylie, *On the classification of gradient Ricci solitons*, Geom. Topol. **14**(2010), no. 4, 2277–2300.
- [Se] H. Seshadri, *On Einstein four-manifolds with S^1 -actions*, Math. Z., **247**(2004), no.3, 487–503.
- [Si] I. M. Singer, *Infinitesimally homogeneous spaces*, Commun. Pure Appl. Math., **13**(1960), 685–697.
- [Vi] C. Villani, *Optimal transport: Old and new*, Grundlehren der Mathematischen Wissenschaften, Vol.338, Springer-Verlag, Berlin, 2009.
- [WW] G. Wei and W. Wylie, *Comparison geometry for the Bakry-Emery Ricci tensor*, J. Differential Geom. **83**(2009), no. 2, 377–405.
- [Zh] Z.-H. Zhang, *Gradient shrinking solitons with vanishing Weyl tensor*, arXiv: 0807.1582.

14 E. PACKER AVE, DEPT. OF MATH, LEHIGH UNIVERSITY, CHRISTMAS-SAUCON HALL, BETHLEHEM, PA 18015

E-mail address: he.chenxu@lehigh.edu

URL: <http://sites.google.com/site/hechenxu/>

520 PORTOLA PLAZA, DEPT OF MATH UCLA, LOS ANGELES, CA 90095

E-mail address: petersen@math.ucla.edu

URL: <http://www.math.ucla.edu/~petersen>

209 S 33RD ST, DEPT. OF MATH, UNIVERSITY OF PENNSYLVANIA, PHILADELPHIA, PA 19104.

E-mail address: wylie@math.upenn.edu

URL: <http://www.math.upenn.edu/~wylie>