

6-7-2006

Near Scale Invariance with Modified Dispersion Relations

Christian Armendariz-Picon

Department of Physics, Syracuse University, Syracuse, NY

Follow this and additional works at: <http://surface.syr.edu/phy>

 Part of the [Physics Commons](#)

Repository Citation

Armendariz-Picon, Christian, "Near Scale Invariance with Modified Dispersion Relations" (2006). *Physics*. Paper 215.
<http://surface.syr.edu/phy/215>

This Article is brought to you for free and open access by the College of Arts and Sciences at SURFACE. It has been accepted for inclusion in Physics by an authorized administrator of SURFACE. For more information, please contact surface@syr.edu.

Near Scale Invariance with Modified Dispersion Relations

C. Armendariz-Picon*

Physics Department, Syracuse University, Syracuse, NY 13244, USA.

(Dated: February 5, 2008)

We describe a novel mechanism to seed a nearly scale invariant spectrum of adiabatic perturbations during a non-inflationary stage. It relies on a modified dispersion relation that contains higher powers of the spatial momentum of matter perturbations. We implement this idea in the context of a massless scalar field in an otherwise perfectly homogeneous universe. The couplings of the field to background scalars and tensors give rise to the required modification of its dispersion relation, and the couplings of the scalar to matter result in an adiabatic primordial spectrum. This work is meant to explicitly illustrate that it is possible to seed nearly scale invariant primordial spectra without inflation, within a conventional expansion history.

I. INTRODUCTION

Inflation has established itself as a key idea in our understanding of the universe [1]. A single stage of accelerated expansion can explain in one shot why the universe is homogeneous and isotropic, spatially flat, what is the origin of the entropy in the universe and why primordial perturbations are adiabatic, Gaussian and nearly scale invariant. For this reason, due to its central role in modern cosmology, it is important to ascertain whether there are alternatives to explain some of the inflationary predictions [2], or whether the universe inevitably had to undergo an inflationary stage [3].

The generation of an adiabatic, Gaussian and scale invariant spectrum of super-horizon perturbations arguably is the most important success of inflation. Indeed, one can reason that it is natural for the universe to begin in a symmetric state, thus accounting for its homogeneity and isotropy. But in an expanding universe with a monotonically decreasing Hubble parameter, there seems to be no way to causally explain the origin of super-horizon perturbations other than inflation, regardless of their spectral index.

In previous works, Eugene Lim and I explored alternative ways of seeding a scale invariant spectrum of primordial perturbations by relaxing some of the assumptions made in our descriptions of the origin of structure. In particular, in [4] we studied whether a contracting sound horizon can lead to the seeding of perturbations, and in [5] I considered the impact of a time-varying scalar field mass on the primordial spectrum. It turns out that in both cases, the generation of scale invariant super-horizon sized primordial perturbations still requires the existence of an inflationary stage, although some of the constraints on its properties can be significantly relaxed.

This work is to some extent a natural continuation of these previous attempts. Our models on the origin of structure are based on extrapolations of low-energy theories to extremely high momenta. However, as we go back in time in the history of the universe, we expect physical

laws to change as typical energies increase. In particular, the dispersion relations used to describe the evolution of structure should receive corrections suppressed by a cut-off scale M , which become important at high energies. A similar phenomenon occurs in crystal lattices; as the wave vector of an electron approaches a Bragg plane, its energy deviates from the conventional quadratic behavior.

In this work, we explore whether modified dispersion relations containing higher powers of the spatial momentum k , can lead to exactly or nearly scale invariant primordial spectra in a non-inflating universe. Ironically, similar dispersion relations have been also considered to study the robustness of inflationary predictions. Specifically, they have been used to analyze whether “trans-Planckian” physics may significantly affect the spectrum of primordial perturbations seeded during inflation [6]. Our goal here is quite different. We would like to determine whether modified dispersion relations may lead to nearly scale invariant spectra in circumstances where we do not expect them to be so.

II. MODIFIED DISPERSION RELATIONS

Consider an expanding, homogeneous, isotropic and spatially flat universe, $ds^2 = a^2(\eta)(-d\eta^2 + d\vec{x}^2)$. Let us assume for simplicity that the universe power-law expands,

$$a(\eta) = \left(\frac{\eta}{\eta_T}\right)^p, \quad (1)$$

where we have arbitrarily set to one the value of the scale factor at an arbitrary time η_T we shall later define. The universe accelerates for $-\infty < p < -1$ and decelerates for $1/2 < p < \infty$, where we have assumed that the dominant energy condition is satisfied. Concretely, a universe with effective equation of state parameter w has

$$p = \frac{2}{1 + 3w}. \quad (2)$$

In Fourier space, the equation of motion of a massless

*Electronic address: armen@phy.syr.edu

scalar field φ in such a universe is

$$v_k'' + \left(\omega^2(k, \eta) - \frac{a''}{a} \right) v_k = 0, \quad (3)$$

where $v_k = a \cdot \varphi_k$, and a prime denotes a derivative with respect to conformal time. In this work we assume that the dispersion relation of the scalar field is

$$\omega^2(k, \eta) = \frac{1}{M^{2n-2}} \frac{k^{2n}}{a^{2m}}, \quad (4)$$

where n and m are constant parameters, and M is a mass scale (recall that we have set $a_T = 1$.) Conventionally, the dispersion relation of a scalar field is $\omega^2 = k^2$, that is, $n = 1$ and $m = 0$. Fluids with a non-constant speed of sound yield a dispersion relation with $n = 1$ and $m \neq 0$. We discuss in Section III how dispersion relations with different values of n and m [7] can arise from generally covariant scalar field actions. In the meantime, the reader can think of equation (4) as a phenomenological description of new physics at momenta k/a that at time η_T are above the cut-off M [8, 9].

A. Initial conditions

The solution of equation (3) with dispersion relation (4) in a universe that expands according to equation (1) is

$$v_k(\eta) = \sqrt{\eta} \left[C_1 H_\nu^{(1)} \left(\frac{\omega \cdot \eta}{mp-1} \right) + C_2 H_\nu^{(2)} \left(\frac{\omega \cdot \eta}{mp-1} \right) \right], \quad (5)$$

where the H_ν are the Hankel functions of the first and second kind, and

$$\nu = \frac{1}{2} \left| \frac{2p-1}{mp-1} \right|. \quad (6)$$

Note that the equation of motion (3) is invariant under $\eta \rightarrow -\eta$. Thus, in order to obtain a solution during inflation (where conformal time is negative), we can simply replace η by $-\eta$. Because the differential equation (3) is second order, its general solution contains two arbitrary integration constants, C_1 and C_2 . Their values are determined by requiring that perturbations be in the adiabatic vacuum at early times. Essentially, this is the only way we know of to specify initial conditions.

The notion of an adiabatic vacuum exists only if the equation of motion admits the approximate solution

$$v_k \approx \frac{1}{\sqrt{2\omega(k, \eta)}} \exp \left(-i \int^\eta \omega(k, \tilde{\eta}) d\tilde{\eta} \right), \quad (7)$$

where ω is a possibly time-dependent frequency [10]. This in fact resembles the vacuum mode functions of a field in Minkowski space. If such an approximate solution exists, perturbations are defined to be in the adiabatic vacuum if they match the approximate solution (7) at early times.

It can be verified that in the limit $\omega \eta \gg 1$, the solution (5) reproduces the adiabatic vacuum (7) if

$$C_1 = \frac{1}{2} \sqrt{\frac{\pi}{mp-1}}, \quad C_2 = 0. \quad (8)$$

Hence, what remains to be verified is whether $\omega \eta \rightarrow \infty$ at early times. Substituting equation (1) into (4) we find that $\omega \eta \gg 1$ amounts to

$$\left(\frac{H}{H_T} \right)^{\frac{mp-1}{p+1}} k^n H_T^{-1} \gg M^{n-1}, \quad (9)$$

where H_T is the value of the Hubble parameter H at the arbitrary time η_T . It is hence clear that the adiabatic condition is satisfied at early times (when $H \rightarrow \infty$) if

$$(1+p)(mp-1) > 0. \quad (10)$$

Therefore, for a conventional dispersion relation, $n = 1$ and $m = 0$, we can specify vacuum initial conditions only when $p < -1$, that is, during inflation. However, if we drop this assumption, we find that an adiabatic solution at early times exists even if $p > 1/2$, that is, even if the universe decelerates. Note that if condition (10) is satisfied, $\omega^2 \cdot \eta^2$ decreases as the universe expands.

B. Horizon Crossing

It is generally argued that only during a stage of inflation it is possible to seed primordial perturbations, regardless of their spectrum. The argument goes as follows. The physical length of a mode is a/k , and the Hubble radius is H^{-1} . Only during a stage of inflation does the length of a mode increase faster than the Hubble radius, that is, only during a stage of inflation can modes begin sub-horizon sized and later become super-horizon sized. In other words, $k^2 \eta^2$ decreases only if the universe accelerates.

Our previous analysis shows the limits of the previous argument (see also [4]). Going back to the equation of motion (3) one realizes that what really matters is whether ω^2 grows faster or slower than a''/a . Only if $\omega^2 \gg a''/a$ at early times there is a chance that we can find solutions that match the adiabatic vacuum (7). We call modes for which $\omega^2 > a''/a$ “short-wavelength” modes, and modes for which $\omega^2 < a''/a$ “long-wavelength” modes. Typically, $a''/a \sim 1/\eta^2$, so a transition between the short and long-wavelength regimes occurs when $\omega^2 \eta^2$ decreases, irrespective of the universe expansion [4]. Moreover, in some (degenerate) cases even this conclusion does not apply. In a radiation dominated universe, a''/a equals zero, and hence all modes (both sub-horizon and super-horizon) are short-wavelength.

Nevertheless, the existence of a short-wavelength regime is not sufficient for the existence of an adiabatic regime, which is the regime needed to specify meaningful

initial conditions. Equation (7) typically is an approximate solution of the equations of motion only if the adiabatic condition $\omega'/\omega^2 \ll 1$ holds [10]. This equation is formally different from the short-wavelength condition $\omega^2 \gg a''/a$, although in practice both are equivalent. Because $\omega'/\omega \sim 1/\eta$ and $a''/a \sim 1/\eta^2$, short-wavelength modes are generally adiabatic. Therefore, the only condition we have to impose on any model for the origin of structure is that $\omega^2 \eta^2$ be monotonically decreasing. Since the period of oscillation of the perturbations (in conformal time) is $1/\omega$, what this means is that the crucial requirement for the causal generation of perturbations is freeze-out ($\omega \eta > 1 \rightarrow \omega \eta < 1$), rather than horizon-crossing ($k \eta > 1 \rightarrow k \eta < 1$).

C. The Power Spectrum

The solution (5) of the equation of motion (3) with adiabatic vacuum initial conditions (8) is

$$v_k = \frac{1}{2} \sqrt{\frac{\pi \eta}{mp-1}} H_\nu^{(1)}(\omega(k, \eta) \cdot \eta). \quad (11)$$

With this exact solution in hand, we are ready to calculate the primordial spectrum of field perturbations. Because, by assumption, φ is in a vacuum state, its expectation value vanishes, $\langle \varphi \rangle = 0$. The two-point function of the field is characterized by the power spectrum

$$\mathcal{P}_\varphi = \frac{k^3}{2\pi^2} |\varphi_k|^2, \quad (12)$$

which is also a measure of the mean squared inhomogeneities of the field on comoving length-scales $1/k$. Using the relation $v_k = a \varphi_k$ and equation (11) we find that at late times ($\omega \eta \ll 1$) the Fourier mode φ_k remains constant for $\text{sgn}[(2p-1)/(mp-1)] = 1$. In this case, the spectrum is given by

$$\mathcal{P}_\varphi = A^2 \cdot \left(\frac{k}{k_T}\right)^{n_s-1}, \quad (13)$$

where the spectral index is

$$n_s - 1 = 3 - n \left| \frac{2p-1}{mp-1} \right|, \quad (14)$$

and the squared amplitude equals

$$A^2 = \frac{\Gamma^2(\nu)}{4\pi^3} \left(\frac{1}{2} \frac{p}{mp-1}\right)^{1-2\nu} H_T^2 \cdot \left(\frac{H_T}{M}\right)^{-2(n-1)\nu}. \quad (15)$$

Note that, in the last equations, k_T is the mode that crosses the Hubble radius at time η_T , $k_T = H_T$. Since, by construction, the equation of motion of the field φ is linear, and because its modes are in a vacuum state, the fluctuations in φ are Gaussian.

The primordial spectrum seeded during a stage of de Sitter inflation ($p = -1$) happens to be scale invariant

for any value of n if $m = n - 1$. This includes the conventional dispersion relation, $n = 1$ and $m = 0$, and the dispersion relation of the ghost condensate [11], $n = 2$ and $m = 1$. But as we see, it is also possible to seed an exactly or nearly scale invariant spectrum of primordial perturbations during a non-inflationary stage of expansion. In particular, many combinations of parameters lead to $n_s \approx 1$ in a non-accelerating universe ($p > 1/2$), given that the only restriction on the model parameters so far is $mp > 1$, from equation (10). And even for $n = 1$ the spectrum seeded during a non-inflationary phase can be scale-invariant if the effective speed of sound is time-dependent and decreasing ($m > 0$), although this realization requires a superluminal *effective* speed of sound [4].

To conclude this section, let us note that current observations actually disfavor an exact scale-invariant spectrum [12] (see however [13]). This does not pose any particular problem in our approach. Although in the realizations we discuss below n has to be an integer, p and m can be arbitrary real numbers. In this case, just as for inflation, departures from scale invariance arise from, e.g., small departures of p from integer values.

D. The scale M

Within a classical description of the universe, nothing prevents us from taking the early-time limit $\eta \rightarrow 0$. However, we expect quantum effects at Planckian energy densities to invalidate our semi-classical description. Hence, rather than requiring that modes be adiabatic as $\eta \rightarrow 0$, we should require that cosmologically relevant modes be in the adiabatic vacuum at the Planckian time at the earliest.

Let us assume that the seeding stage (1) is followed by the conventional period of radiation domination, and let us assume that the transition occurs at time η_T . Requiring the adiabatic condition $\omega \eta \gg 1$ to hold at the Planck time for a mode of the size of our present horizon, we arrive at the upper bound

$$\left(\frac{M}{M_P}\right)^{n-1} \ll 10^{-29n} \left(\frac{H_T}{M_P}\right)^{n/2-p(m+1)/(p+1)}, \quad (16)$$

where $M_P = G^{-1/2} = 10^{19}$ GeV is the Planck mass. We shall later argue that the perturbations in the field φ are converted into adiabatic perturbations at the time of the transition between the seeding stage and radiation domination. Because in our calculation of the spectrum we assumed that the relevant modes are long-wavelength, we have to make sure that this condition is satisfied at the time of the transition. Conservatively requiring than five decades in k above our present horizon be long-wavelength at that time we arrive then at the lower bound

$$\left(\frac{H_T}{M_P}\right)^{n/2-1} \ll \left(\frac{M}{M_P}\right)^{n-1} 10^{24n}. \quad (17)$$

Note that the universe has to be radiation dominated by the time of nucleosynthesis at the latest, which implies the additional constraint $H_T > 10^{-44} M_P$. Conditions (16) and (17) can be simultaneously satisfied with parameters that lead to a nearly scale invariant spectrum. Set for instance $p \approx 2$, $n = 1$ and $m = 1$. In this case, the mass scale M does not enter the equations. Choose in addition $H_T = 10^{-40} M_P$, which amounts to a temperature at the transition $T_T \sim 100$ MeV. All the required conditions are then met. If the parameters are instead, say, $p \approx 1$, $n = 3$ and $m = 2$, the spectrum is again nearly scale invariant, and equations (16) and (17) are satisfied with $M = 10^{-17}$ eV and $H_T = 10^{-40} M_P$. Note the tiny value of the scale M , which is a reflection of the horizon problem; in a non-accelerating universe, our current horizon has a size much larger than the Hubble radius at early times. However, the value of this parameter should be interpreted with care, as the scale M is only defined modulo an appropriately normalized time-dependent function (see below).

III. REALIZATIONS

We still have to answer whether the dispersion relation (4) follows from any generally covariant scalar field theory. What we have to explain is *i*) the origin of the higher powers in spatial momentum k , and *ii*) why these powers are not accompanied by corresponding powers of the scale factor a . As we shall see, ingredient *i*) arises in any low-energy effective theory, while ingredient *ii*) follows from couplings between φ and background scalars and tensors.

A. Effective Field Theories

Let us begin by addressing the origin of the higher powers of spatial momenta in equation (4). Suppose we deal with a low energy effective action with cut-off scale M . The action is expected to contain all possible terms compatible with the symmetries of the theory. Generically no symmetry forbids a term

$$L_{\text{corr}} = \frac{1}{2M^{2n-2}} \varphi \square^n \varphi, \quad (18)$$

where $\square = \nabla^\mu \nabla_\mu$, so we expect the action to contain this operator, in addition to many other similar ones. Upon variation of the action, this term leads to corrections to the equation of motion, $\square \varphi + M^{2-2n} \square^n \varphi = 0$. In a spatially flat Friedman-Robertson-Walker universe with metric (1) these corrections have the structure

$$\square^n \varphi = \frac{1}{a^{2n+1}} \sum_{p+2q+s=2n} c^n{}_{pq} \cdot \eta^{-p} k^{2q} v_k^{(s)}, \quad (19)$$

where $c^n{}_{pq}$ are dimensionless coefficients, a superscript (s) denotes s derivatives with respect to conformal time and we have reintroduced $v_k = a \varphi_k$.

Although the equation of motion then contains the desired term, proportional to k^{2n} , it includes in addition many others that actually dominate over the gradients. Compare for instance $k^{2n} v_k$ with $(k^{2n-2}/\eta^2) v_k$. The first one dominates when $k^2 > \eta^{-2}$, that is, for sub-horizon modes. But without inflation, cosmologically relevant scales are well outside the horizon in the past! Hence, higher-dimensional operators like the one in equation (18) do not lead to the dispersion relation (4).

B. Spatial Projectors

Upon closer inspection of the structure of the dispersion relation (4) one realizes that the key point is the presence of higher-order spatial derivatives in the equation of motion, but the absence at the same time of higher-order time derivatives. This structure naturally appears in the presence of appropriate second rank background tensors [14].

Imagine a spacetime endowed with a symmetric tensor $h_{\mu\nu}$ that projects onto the space orthogonal to a timelike vector field u^μ . By definition, the projector $h_{\mu\nu}$ satisfies $h_{\mu\nu} h^\nu{}_\rho = h_{\mu\rho}$ and $h_{\mu\nu} u^\nu = 0$. In our context, the reader can think of the vector field u^μ as the four velocity of observers for which the universe appears to be isotropic. In conformal time coordinates, this vector has components $u^\mu = (a^{-1}, 0, 0, 0)$, so the projection tensor is $h_{\mu\nu} = a^2 \cdot \text{diag}(0, 1, 1, 1)$. If this is the case, it then follows that, acting on a scalar,

$$D_\mu = h_\mu{}^\nu \nabla_\nu \quad (20)$$

is orthogonal to u^μ and has only spatial components, i. e., is a spatial derivative. On the other hand

$$d_\mu = (\delta_\mu{}^\nu - h_\mu{}^\nu) \nabla_\nu \quad (21)$$

has vanishing spatial components and hence represents a time derivative. Therefore, by combining these two generally covariant derivatives, we can obtain any combination of time and spatial derivatives. In particular,

$$\varphi \cdot (D_\mu D^\mu)^n \varphi = (-1)^n \frac{k^{2n}}{a^{2n}} \varphi_k^2. \quad (22)$$

C. Time functions

Note that in equation (22) every factor of k is accompanied by a corresponding factor of the scale factor a . In order to justify the presence of different powers of a in the dispersion relation (4), a non-vanishing background scalar r is sufficient. Because of the symmetry of the spacetime, this scalar can depend only on time, $r = r(\eta)$, so this is why we shall call r a time function. At this point let us simply assume that $r = a^{2(n-m-1)}$. We shall justify this functional dependence below; what matters

here is that at time η_T , $r = 1$. Given equation (1), it follows that the Lagrangian

$$L = -\frac{1}{2}d_\mu\varphi d^\mu\varphi - \frac{(-1)^n}{2M^{2n-2}} \cdot r \cdot \varphi \cdot (D_\mu D^\mu)^n \varphi \quad (23)$$

yields the equation of motion (3) with dispersion relation given by (4), provided we assume that the fluctuations in r , $g_{\mu\nu}$ and $h_{\mu\nu}$ are negligible; in other words, provided that the universe is highly homogeneous and isotropic. Notice that $n = m + 1$ leads to a simpler model, since in that case r is a constant. In the following, we present possible different origins of the spatial tensor $h_{\mu\nu}$ and the time function r . Similar constructions of the projection tensor involving vector fields [14] are also possible.

a. Non-minimal gravitational couplings The existence of an appropriate tensor $h_{\mu\nu}$ is quite natural in a Friedman-Robertson-Walker universe, and does not imply the breaking of any spacetime symmetry. Consider for instance

$$h_{\mu\nu} = \frac{3(p-1)}{p+1} \frac{R_{\mu\nu}}{R} + \frac{3g_{\mu\nu}}{2(p+1)} \text{ and } r = \left(\frac{R}{R_T} \right)^{-\frac{p(n-m-1)}{p+1}} \quad (24)$$

when $p \neq 1$. The reader can readily verify that the tensor $h_{\mu\nu}$ and the scalar r have the required properties in a universe whose scale factor evolves according to equation (1). This choice of r defines η_T to be the time at which the Ricci scalar equals the free parameter R_T . The presence of the additional curvature terms however leads to modifications in the gravitational equations. Inverse powers of the Ricci scalar have been also postulated as explanations for cosmic acceleration in [15]. Note that the scalar curvature does not necessarily vanish during radiation-domination, say, due to the presence of dark matter particles.

b. Couplings to other fields Even though in our analysis all we had to assume is the appropriate form of the scalar field equation of motion, regardless of the dynamics of gravity, it might be phenomenologically desirable to restrict ourselves to actions where gravity is described by general relativity. If χ is a homogeneous time-evolving scalar field in a FRW universe, then, the projector

$$h_{\mu\nu} = g_{\mu\nu} - \frac{\partial_\mu\chi \partial_\nu\chi}{\partial_\alpha\chi \partial^\alpha\chi} \quad (25)$$

also has the required properties. In addition, if the scalar field χ is rolling down an exponential potential

$$L_\chi = -\frac{1}{2}\partial_\mu\chi\partial^\mu\chi - V_0 \exp\left(-\sqrt{\frac{16\pi(1+p)}{p}} \frac{\chi}{M_P}\right), \quad (26)$$

the time-keeping function

$$r = \exp\left[(n-m-1)\sqrt{\frac{16\pi p}{1+p}} \frac{\chi - \chi_T}{M_P}\right] \quad (27)$$

leads to the desired power of the scale factor in the dispersion relation.¹ In this case, η_T is the time at which the scalar field χ equals χ_T .

IV. METRIC PERTURBATIONS

The mechanism we have presented generates a nearly scale invariant spectrum of perturbations in a subdominant scalar field φ , that is, it leads to *isocurvature* perturbations. Observations however suggest that primordial perturbations are not only nearly scale invariant, but also *adiabatic* and Gaussian. Hence, we still have to explain how these isocurvature perturbations are converted into adiabatic ones. We shall discuss this conversion in the context of the realization of our model where the spatial projectors arises from an evolving and homogeneous scalar field χ .

Imagine that the early universe is dominated by a scalar field χ with an exponential potential, as in equation (26). Assume that this potential has a minimum at χ_{\min} , that is, the potential deviates from an exponential around $\chi \sim \chi_{\min}$. Once the field reaches this minimum, it will start to oscillate and decay into the fields ψ it couples to, thus reheating the universe. Now, assume that the couplings of χ to ψ are not constants, but depend on the fluctuating field φ [16],

$$L_{\text{int}} = -\lambda_0 \cdot \left(1 + f \frac{\varphi}{M}\right) \chi \bar{\psi} \psi, \quad (28)$$

where λ_0 and f are two dimensionless constants. Because the reheating temperature depends on the coupling of χ to matter, the fluctuations in φ lead then to fluctuations in the reheating temperature and the Newtonian potential Φ [5, 16],

$$\Phi \sim \frac{\delta T}{T} \sim f \frac{\varphi}{M}. \quad (29)$$

Combining this equation with our results in equation (13) we find that in cases where the spectrum of φ is nearly scale invariant, the decay of χ results again in a nearly scale independent spectrum of adiabatic perturbations with squared amplitude

$$\mathcal{P}_\Phi \sim f^2 \cdot \left(\frac{H_T}{M}\right)^{\frac{3-n}{n}}. \quad (30)$$

These perturbations are nearly Gaussian (f_{NL} equals a few), provided that the decay rate of χ at the end of the seeding stage is not too high [17].

Cosmic microwave background measurements [12] require $\mathcal{P}_\Phi \sim 10^{-10}$, which fixes the parameter f for given

¹ Since φ vanishes classically, $\varphi_{\text{cl}} \equiv \langle \varphi \rangle = 0$, the terms in the Lagrangian (23) do not alter the χ equation of motion.

H_T and M . Hence, the amplitude of metric perturbations is not uniquely determined by the amplitude of the perturbations in φ . For instance, for $n = 3$ the amplitude of primordial perturbations is just fixed by f , and observations hence require $f \sim 10^{-5}$.

V. HOMOGENEITY

Some inflationary models not only explain the origin of a nearly scale invariant spectrum of primordial perturbations, but also why the universe is homogeneous, isotropic and spatially flat. Whereas our scenario has nothing to say about the spatial geometry of the universe, it can indeed partially address the homogeneity of space.

To that purpose, consider the energy density of an inhomogeneous field φ in an expanding universe. The energy momentum tensor of the scalar field has been calculated in [18]. In the case of a scalar field with action (23) the energy density due to the inhomogeneities in φ is

$$\rho_{\text{inh}} \approx \frac{1}{2} \frac{\varphi_k'^2}{a^2} + \frac{1}{2} \frac{\omega^2}{a^2} \varphi_k^2. \quad (31)$$

We would like to compare the energy density of these inhomogeneities with the total energy density of the universe, $\rho_{\text{tot}} \sim M_P^2/(a^2\eta^2)$. We have seen that as the universe expands, modes leave the short-wavelength regime and become long-wavelength, that is $\omega\eta$ decreases. In the limit $\omega\eta \ll 1$, the solution of the equation of motion is $\varphi_k = \text{const}$. Therefore, the ratio of energy densities approaches

$$\frac{\rho_{\text{inh}}}{\rho_{\text{tot}}} \sim \omega^2 \eta^2 \frac{\varphi_k^2}{M_P^2}. \quad (32)$$

and, hence, just as during inflation, decreases. Clearly, this argument is somewhat heuristic, but it nevertheless suggests that inhomogeneities are “washed-out” by the expansion even if the universe does not inflate.

VI. CONCLUSIONS

In this article we have described a way to seed a nearly scale invariant spectrum of adiabatic primordial pertur-

bations in a non-inflating universe. The key element is the modified dispersion relation of a matter component in the universe. No epoch of contraction is required, and the Hubble parameter does not have to increase at any time during the past history of the universe. For concreteness, we have illustrated this idea with a scalar field, but other realizations of this mechanism might be also possible. In this scalar field context, the modified dispersion relations arises from rather non-generic couplings of the scalar field to background scalars and tensors. Typically, these couplings have to be suppressed by a tiny scale M , a reflection of the horizon problem. Because the spectrum of primordial perturbations is seeded during a non-inflationary stage in an otherwise perfectly homogeneous universe, no gravitational waves are produced. Perturbations are also expected to be non-Gaussian, with an f_{NL} of a few, and thus potentially detectable in the near future. The mechanism does not address the spatial flatness of the universe, but might have the potential to tackle the homogeneity problem.

It is natural to compare our scenario with the inflationary benchmark. Certainly, *some* inflationary models still provide the simplest explanation for many of the features of our universe. But this work should be interpreted from a different perspective anyway. At this stage, it merely shows that the causal generation of the primordial perturbations we observe in our universe does not necessarily require a stage of inflation or an unconventional expansion history.

Acknowledgments

The author thanks the Yukawa Institute for Theoretical Physics and the Max-Planck-Institut für Gravitationsphysik for their kind hospitality. I am also indebted to Shinji Mukohyama for useful conversations and remarks. This work was supported in part by the NSF under grant PHY-0604760.

-
- [1] A. D. Linde, “Particle Physics and Inflationary Cosmology,” arXiv:hep-th/0503203. V. Mukhanov, “Physical Foundations of Cosmology,” (Cambridge University Press, Cambridge, 2005)
- [2] M. Gasperini and G. Veneziano, “Pre - big bang in string cosmology,” *Astropart. Phys.* **1**, 317 (1993) [arXiv:hep-th/9211021]. P. J. Steinhardt and N. Turok,

“Cosmic evolution in a cyclic universe,” *Phys. Rev. D* **65**, 126003 (2002) [arXiv:hep-th/0111098]. S. Hollands and R. M. Wald, “An alternative to inflation,” *Gen. Rel. Grav.* **34**, 2043 (2002) [arXiv:gr-qc/0205058]. A. Nayeri, R. H. Brandenberger and C. Vafa, “Producing a scale-invariant spectrum of perturbations in a Hagedorn phase of string cosmology,” arXiv:hep-th/0511140.

- [3] N. Kaloper, R. Madden and K. A. Olive, “Towards a singularity - free inflationary universe?,” Nucl. Phys. B **452**, 677 (1995) [arXiv:hep-th/9506027]. D. H. Lyth, “The failure of cosmological perturbation theory in the new ekpyrotic scenario,” Phys. Lett. B **526**, 173 (2002) [arXiv:hep-ph/0110007]. L. Kofman, A. Linde and V. F. Mukhanov, “Inflationary theory and alternative cosmology,” JHEP **0210**, 057 (2002) [arXiv:hep-th/0206088].
- [4] C. Armendariz-Picon and E. A. Lim, “Scale invariance without inflation?,” JCAP **0312**, 002 (2003) [arXiv:astro-ph/0307101].
- [5] C. Armendariz-Picon, “Liberating the inflaton from primordial spectrum constraints,” JCAP **0403**, 009 (2004) [arXiv:astro-ph/0310512].
- [6] J. Martin and R. Brandenberger, “On the dependence of the spectra of fluctuations in inflationary cosmology on trans-Planckian physics,” Phys. Rev. D **68**, 063513 (2003) [arXiv:hep-th/0305161].
- [7] R. H. Brandenberger and J. Martin, “The robustness of inflation to changes in super-Planck-scale physics,” Mod. Phys. Lett. A **16**, 999 (2001) [arXiv:astro-ph/0005432].
- [8] W. G. Unruh, “Sonic analog of black holes and the effects of high frequencies on black hole evaporation,” Phys. Rev. D **51**, 2827 (1995).
- [9] S. Corley and T. Jacobson, “Hawking Spectrum and High Frequency Dispersion,” Phys. Rev. D **54**, 1568 (1996) [arXiv:hep-th/9601073].
- [10] N. D. Birrell and P. C. Davies, “Quantum fields in curved space,” (Cambridge University Press, Cambridge, 1982)
- [11] N. Arkani-Hamed, P. Creminelli, S. Mukohyama and M. Zaldarriaga, “Ghost inflation,” JCAP **0404**, 001 (2004) [arXiv:hep-th/0312100].
- [12] D. N. Spergel *et al.*, “Wilkinson Microwave Anisotropy Probe (WMAP) three year results: Implications for cosmology,” arXiv:astro-ph/0603449.
- [13] J. Magueijo and R. D. Sorkin, “Occam’s razor meets WMAP,” arXiv:astro-ph/0604410. D. Parkinson, P. Mukherjee and A. R. Liddle, “A Bayesian model selection analysis of WMAP3,” arXiv:astro-ph/0605003. H. K. Eriksen *et al.*, “A re-analysis of the three-year WMAP temperature power spectrum and likelihood,” arXiv:astro-ph/0606088.
- [14] T. Jacobson and D. Mattingly, “Generally covariant model of a scalar field with high frequency dispersion and the cosmological horizon problem,” Phys. Rev. D **63**, 041502 (2001) [arXiv:hep-th/0009052].
- [15] S. M. Carroll, V. Duvvuri, M. Trodden and M. S. Turner, “Is cosmic speed-up due to new gravitational physics?,” Phys. Rev. D **70**, 043528 (2004) [arXiv:astro-ph/0306438].
- [16] G. Dvali, A. Gruzinov and M. Zaldarriaga, “A new mechanism for generating density perturbations from inflation,” Phys. Rev. D **69**, 023505 (2004) [arXiv:astro-ph/0303591].
- [17] M. Zaldarriaga, “Non-Gaussianities in models with a varying inflaton decay rate,” Phys. Rev. D **69**, 043508 (2004) [arXiv:astro-ph/0306006].
- [18] M. Lemoine, M. Lubo, J. Martin and J. P. Uzan, “The stress-energy tensor for trans-Planckian cosmology,” Phys. Rev. D **65**, 023510 (2002) [arXiv:hep-th/0109128].