

7-31-2011

Strong Approximation of Homeomorphisms of Finite Dirichlet Energy

Tadeusz Iwaniec

Syracuse University and University of Helsinki

Leonid V. Kovalev

Syracuse University

Jani Onninen

Syracuse University

Follow this and additional works at: <https://surface.syr.edu/mat>

 Part of the [Mathematics Commons](#)

Recommended Citation

Iwaniec, Tadeusz; Kovalev, Leonid V.; and Onninen, Jani, "Strong Approximation of Homeomorphisms of Finite Dirichlet Energy" (2011). *Mathematics - Faculty Scholarship*. 56.

<https://surface.syr.edu/mat/56>

This Article is brought to you for free and open access by the Mathematics at SURFACE. It has been accepted for inclusion in Mathematics - Faculty Scholarship by an authorized administrator of SURFACE. For more information, please contact surface@syr.edu.

STRONG APPROXIMATION OF HOMEOMORPHISMS OF FINITE DIRICHLET ENERGY

TADEUSZ IWANIEC, LEONID V. KOVALEV, AND JANI ONNINEN

ABSTRACT. Let $\mathbb{X} \subset \mathbb{C}$ and $\mathbb{Y} \subset \mathbb{C}$ be Jordan domains of the same finite connectivity, \mathbb{Y} being inner chordarc regular (such are Lipschitz domains). Every homeomorphism $h: \mathbb{X} \rightarrow \mathbb{Y}$ in the Sobolev space $\mathcal{W}^{1,2}$ extends to a continuous map $h: \bar{\mathbb{X}} \rightarrow \bar{\mathbb{Y}}$. We prove that there exist homeomorphisms $h_k: \bar{\mathbb{X}} \rightarrow \bar{\mathbb{Y}}$ which converge to h uniformly and in $\mathcal{W}^{1,2}(\mathbb{X}, \mathbb{Y})$. The problem of approximation of Sobolev homeomorphisms, raised by J. M. Ball and L. C. Evans, is deeply rooted in a study of energy-minimal deformations in nonlinear elasticity. The new feature of our main result is that approximation takes place also on the boundary, where the original map need not be a homeomorphism.

1. INTRODUCTION

Throughout this text \mathbb{X} and \mathbb{Y} are finitely connected Jordan domains in the complex plane $\mathbb{C} \simeq \mathbb{R}^2$. We shall consider orientation preserving homeomorphisms $h: \mathbb{X} \xrightarrow{\text{onto}} \mathbb{Y}$ of finite Dirichlet energy

$$\mathcal{E}_x[h] = \iint_{\mathbb{X}} |Dh(z)|^2 dx dy < \infty, \quad z = x + iy$$

where $|Dh|$ stands for the Hilbert-Schmidt norm of the derivative matrix Dh . Such a mapping has a continuous extension $h: \bar{\mathbb{X}} \xrightarrow{\text{onto}} \bar{\mathbb{Y}}$ which is not necessarily a homeomorphism [15]. In this paper we show that h can be strongly approximated by homeomorphisms between closed domains, provided \mathbb{Y} is inner chordarc regular, see Definition 2.3. In particular, \mathbb{Y} can be a Lipschitz domain.

Theorem 1.1. *Let \mathbb{X} and \mathbb{Y} be finitely connected Jordan domains, \mathbb{Y} being inner chordarc. Let $h: \mathbb{X} \xrightarrow{\text{onto}} \mathbb{Y}$ be a homeomorphism in the Sobolev space $\mathcal{W}^{1,2}(\mathbb{X}, \mathbb{Y})$. Then:*

- (a) *there exist homeomorphisms $h_k: \bar{\mathbb{X}} \xrightarrow{\text{onto}} \bar{\mathbb{Y}}$, $k = 1, 2, \dots$, that converge to $h: \bar{\mathbb{X}} \xrightarrow{\text{onto}} \bar{\mathbb{Y}}$ uniformly and strongly in $\mathcal{W}^{1,2}(\mathbb{X})$.*
- (b) *Moreover, for every compact subset $\mathbb{G} \subset \mathbb{X}$, we have $h_k \equiv h$ on \mathbb{G} , provided $k = k(\mathbb{G})$ is sufficiently large.*
- (c) *If, in addition, $h: \mathfrak{X} \rightarrow \partial\mathbb{Y}$ is injective on a compact subset $\mathfrak{X} \subset \partial\mathbb{X}$, then h_k can be chosen so that $h_k \equiv h$ on \mathfrak{X} .*

2010 *Mathematics Subject Classification.* Primary 30E10; Secondary 46E35, 58E20.

Key words and phrases. Approximation, Homeomorphisms, Dirichlet energy, Lipschitz domain, Inner chordarc domain, harmonic mapping.

T. Iwaniec was supported by the NSF grant DMS-0800416 and the Academy of Finland project 1128331. L. Kovalev was supported by the NSF grant DMS-0968756. J. Onninen was supported by the NSF grant DMS-1001620.

The motivation for part (c) comes from variational problems for mappings between quadrilaterals, i.e., Jordan domains with four distinguished points which constitute the injectivity set \mathfrak{X} . We will pursue such applications of Theorem 1.1 elsewhere.

Let us compare Theorem 1.1 to the known results in this field. The problem of approximation of Sobolev homeomorphisms was raised by J. M. Ball and L. C. Evans in 1990s [1, 2]. There have been several recent advances in this direction [4, 8, 13, 14, 18]. In particular, in [13, 14] we proved that any $\mathscr{W}^{1,p}$ -homeomorphism $h: \mathbb{U} \xrightarrow{\text{onto}} \mathbb{V}$ between open subsets of \mathbb{R}^2 can be approximated in the $\mathscr{W}^{1,p}$ norm by diffeomorphisms of these open subsets. The new feature of Theorem 1.1 is that approximation takes place also on the boundary, where the original map need not be a homeomorphism. This is why Theorem 1.1 imposes regularity assumptions on the boundaries of \mathbb{X} and \mathbb{Y} .

Combining Theorem 1.1 with [13, Theorem 1.2] yields the following result.

Corollary 1.2. *If $h: \mathbb{X} \xrightarrow{\text{onto}} \mathbb{Y}$ is a \mathcal{C}^∞ -diffeomorphism in $\mathscr{W}^{1,2}(\mathbb{X})$ then, in addition to all properties listed in Theorem 1.1, the mappings $h_k: \mathbb{X} \xrightarrow{\text{onto}} \mathbb{Y}$ can also be found as \mathcal{C}^∞ -diffeomorphisms. If h is only a homeomorphism, such an approximation by diffeomorphisms is still available, except for the property (b).*

One of the fundamental problems in topology is to approximate continuous mappings by homeomorphisms. The approximation procedures, still only partially understood, have led topologists to the concept of monotone mappings [19] and somewhat subtle concept of cellular mappings [5]. We refer the interested reader to [17, 23]. In the mathematical theory of hyperelasticity, on the other hand, we are concerned with the energy-minimal deformations $h: \mathbb{X} \xrightarrow{\text{onto}} \mathbb{Y}$, so having additional Sobolev type regularity [3, 6, 20]. However, very often the injectivity of the energy minimal mappings is lost, though they enjoy some features of homeomorphisms, like monotonicity. In particular, the question of approximation of a monotone mapping in the Sobolev space $\mathscr{W}^{1,2}(\mathbb{X})$ by homeomorphisms $h_k: \overline{\mathbb{X}} \xrightarrow{\text{onto}} \overline{\mathbb{Y}}$ gains interest in the mathematical models of elasticity. A novelty in these directions is the following corollary of Theorem 1.1 and [7, Theorem 1.6].

Corollary 1.3. *Suppose $h: \mathbb{X} \xrightarrow{\text{onto}} \mathbb{Y}$ lies in the Sobolev space $W^{1,2}(\mathbb{X})$ and extends to a continuous monotone map $h: \overline{\mathbb{X}} \xrightarrow{\text{onto}} \overline{\mathbb{Y}}$. Then there exist homeomorphisms $h_k: \overline{\mathbb{X}} \xrightarrow{\text{onto}} \overline{\mathbb{Y}}$, $k = 1, 2, \dots$, that converge to $h: \overline{\mathbb{X}} \xrightarrow{\text{onto}} \overline{\mathbb{Y}}$ uniformly and strongly in $\mathscr{W}^{1,2}(\mathbb{X})$.*

We conclude this introduction with two open questions.

Question 1.4. Does Theorem 1.1 remain valid when target \mathbb{Y} is an arbitrary finitely connected Jordan domain?

Question 1.5. Can Theorem 1.1 be extended to Sobolev spaces $\mathscr{W}^{1,p}$, $p \in (1, \infty)$? Or to dimensions $n > 2$?

2. PRELIMINARIES

Royden Algebra. The Royden algebra $\mathscr{R}(\mathbb{X})$ consists of continuous functions $g: \overline{\mathbb{X}} \rightarrow \mathbb{C}$ which have finite energy. The norm is given by

$$\|g\|_{\mathscr{R}(\mathbb{X})} = \sup_{z \in \mathbb{X}} |g(z)| + \left(\iint_{\mathbb{X}} |Dg(z)|^2 dx dy \right)^{\frac{1}{2}}$$

Let $\mathcal{R}_0(\mathbb{X})$ denote the completion of $\mathcal{C}_0^\infty(\mathbb{X})$ in this norm. Any conformal mapping $\varphi: \mathbb{X}' \xrightarrow{\text{onto}} \mathbb{X}$ between Jordan domains induces an isometry $\varphi^\#: \mathcal{R}(\mathbb{X}) \xrightarrow{\text{onto}} \mathcal{R}(\mathbb{X}')$ by the rule $\varphi^\#(h) = h \circ \varphi$. Thus the domain of definition of h is permitted to be changed by any conformal transformation. The following observation will allow us to transform the target.

Lemma 2.1. *Let $h, h_k: \mathbb{X} \xrightarrow{\text{onto}} \mathbb{Y}$ be homeomorphisms in $\mathcal{R}(\mathbb{X})$ and $\Phi: \mathbb{Y} \xrightarrow{\text{onto}} \mathbb{Y}'$ be a \mathcal{C}^1 -diffeomorphism that extends to a homeomorphism $\Phi: \overline{\mathbb{Y}} \xrightarrow{\text{onto}} \overline{\mathbb{Y}'}$. Assume that both the gradient matrix $D\Phi$ and its inverse $(D\Phi)^{-1}$ are bounded in \mathbb{Y} . Then $\Phi \circ h_k \rightarrow \Phi \circ h$ in $\mathcal{R}(\mathbb{X})$ if and only if $h_k \rightarrow h$ in $\mathcal{R}(\mathbb{X})$.*

Our primary appliance for strong approximation in Theorem 1.1 will be local harmonic replacements near the boundaries of \mathbb{X} . We will rely on a well-known fact, see [13].

Lemma 2.2. *In a finitely connected Jordan domain $\mathbb{X} \subset \mathbb{C}$, we consider a function $g \in \mathcal{R}(\mathbb{X})$. Let $h: \overline{\mathbb{X}} \rightarrow \mathbb{C}$ denote the continuous harmonic extension of the boundary map $g: \partial\mathbb{X} \rightarrow \mathbb{C}$ into \mathbb{X} . Then $h \in g + \mathcal{R}_0(\mathbb{X})$. Moreover,*

$$(2.1) \quad 0 \leq \mathcal{E}_x[g - h] = \mathcal{E}_x[g] - \mathcal{E}_x[h] \leq \mathcal{E}_x[g].$$

Inner Chordarc Domains.

Definition 2.3. A finitely connected Jordan domain \mathbb{Y} is *inner chordarc* if there exists a constant C with the following property. Suppose that a, b belong to the same boundary component of \mathbb{Y} and $\gamma \subset \mathbb{Y}$ is an open Jordan arc with endpoints at a and b . Then the shortest connection from a to b along $\partial\mathbb{Y}$ has length at most $C \cdot \text{length}(\gamma)$.

Inner chordarc domains were studied in Geometric Function Theory since 1980s [11, 12, 16, 21, 24, 25, 26]. They are more general than Lipschitz domains. For instance they allow inward cusps and logarithmic spiraling, see Figure 1.

Theorem 2.4. [25] *A simply connected Jordan domain Ω is inner chordarc if and only if there exists a \mathcal{C}^1 -diffeomorphism F from Ω onto the unit disk \mathbb{D} that extends to a homeomorphism $F: \overline{\Omega} \xrightarrow{\text{onto}} \overline{\mathbb{D}}$ such that both gradient matrices DF and $(DF)^{-1}$ are bounded in Ω .*

We remark that [25, Theorem 3.8] is stated in terms of BLD homeomorphisms, which are not necessarily smooth. However, the mapping constructed in its proof is a diffeomorphism.




FIGURE 1. Inner chordarc domains

Figure 1 illustrates two such mappings from the unit disk onto a non-Lipschitz domain: $\Phi(z) = |z - 1|^{4i}(z - 1)$ (the image is a spiral domain) and $\Psi(z) = (z + 1)^2/|z + 1|$ (the image contains an inward cusp).

Monotone mappings. Recall that a continuous map $h : \overline{\mathbb{X}} \xrightarrow{\text{onto}} \overline{\mathbb{Y}}$ is monotone if the preimage of every continuum (connected compact set) in $\overline{\mathbb{Y}}$ is a continuum in $\overline{\mathbb{X}}$. A point $y_0 \in \overline{\mathbb{Y}}$ is said to be a *simple value* of h if its preimage $h^{-1}\{y_0\}$ is a single point in $\overline{\mathbb{X}}$.

We need an elementary lemma.

Lemma 2.5. *Let $h : \mathbb{X} \xrightarrow{\text{onto}} \mathbb{Y}$ be a homeomorphism between ℓ -connected Jordan domains and suppose that it extends continuously up to the boundary. Then the extended map $h : \overline{\mathbb{X}} \xrightarrow{\text{onto}} \overline{\mathbb{Y}}$ is monotone. Furthermore, the inverse map $h^{-1} : \mathbb{Y} \xrightarrow{\text{onto}} \mathbb{X}$ extends continuously up to simple values of h which are virtually all points in $\partial\mathbb{Y}$, except for a countable number of them.*

Every homeomorphism with finite Dirichlet energy has a continuous extension to the boundary [15]. Precisely,

Theorem 2.6. *Every finite energy homeomorphism $h : \mathbb{X} \xrightarrow{\text{onto}} \mathbb{Y}$ between ℓ -connected Jordan domains extends to a continuous map between the closures, again denoted by $h : \overline{\mathbb{X}} \xrightarrow{\text{onto}} \overline{\mathbb{Y}}$. This map is monotone, though the inverse $h^{-1} : \mathbb{Y} \xrightarrow{\text{onto}} \mathbb{X}$ may not admit continuous extension to the closure of \mathbb{Y} , unless it also has finite energy.*

Harmonic Homeomorphisms. We shall make use of the following strong version of the Radó-Kneser-Choquet theorem, see [10, §3.2].

Theorem 2.7 (Radó-Kneser-Choquet). *Let $\mathbb{U} \subset \mathbb{C}$ be a simply connected Jordan domain and $\Omega \subset \mathbb{C}$ a bounded convex domain. Suppose we are given a continuous monotone map $h : \partial\mathbb{U} \xrightarrow{\text{onto}} \partial\Omega$, not necessarily a homeomorphism. Then its continuous harmonic extension, denoted by $H : \overline{\mathbb{U}} \rightarrow \mathbb{C}$, defines a \mathcal{C}^∞ -diffeomorphism $H : \mathbb{U} \xrightarrow{\text{onto}} \Omega$.*

3. PROOF OF THEOREM 1.1



FIGURE 2. A finitely connected inner chordarc domain, not Lipschitz

Each boundary $\partial\mathbb{X}$ and $\partial\mathbb{Y}$ consists of ℓ disjoint Jordan curves. We reserve the notation,

$$\mathfrak{X}_1, \mathfrak{X}_2, \dots, \mathfrak{X}_\ell, \quad \text{for the components of } \partial\mathbb{X}$$

$$\Upsilon_1, \Upsilon_2, \dots, \Upsilon_\ell, \quad \text{for the components of } \partial\mathbb{Y}$$

The components are numbered so that h sends \mathfrak{X}_ν onto Υ_ν for $\nu = 1, \dots, \ell$. Figure 2 illustrates a domain \mathbb{Y} that satisfies the assumptions of the theorem.

We recall that the boundary map $h : \partial\mathbb{X} \xrightarrow{\text{onto}} \partial\mathbb{Y}$ is injective on a compact subset $\mathfrak{X} \subset \partial\mathbb{X} = \mathfrak{X}_1 \cup \mathfrak{X}_2 \cup \dots \cup \mathfrak{X}_\ell$, possibly empty. Let us fix an $\varepsilon > 0$ and a compact $\mathbb{G} \subset \mathbb{X}$. We need to construct a homeomorphism $h_\varepsilon : \overline{\mathbb{X}} \xrightarrow{\text{onto}} \overline{\mathbb{Y}}$ of Sobolev class $\mathscr{W}^{1,2}(\mathbb{X}, \mathbb{Y})$ which coincides with h on $\mathfrak{X} \cup \mathbb{G}$ and

$$(3.1) \quad \|h_\varepsilon - h\|_{\mathscr{A}(\mathbb{X})} \preccurlyeq \varepsilon$$

Hereafter the symbol \preccurlyeq indicates that the inequality holds with an *implied* multiplicative constant. The implied constants will vary from line to line but remain **independent of** ε as long as ε is sufficiently small.

To prove (3.1) we shall set up a chain of homeomorphisms $h^0, h^1, \dots, h^\ell : \mathbb{X} \xrightarrow{\text{onto}} \mathbb{Y}$ in the Sobolev space $\mathscr{W}^{1,2}(\mathbb{X}, \mathbb{Y})$ whose continuous extensions, still denoted by $h^0, h^1, \dots, h^\ell : \overline{\mathbb{X}} \xrightarrow{\text{onto}} \overline{\mathbb{Y}}$, satisfy

- $h^0 \equiv h$
- $h^1 : \mathfrak{X}_1 \cup \mathfrak{X} \rightarrow \partial\mathbb{Y}$ is injective,
 $h^1 \equiv h$ on $\mathfrak{X} \cup \mathbb{G}$,
 $\|h^1 - h^0\|_{\mathscr{A}(\mathbb{X})} \preccurlyeq \varepsilon$
- For $2 \leq \nu \leq \ell$,
 $h^\nu : \mathfrak{X}_1 \cup \dots \cup \mathfrak{X}_\nu \cup \mathfrak{X} \rightarrow \partial\mathbb{Y}$ is injective,
 $h^\nu \equiv h^{\nu-1}$ on $\mathfrak{X}_1 \cup \dots \cup \mathfrak{X}_{\nu-1} \cup \mathfrak{X} \cup \mathbb{G}$,
 $\|h^\nu - h^{\nu-1}\|_{\mathscr{A}(\mathbb{X})} \preccurlyeq \varepsilon$

Thus the final term h^ℓ works for the desired homeomorphism $h_\varepsilon : \overline{\mathbb{X}} \xrightarrow{\text{onto}} \overline{\mathbb{Y}}$, by the triangle inequality. We proceed by induction on ν . The induction begins with h^0 , which is obvious. Suppose we are given the map $h^\nu : \overline{\mathbb{X}} \xrightarrow{\text{onto}} \overline{\mathbb{Y}}$ for some $0 \leq \nu < \ell$. The construction of $h^{\nu+1} : \overline{\mathbb{X}} \xrightarrow{\text{onto}} \overline{\mathbb{Y}}$ will be made in 5 steps.

Step I. *Transition to the case when $\Upsilon_{\nu+1} = \mathbb{T}$ is the outer boundary.* To make this transition rigorous, let us perform the following transformations of $\overline{\mathbb{Y}}$. First, we reduce ourselves to the case in which $\Upsilon_{\nu+1}$ is the outer boundary of \mathbb{Y} by applying an inversion if necessary. Note that such an inversion is a diffeomorphism in a neighborhood of $\overline{\mathbb{Y}}$. Once $\Upsilon_{\nu+1}$ is the outer boundary of \mathbb{Y} we apply Theorem 2.4 to transform the bounded component of $\mathbb{C} \setminus \Upsilon_{\nu+1}$, denoted by Ω , onto the unit disk. Let this transformation be denoted by $F : \Omega \xrightarrow{\text{onto}} \mathbb{D}$. This map extends to a homeomorphism $F : \overline{\Omega} \xrightarrow{\text{onto}} \overline{\mathbb{D}}$, and has both matrix functions DF and $(DF)^{-1}$ continuous and bounded in Ω . Let $\mathbb{Y}' = F(\mathbb{Y})$. By virtue of Lemma 2.1 the composition with the mapping $F : \mathbb{Y} \xrightarrow{\text{onto}} \mathbb{Y}'$ transforms converging sequences of mappings $h_k : \mathbb{X} \xrightarrow{\text{onto}} \mathbb{Y}$ into converging sequences of mappings $F \circ h_k : \mathbb{X} \xrightarrow{\text{onto}} \mathbb{Y}'$. The inverse F^{-1} has the same property; therefore we can work with the target \mathbb{Y}' instead of \mathbb{Y} . In what follows we simply assume that $\Upsilon_{\nu+1} = \mathbb{T}$ is the outer boundary of \mathbb{Y} .

Step II. Harmonic Replacements Near $\mathfrak{X}_{\nu+1}$. The idea is to alter h^ν in a thin neighborhood of $\mathfrak{X}_{\nu+1}$ to gain piecewise harmonicity therein. In this step we change neither the boundary map $h^\nu : \partial\mathbb{X} \xrightarrow{\text{onto}} \partial\mathbb{Y}$ nor the values of h^ν on the given compact $\mathbb{G} \subset \mathbb{X}$.

Recall from Lemma 2.5 that all but a countable number of points in \mathbb{T} are simple values of the map $h^\nu : \mathfrak{X}_{\nu+1} \xrightarrow{\text{onto}} \mathbb{T}$. They are dense, so one can partition \mathbb{T} into arbitrarily small *closed* circular arcs whose ends are simple values of h^ν ,

$$\mathbb{T} = \bigcup_{\kappa=1}^N \mathcal{C}_\kappa, \quad \text{diam } \mathcal{C}_\kappa \leq \varepsilon < 2, \quad \text{for all } \kappa = 1, \dots, N$$

Let \mathbb{S}_κ denote the open region between the arc \mathcal{C}_κ and the closed line interval \mathbf{I}_κ connecting the endpoints of \mathcal{C}_κ . We call \mathbf{I}_κ the *base* of the *segment* \mathbb{S}_κ . We require the partition of \mathbb{T} to be fine enough so that the compact set $h^\nu(\mathbb{G})$ intersects none of the segments \mathbb{S}_κ , $\kappa = 1, \dots, N$. One further restriction on the partition comes from the following observation: the finer the partition the closer to \mathbb{T} are the segments \mathbb{S}_κ . Since $h^\nu : \mathbb{X} \xrightarrow{\text{onto}} \mathbb{Y}$ is a homeomorphism, it follows that the preimages of \mathbb{S}_κ under h^ν , denoted by \mathbb{X}_κ , can be as close to $\mathfrak{X}_{\nu+1}$ as we wish. In particular, we may ensure that

$$(3.2) \quad \sum_{\kappa=1}^N \iint_{\mathbb{X}_\kappa} |Dh^\nu|^2 \leq \varepsilon^2$$

That is all what we require to determine the partition of \mathbb{T} . This partition will remain fixed for the rest of the proof. Now we observe that each \mathbb{X}_κ is a simply connected Jordan domain. Its boundary consists of two closed Jordan arcs with common endpoints. The one in $\mathfrak{X}_{\nu+1}$ is denoted by $\Gamma_\kappa = \overline{\mathbb{X}_\kappa} \cap \mathfrak{X}_{\nu+1}$ and the open arc in \mathbb{X} is denoted by $\gamma_\kappa \stackrel{\text{def}}{=} \partial\mathbb{X}_\kappa \cap \mathbb{X}$. It is at this point that we take advantage of the condition that the endpoints of \mathcal{C}_κ are simple values of h^ν . This condition implies that the inverse map $(h^\nu)^{-1} : \mathbf{I}_\kappa \xrightarrow{\text{onto}} \overline{\gamma}_\kappa$ is a homeomorphism. On the other hand the preimage $\Gamma_\kappa = (h^\nu)^{-1}(\mathcal{C}_\kappa)$ is a closed arc, because of monotonicity of $h^\nu : \partial\mathbb{X} \xrightarrow{\text{onto}} \partial\mathbb{Y}$. Therefore, the open Jordan arc $\gamma_\kappa \subset \mathbb{X}$ and the closed arc $\Gamma_\kappa \subset \mathfrak{X}_{\nu+1}$ form a closed Jordan curve; precisely, the boundary of \mathbb{X}_κ . In summary,

- $\partial\mathbb{X}_\kappa = \gamma_\kappa \cup \Gamma_\kappa$, $h^\nu : \mathbb{X}_\kappa \xrightarrow{\text{onto}} \mathbb{S}_\kappa$,
- $h^\nu : \partial\mathbb{X}_\kappa \xrightarrow{\text{onto}} \partial\mathbb{S}_\kappa$ is continuous and monotone
- This latter boundary map is injective on the compact subset $\mathfrak{X}_{\nu+1}^\kappa \stackrel{\text{def}}{=} (\mathfrak{X} \cap \mathfrak{X}_{\nu+1}) \cup \overline{\gamma}_\kappa \subset \partial\mathbb{X}_\kappa$

Now we appeal to Theorem 2.7 of Radó-Kneser-Choquet which allows us to replace $h^\nu : \mathbb{X}_\kappa \xrightarrow{\text{onto}} \mathbb{S}_\kappa$ by the harmonic extension of its boundary map $h^\nu : \partial\mathbb{X}_\kappa \xrightarrow{\text{onto}} \partial\mathbb{S}_\kappa$. We need to introduce, for a little while, more notation.

- $h_\kappa^\nu : \overline{\mathbb{X}_\kappa} \xrightarrow{\text{onto}} \overline{\mathbb{S}_\kappa}$ — harmonic extension of $h^\nu : \partial\mathbb{X}_\kappa \xrightarrow{\text{onto}} \partial\mathbb{S}_\kappa$
- $\mathbf{h}^\nu : \mathbb{X} \xrightarrow{\text{onto}} \mathbb{Y}$ — homeomorphism of class $\mathscr{H}^{1,2}(\mathbb{X}, \mathbb{Y})$, defined by

$$\mathbf{h}^\nu = \begin{cases} h_\kappa^\nu & \text{on } \mathbb{X}_\kappa, \quad \kappa = 1, 2, \dots, N \\ h^\nu & \text{otherwise} \end{cases}$$

The continuous extension $\mathbf{h}^\nu : \overline{\mathbb{X}} \xrightarrow{\text{onto}} \overline{\mathbb{Y}}$ agrees with h^ν on $\partial\mathbb{X}$ and on the compact $\mathbb{G} \subset \mathbb{X}$ as well.

Let us estimate the difference $h^\nu - \mathbf{h}^\nu$ in the norm of Royden algebra. First we find that

$$\|h^\nu - \mathbf{h}^\nu\|_{\mathcal{C}^\infty(\mathbb{X})} \leq \sup_{1 \leq \kappa \leq N} \|h^\nu - \mathbf{h}^\nu\|_{\mathcal{C}^\infty(\mathbb{X}_\kappa)} \leq \sup_{1 \leq \kappa \leq N} \text{diam } \mathbb{S}_\kappa \leq \varepsilon$$

Secondly, in view of (2.1), we see that

$$\mathcal{E}_x[h^\nu - \mathbf{h}^\nu] = \sum_{\kappa=1}^N \mathcal{E}_{x_\kappa}[h^\nu - h_{j_\kappa}^\nu] \leq \sum_{\kappa=1}^N \mathcal{E}_{x_\kappa}[h^\nu] \leq \varepsilon^2$$

by (3.2). Hence

$$\|h^\nu - \mathbf{h}^\nu\|_{\mathcal{A}(\mathbb{X})} \preceq \varepsilon$$

Summarizing, the construction of $h^{\nu+1}$ in an ε proximity to h^ν will be done once the similar construction is in hand for \mathbf{h}^ν . In what follows, instead of using \mathbf{h}^ν , we assume that the original map h^ν was already harmonic in every \mathbb{X}_κ . This simplifies writing and causes no loss of generality.

Step III. Reduction of the domain to the unit disk. The idea is to construct for each $\kappa = 1, 2, \dots, N$ a sequence of homeomorphisms $h_j^{\kappa, \nu} : \overline{\mathbb{X}_\kappa} \xrightarrow{\text{onto}} \overline{\mathbb{S}_\kappa}$, $j = 1, 2, \dots$, that converge to $h^\nu : \overline{\mathbb{X}_\kappa} \xrightarrow{\text{onto}} \overline{\mathbb{S}_\kappa}$ uniformly and in $\mathcal{W}^{1,2}(\mathbb{X}_\kappa, \mathbb{S}_\kappa)$. In addition to that, we require that each $h_j^{\kappa, \nu}$ agrees with h^ν on the compact subset $\mathfrak{X}_{\nu+1}^\kappa \stackrel{\text{def}}{=} (\mathfrak{X} \cap \mathfrak{X}_{\nu+1}) \cup \overline{\gamma}_\kappa \subset \partial \mathbb{X}_\kappa$. Recall that the boundary map $h^\nu : \partial \mathbb{X}_\kappa \xrightarrow{\text{onto}} \partial \mathbb{S}_\kappa$ is injective on $\mathfrak{X}_{\nu+1}^\kappa$. Once this is done, the construction of $h^{\nu+1}$ will be completed in the following way: for each κ we choose and fix $j = j_\kappa$ sufficiently large so that

$$\|h^\nu - h_{j_\kappa}^{\kappa, \nu}\|_{\mathcal{A}(\mathbb{X}_\kappa)} \leq \varepsilon$$

Then we replace each $h^\nu : \overline{\mathbb{X}_\kappa} \xrightarrow{\text{onto}} \overline{\mathbb{S}_\kappa}$ by $h_{j_\kappa}^{\kappa, \nu} : \overline{\mathbb{X}_\kappa} \xrightarrow{\text{onto}} \overline{\mathbb{S}_\kappa}$ to obtain the desired map

$$h^{\nu+1} \stackrel{\text{def}}{=} \begin{cases} h_{j_\kappa}^{\kappa, \nu} & \text{on } \mathbb{X}_\kappa, \quad \kappa = 1, 2, \dots, n \\ h^\nu & \text{otherwise} \end{cases}$$

Thus we are reduced to finding the sequence $h_j^{\kappa, \nu} : \overline{\mathbb{X}_\kappa} \xrightarrow{\text{onto}} \overline{\mathbb{S}_\kappa}$, $j = 1, 2, \dots$. Before proceeding to somewhat involved computation we need to simplify the domain and the target of $h^\nu : \overline{\mathbb{X}_\kappa} \xrightarrow{\text{onto}} \overline{\mathbb{S}_\kappa}$. Since the problem is clearly unaffected by a rotation of the target (harmonicity of the map is not compromised), we may confine ourselves to the segment of the form

$$\mathbb{S}_\kappa = \mathbb{S} \stackrel{\text{def}}{=} \{ \xi : |\xi| < 1 : \cos \omega < \Re \xi < 1 \}, \quad \text{for some } 0 < \omega < \frac{\pi}{2}$$

Thus its arc \mathcal{C} is $\{ \xi = e^{i\phi} : -\omega \leq \phi \leq \omega \}$, the base \mathbf{I} is $\{ \xi = \cos \omega + i\tau : -\sin \omega \leq \tau \leq \sin \omega \}$ and the corners are $\xi^+ = e^{i\omega}$, $\xi^- = e^{-i\omega}$. Regarding the domain \mathbb{X}_κ , it is legitimate to conformally transform it onto the unit disk \mathbb{D} ; any conformal mapping between two Jordan domains induces an isometry of their Royden algebras. Thus we consider a conformal map $\chi : \mathbb{D} \xrightarrow{\text{onto}} \mathbb{X}_\kappa$ and the pullback $f \stackrel{\text{def}}{=} h^\nu \circ \chi : \mathbb{D} \xrightarrow{\text{onto}} \mathbb{S}$. Recall that χ is a homeomorphism up to the boundaries. One extra condition turns out to be useful; namely, we may choose χ to be normalized at three boundary points so that the map $f \stackrel{\text{def}}{=} h^\nu \circ \chi$ satisfies,

$$(3.3) \quad f(e^{i\omega}) = e^{i\omega}, \quad f(e^{-i\omega}) = e^{-i\omega}, \quad f(-1) = \cos \omega$$

The first two values of f are the endpoints of the base $\mathbf{I} \subset \partial\mathbb{S}$ and the last one is its midpoint. Let us state clearly what we aim to show to complete the proof of Theorem 1.1.

Proposition 3.1. *Let $f : \mathbb{D} \xrightarrow{\text{onto}} \mathbb{S}$ be an orientation preserving harmonic homeomorphism of finite energy and let its continuous extension $f : \overline{\mathbb{D}} \xrightarrow{\text{onto}} \overline{\mathbb{S}}$ satisfy (3.3). Thus f maps the closed arc \mathcal{C} monotonically onto itself and \mathbf{T} homeomorphically onto the base $\mathbf{I} \subset \partial\mathbb{S}$. Suppose, in addition, that $f : \mathcal{C} \xrightarrow{\text{onto}} \mathcal{C}$ is injective on a compact subset $\mathbf{K} \subset \mathcal{C}$. Then there exist homeomorphisms $f_m : \overline{\mathbb{D}} \xrightarrow{\text{onto}} \overline{\mathbb{D}}$ converging to $f : \overline{\mathbb{D}} \xrightarrow{\text{onto}} \overline{\mathbb{S}}$ uniformly and in the Sobolev space $\mathcal{W}^{1,2}(\mathbb{D}, \mathbb{D})$. Moreover $f_m \equiv f$ on \mathbf{T} and \mathbf{K} , for $m = 4, 5, \dots$*

Thus, in steps IV and V, we shall concern ourselves only with the proof of this proposition.




FIGURE 3. Harmonic map $f : \mathbb{D} \xrightarrow{\text{onto}} \mathbb{S}$.

Step IV. *Good approximation of the boundary map $f : \partial\mathbb{D} \xrightarrow{\text{onto}} \partial\mathbb{S}$.* The aim is to properly approximate the continuous monotone boundary map $f : \partial\mathbb{D} \xrightarrow{\text{onto}} \partial\mathbb{S}$ by homeomorphisms $f_m : \partial\mathbb{D} \xrightarrow{\text{onto}} \partial\mathbb{S}$ which agree with f on both sets $\mathbf{T} = \overline{\mathbb{T}} \setminus \overline{\mathcal{C}}$ and $\mathbf{K} \subset \mathcal{C}$. Once such an approximation is in hand, we shall extend each f_m harmonically inside the disk. Certainly, uniform convergence $f_m \rightrightarrows f$ on \mathbf{T} would suffice to deduce uniform convergence in the entire disk, by the maximum principle. However, it is not obvious at all how to make the approximation of the boundary map in order to control the energy of the extended mappings. The Douglas criterion will come into play.

We need only work to construct homeomorphisms $f_m : \mathcal{C} \xrightarrow{\text{onto}} \mathcal{C}$, as their values on $\mathbf{T} = \overline{\mathbb{T}} \setminus \overline{\mathcal{C}}$ are already known; $f_m \equiv f : \mathbf{T} \xrightarrow{\text{onto}} \mathbf{T}$. Let us write $f : \mathcal{C} \xrightarrow{\text{onto}} \mathcal{C}$ as $f(e^{i\theta}) = e^{i\phi(\theta)}$, where $\phi : [-\omega, \omega] \xrightarrow{\text{onto}} [-\omega, \omega]$ is a nondecreasing continuous function such that $\phi(-\omega) = -\omega$ and $\phi(\omega) = \omega$. We also require that $f_m \equiv f$ on $\mathbf{K} \subset \mathcal{C}$. The complement $\mathcal{C} \setminus \mathbf{K}$ consists of a countable number of disjoint open circular arcs (components) whose endpoints belong to \mathbf{K} . Let $\mathcal{C}_\alpha^\beta \stackrel{\text{def}}{=} \{e^{i\theta} ; \alpha \leq$

$\theta \leq \beta$ be one of such arcs together with its endpoints, where $-\omega \leq \alpha < \beta \leq \omega$, and similarly for the image $-\omega \leq \phi(\alpha) < \phi(\beta) \leq \omega$. The latter strict inequality is justified by the fact that $e^{i\alpha}$ and $e^{i\beta}$ are points in \mathbf{K} and the map f , being injective on \mathbf{K} , assumes distinct values $e^{i\phi(\alpha)}$ and $e^{i\phi(\beta)}$ at these points.

We now define homeomorphisms between closed arcs $f_m : \mathcal{C}_\alpha^\beta \xrightarrow{\text{onto}} \mathcal{C}_{\phi(\alpha)}^{\phi(\beta)}$ ($m \geq 4$) by the rule; $f_m(e^{i\theta}) = e^{i\phi_m(\theta)}$, where

$$(3.4) \quad \phi_m(\theta) = \left[1 - \frac{\beta - \alpha}{m}\right] \cdot [\phi(\theta) - \phi(\alpha)] + \frac{\phi(\beta) - \phi(\alpha)}{m} \cdot (\theta - \alpha) + \phi(\alpha)$$

for $\alpha \leq \theta \leq \beta$. We have $\phi_m(\alpha) = \phi(\alpha)$ and $\phi_m(\beta) = \phi(\beta)$. The first term defining ϕ_m is nondecreasing in θ while the second term is strictly increasing. Therefore, $f_m : \mathcal{C}_\alpha^\beta \xrightarrow{\text{onto}} \mathcal{C}_{\phi(\alpha)}^{\phi(\beta)}$ is a homeomorphism. Formula (3.4), applied to every arc component of $\mathcal{C} \setminus \mathbf{K}$, gives homeomorphisms which agree with f at the endpoints of the arcs. We glue them together with f at the endpoints to obtain a homeomorphism $f_m : \mathbb{T} \xrightarrow{\text{onto}} \mathbb{T}$ which coincides with $f : \mathbb{T} \xrightarrow{\text{onto}} \mathbb{T}$ on $\mathbf{T} \cup \mathbf{K}$. Further analysis of f_m is necessary to deduce proper convergence as $m \rightarrow \infty$. First note that on each arc component of $\mathcal{C} \setminus \mathbf{K}$ we have

$$|f_m(e^{i\theta}) - f(e^{i\theta})| \leq |\phi_m(\theta) - \phi(\theta)| \leq \frac{8\omega^2}{m} < \frac{21}{m}$$

Hence

$$(3.5) \quad |f_m(\xi) - f(\xi)| \leq \frac{21}{m}, \quad \text{for every } \xi \in \mathbb{T} \text{ and } m = 4, 5, \dots$$

In particular, $f_m \rightrightarrows f$ uniformly on \mathbb{T} .

Lemma 3.2. *For all $\xi_1, \xi_2 \in \mathbb{T}$ and $m = 4, 5, \dots$, we have*

$$(3.6) \quad |f_m(\xi_1) - f_m(\xi_2)| \leq \frac{5}{\sin(\omega/4)} |f(\xi_1) - f(\xi_2)| + 4|\xi_1 - \xi_2|$$

Proof. There are three cases to consider:

Case 1. We first do the case when both ξ_1 and ξ_2 belong to the closure of the same component of $\mathcal{C} \setminus \mathbf{K}$; say, $\xi_1 = e^{i\theta_1} \in \mathcal{C}_\alpha^\beta$ and $\xi_2 = e^{i\theta_2} \in \mathcal{C}_\alpha^\beta$, where $\alpha \leq \theta_1, \theta_2 \leq \beta$. It follows from formula (3.4) that

$$\begin{aligned} |f_m(e^{i\theta_1}) - f_m(e^{i\theta_2})| &\leq |\phi_m(\theta_1) - \phi_m(\theta_2)| \leq |\phi(\theta_1) - \phi(\theta_2)| + |\theta_1 - \theta_2| \\ &\leq 2|f(e^{i\theta_1}) - f(e^{i\theta_2})| + 2|e^{i\theta_1} - e^{i\theta_2}| \end{aligned}$$

Case 2. Both $\xi_1 = e^{i\theta_1}$ and $\xi_2 = e^{i\theta_2}$ belong to \mathcal{C} . We assume that the closed set $\{\tau : \theta_1 \leq \tau \leq \theta_2, e^{i\tau} \in \mathbf{K}\}$ is not empty. Otherwise, ξ_1 and ξ_2 would belong to the same arc component of $\mathcal{C} \setminus \mathbf{K}$. Let us set the notation,

$$\tau_1 \stackrel{\text{def}}{=} \min \{ \tau : \theta_1 \leq \tau \leq \theta_2, e^{i\tau} \in \mathbf{K} \}, \quad \widehat{\xi}_1 \stackrel{\text{def}}{=} e^{i\tau_1} \in \mathbf{K}$$

$$\tau_2 \stackrel{\text{def}}{=} \max \{ \tau : \theta_1 \leq \tau \leq \theta_2, e^{i\tau} \in \mathbf{K} \}, \quad \widehat{\xi}_2 \stackrel{\text{def}}{=} e^{i\tau_2} \in \mathbf{K}$$

and note that the points $\xi_1, \widehat{\xi}_1$ belong to the closure of one arc component in $\mathcal{C} \setminus \mathbf{K}$. The same applies to the pair $\xi_2, \widehat{\xi}_2$. Therefore, using *Case 1*, one can write the following chain of inequalities

$$\begin{aligned}
& |f_m(\xi_1) - f_m(\xi_2)| \\
& \leq |f_m(\xi_1) - f_m(\widehat{\xi}_1)| + |f_m(\widehat{\xi}_1) - f_m(\widehat{\xi}_2)| + |f_m(\widehat{\xi}_2) - f_m(\xi_2)| \\
& \leq 2|f(\xi_1) - f(\widehat{\xi}_1)| + 2|\xi_1 - \widehat{\xi}_1| + |f(\widehat{\xi}_1) - f(\widehat{\xi}_2)| \\
& \quad + 2|f(\widehat{\xi}_2) - f(\xi_2)| + 2|\widehat{\xi}_2 - \xi_2|
\end{aligned}$$

Since $\widehat{\xi}_1$ and $\widehat{\xi}_2$ lay in the shorter circular arc between ξ_1 and ξ_2 , it follows that

$$|\xi_1 - \widehat{\xi}_1| \leq |\xi_1 - \xi_2| \quad \text{and} \quad |\widehat{\xi}_2 - \xi_2| \leq |\xi_1 - \xi_2|$$

The same argument applies to the images of these point under the monotone map $f : \mathcal{C} \xrightarrow{\text{onto}} \mathcal{C}$. Thus,

$$|f(\xi_1) - f(\widehat{\xi}_1)| \leq |f(\xi_1) - f(\xi_2)| \quad \text{and} \quad |f(\widehat{\xi}_2) - f(\xi_2)| \leq |f(\xi_1) - f(\xi_2)|$$

and also $|f(\widehat{\xi}_1) - f(\widehat{\xi}_2)| \leq |f(\xi_1) - f(\xi_2)|$. Substitute these inequalities into the chain above to conclude with the desired inequality

$$(3.7) \quad |f_m(\xi_1) - f_m(\xi_2)| \leq 5|f(\xi_1) - f(\xi_2)| + 4|\xi_1 - \xi_2|$$

The case $\xi_1, \xi_2 \in \mathbf{T}$ is trivial, because $f_m(\xi_1) - f_m(\xi_2) = f(\xi_1) - f(\xi_2)$. Thus, all that remains is to consider

Case 3. Let $\xi_1 \in \mathcal{C}$ and $\xi_2 \in \mathbf{T}$. By symmetry we may take $\xi_1 = e^{i\theta_1}$, where $0 \leq \theta_1 < \omega$. We may also assume that $|\xi_1 - \xi_2| \leq \sin \omega$. Otherwise, inequality (3.6) holds; namely, $|f_m(\xi_1) - f_m(\xi_2)| \leq \text{diam } \mathbb{S} = 2 \sin \omega < 2|\xi_1 - \xi_2|$. Geometrically, the assumption $|\xi_1 - \xi_2| \leq \sin \omega$ tells us that ξ_2 cannot lay in the lower half of the arc \mathbf{T} . Thus $\xi_2 = e^{i\theta_2}$, where $\omega < \theta_2 \leq \pi$. Let the upper corner of the segment \mathbb{S} be denoted by $\xi^+ = e^{i\omega}$. The location of $f(\xi_2)$ is restricted to the upper half of the base of the segment \mathbb{S} , because $f : \mathbb{T} \xrightarrow{\text{onto}} \mathbb{T}$ is monotone and $f(e^{i\pi}) = \cos \omega$ (the midpoint of the base) due to normalization at (3.3). Regarding the position of $f(\xi_1) \in \mathcal{C}$, we may assume that this value also lies in the upper half of the arc \mathcal{C} . Otherwise, we would have $|f(\xi_1) - f(\xi_2)| \geq 1 - \cos \omega = 2 \sin^2 \frac{\omega}{2}$ while, on the other hand,

$$\begin{aligned}
|f_m(\xi_1) - f_m(\xi_2)| & \leq \text{diam } \mathbb{S} = 2 \sin \omega < \frac{5}{\sin(\omega/4)} \cdot 2 \sin^2 \frac{\omega}{2} \\
& \leq \frac{5}{\sin(\omega/4)} |f(\xi_1) - f(\xi_2)|
\end{aligned}$$

which implies (3.6).

We are ready to complete Case 3. First we use the triangle inequality and (3.7),

$$\begin{aligned}
|f_m(\xi_1) - f_m(\xi_2)| & \leq |f_m(\xi_1) - f_m(\xi^+)| + |f_m(\xi^+) - f_m(\xi_2)| \\
& \leq 5|f(\xi_1) - f(\xi^+)| + 4|\xi_1 - \xi^+| + |f(\xi^+) - f(\xi_2)| \\
& \leq 5\{|f(\xi_1) - f(\xi^+)| + |f(\xi^+) - f(\xi_2)|\} + 4|\xi_1 - \xi_2|
\end{aligned}$$

Then comes a geometric fact about the term within the curled braces. Certainly, we have $f(\xi^+) \neq f(\xi_2)$, because f is injective on \mathbf{T} . If, incidentally, $f(\xi^+) = f(\xi_1)$ then the latter estimate yields (3.6). Thus we may assume that three points $A \stackrel{\text{def}}{=} f(\xi_1)$, $B \stackrel{\text{def}}{=} f(\xi_2)$ and $C \stackrel{\text{def}}{=} f(\xi^+)$ are vertices of a triangle. Let $a = |B - C|$, $b = |A - C|$ and $c = |A - B|$. Since A lies in the arc of \mathbb{S} , B lies in the base of \mathbb{S} and C is the corner of \mathbb{S} , all of them in the upper half of \mathbb{S} , it follows (from

geometry of the segment \mathbb{S}) that the angle opposite to the side \overline{AB} , denoted by γ , satisfies: $\frac{\omega}{2} \leq \gamma < \omega$. The law of cosines tells us that

$$c^2 = a^2 + b^2 - 2ab \cos \gamma \geq a^2 + b^2 - 2ab \cos(\omega/2) \geq (a + b)^2 \sin^2 \frac{\omega}{4}.$$

Hence $a + b \leq \frac{c}{\sin(\omega/4)}$; that is,

$$|f(\xi_1) - f(\xi^+)| + |f(\xi^+) - f(\xi_2)| \leq \frac{1}{\sin(\omega/4)} |f(\xi_1) - f(\xi_2)|,$$

completing the proof of Lemma 3.2. \square

Step V. *Harmonic extension and strong convergence in $\mathcal{W}^{1,2}(\mathbb{D})$.* The boundary homeomorphisms $f_m : \mathbb{T} \xrightarrow{\text{onto}} \partial\mathbb{S}$ will now be extended harmonically inside the unit disk. We use the same label for the extensions, $f_m : \overline{\mathbb{D}} \xrightarrow{\text{onto}} \overline{\mathbb{S}}$. These mappings are homeomorphisms, due to Theorem 2.7. Since both f and f_m are harmonic in \mathbb{D} , the sequence f_m converges to f uniformly in $\overline{\mathbb{D}}$, by the maximum principle. The key point here is that they also belong to the Sobolev space $\mathcal{W}^{1,2}(\mathbb{D})$, and converge in the Sobolev norm as well. To see this we recall the Douglas criterion [9] which asserts that any function g that is continuous on $\overline{\mathbb{D}}$ and harmonic in \mathbb{D} satisfies

$$(3.8) \quad \mathcal{E}_{\mathbb{D}}[g] \stackrel{\text{def}}{=} \iint_{\mathbb{D}} |Dg|^2 = \frac{1}{2\pi} \int_{\mathbb{T}} \int_{\mathbb{T}} \left| \frac{g(\xi) - g(\zeta)}{\xi - \zeta} \right|^2 |d\xi| \cdot |d\zeta|.$$

Recall that by (3.6) the mappings $f_m : \mathbb{T} \xrightarrow{\text{onto}} \partial\mathbb{S}$ satisfy

$$\left| \frac{f_m(\xi) - f_m(\zeta)}{\xi - \zeta} \right|^2 \preceq \left| \frac{f(\xi) - f(\zeta)}{\xi - \zeta} \right|^2 + 1$$

where the implied constant does not depend on m . By virtue of (3.8) this implies

$$\mathcal{E}_{\mathbb{D}}[f_m] \preceq \mathcal{E}_{\mathbb{D}}[f] + 1 < \infty.$$

Therefore f_m have uniformly bounded energy. It follows that f_m converge to f not only uniformly but also weakly in $\mathcal{W}^{1,2}(\mathbb{D})$. In particular, $\mathcal{E}[f] \leq \liminf \mathcal{E}[f_m]$. It is crucial to notice, using Dominated Convergence Theorem, that in fact we have equality

$$(3.9) \quad \begin{aligned} \mathcal{E}_{\mathbb{D}}[f] &= \frac{1}{2\pi} \int_{\mathbb{T}} \int_{\mathbb{T}} \left| \frac{f(\xi) - f(\zeta)}{\xi - \zeta} \right|^2 |d\xi| \cdot |d\zeta| \\ &= \lim \frac{1}{2\pi} \int_{\mathbb{T}} \int_{\mathbb{T}} \left| \frac{f_m(\xi) - f_m(\zeta)}{\xi - \zeta} \right|^2 |d\xi| \cdot |d\zeta| = \lim \mathcal{E}_{\mathbb{D}}[f_m]. \end{aligned}$$

This shows that f_m converge to f strongly in $\mathcal{W}^{1,2}(\mathbb{D})$, completing the proof of Proposition 3.1 and thus of Theorem 1.1. \square

REFERENCES

1. J. M. Ball, *Singularities and computation of minimizers for variational problems*, Foundations of computational mathematics (Oxford, 1999), 1–20, London Math. Soc. Lecture Note Ser., 284, Cambridge Univ. Press, Cambridge, 2001.
2. J. M. Ball, *Progress and puzzles in nonlinear elasticity*, in “Poly-, quasi- and rank-one convexity in applied mechanics” (eds. J. Schröder and P. Neff), Proceedings of CISM International Centre for Mechanical Sciences, vol. 516 (2010), 1–15.

3. P. Bauman, D. Phillips, and N. Owen, *Maximum principles and a priori estimates for an incompressible material in nonlinear elasticity*, Comm. Partial Differential Equations **17** (1992), no. 7-8, 1185–1212.
4. J. C. Bellido and C. Mora-Corral, *Approximation of Hölder continuous homeomorphisms by piecewise affine homeomorphisms*, Houston J. Math., **37**, no. 2 (2011) 449–500.
5. M. Brown, *A proof of the generalized Schoenflies theorem*, Bull. Amer. Math. Soc. **66** (1960) 74–76.
6. S. Conti and C. De Lellis, *Some remarks on the theory of elasticity for compressible Neo-hookean materials*, Ann. Sc. Norm. Super. Pisa Cl. Sci. (5) **2** (2003) 521–549.
7. J. Cristina, T. Iwaniec, L. V. Kovalev and J. Onninen, *Lipschitz regularity for the Hopf-Laplace equation*, arXiv:1011.5934.
8. S. Daneri and A. Pratelli, *Smooth approximation of bi-Lipschitz orientation-preserving homeomorphisms*, arXiv:1106.1192.
9. J. Douglas, *Solution of the problem of Plateau*, Trans. Amer. Math. Soc., **33**, (1931), 231–321.
10. P. Duren, *Harmonic mappings in the plane*, Cambridge Tracts in Mathematics, 156. Cambridge University Press, Cambridge, 2004.
11. D. M. Freeman and D. A. Herron, *Bilipschitz homogeneity and inner diameter distance*, J. Anal. Math. **111** (2010), no. 1, 1–46.
12. D. A. Herron and T. S. Sullivan, *Fractal inner chordarc disks*, J. Anal. Math. **84** (2001), no. 1, 173–205.
13. T. Iwaniec, L. V. Kovalev and J. Onninen, *Hopf differentials and smoothing Sobolev homeomorphisms*, Int. Math. Res. Not. IMRN, to appear.
14. T. Iwaniec, L. V. Kovalev and J. Onninen, *Diffeomorphic approximation of Sobolev homeomorphisms* Arch. Rat. Mech. Anal., to appear.
15. T. Iwaniec and J. Onninen, *Deformations of finite conformal energy: Boundary behavior and limit theorems*, Trans. Amer. Math. Soc. **363** (2011), no. 11, 5605–5648.
16. T. G. Latfullin, *Geometric characterization of the quasi-isometric image of a half plane*, in “Theory of mappings, its generalizations and applications”, 116–126, Naukova Dumka, Kiev, 1982.
17. L. F. McAuley, *Some fundamental theorems and problems related to monotone mappings* 1971 Proc. First Conf. on Monotone Mappings and Open Mappings (SUNY at Binghamton, Binghamton, N.Y., 1970). 1–36. State Univ. of New York at Binghamton, N.Y.
18. C. Mora-Corral, *Approximation by piecewise affine homeomorphisms of Sobolev homeomorphisms that are smooth outside a point*, Houston J. Math. **35** (2009), no. 2, 515–539.
19. C. B. Morrey, *The Topology of (Path) Surfaces*, Amer. J. Math. **57** (1935), no. 1, 17–50.
20. S. Müller, S. J. Spector, and Q. Tang, *Invertibility and a topological property of Sobolev maps*, SIAM J. Math. Anal. **27** (1996), no. 4, 959–976.
21. Ch. Pommerenke, *One-sided smoothness conditions and conformal mapping*, J. London Math. Soc. (2) **26** (1982), no. 1, 77–88.
22. S. W. Semmes, *Quasiconformal mappings and chord-arc curves*, Trans. Amer. Math. Soc. **306** (1988), no. 1, 233–263.
23. T. Radó, *Length and Area*, American Mathematical Society, New York, 1948.
24. P. Tukia, *The planar Schönflies theorem for Lipschitz maps*, Ann. Acad. Sci. Fenn. Ser. A I Math. **5** (1980), no. 1, 49–72.
25. J. Väisälä, *Homeomorphisms of bounded length distortion*, Ann. Acad. Sci. Fenn. Ser. A I Math. **12** (1987), no. 2, 303–312.
26. J. Väisälä, *Quasiconformal maps of cylindrical domains*, Acta Math. **162** (1989), no. 3-4, 201–225.
27. J. W. T. Youngs, *The topological theory of Fréchet surfaces*, Ann. of Math. (2) **45** (1944), 753–785.

DEPARTMENT OF MATHEMATICS, SYRACUSE UNIVERSITY, SYRACUSE, NY 13244, USA AND DE-
PARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF HELSINKI, FINLAND
E-mail address: `tiwaniec@syr.edu`

DEPARTMENT OF MATHEMATICS, SYRACUSE UNIVERSITY, SYRACUSE, NY 13244, USA
E-mail address: `lvkovaie@syr.edu`

DEPARTMENT OF MATHEMATICS, SYRACUSE UNIVERSITY, SYRACUSE, NY 13244, USA
E-mail address: `jkonnine@syr.edu`