

Syracuse University

SURFACE

Mathematics - Faculty Scholarship

Mathematics

4-14-2011

A Direct Limit for Limit Hilbert-Kunz Multiplicity for Smooth Projective Curves

Holger Brenner
Universitat Osnabruck,

Jinjia Li
Syracuse University and University of Louisville

Claudia Miller
Syracuse University

Follow this and additional works at: <https://surface.syr.edu/mat>



Part of the [Mathematics Commons](#)

Recommended Citation

Brenner, Holger; Li, Jinjia; and Miller, Claudia, "A Direct Limit for Limit Hilbert-Kunz Multiplicity for Smooth Projective Curves" (2011). *Mathematics - Faculty Scholarship*. 12.
<https://surface.syr.edu/mat/12>

This Article is brought to you for free and open access by the Mathematics at SURFACE. It has been accepted for inclusion in Mathematics - Faculty Scholarship by an authorized administrator of SURFACE. For more information, please contact surface@syr.edu.

A DIRECT LIMIT FOR LIMIT HILBERT-KUNZ MULTIPLICITY FOR SMOOTH PROJECTIVE CURVES

HOLGER BRENNER, JINJIA LI, AND CLAUDIA MILLER

ABSTRACT. This paper concerns the question of whether a more direct limit can be used to obtain the limit Hilbert Kunz multiplicity, a possible candidate for a characteristic zero Hilbert-Kunz multiplicity. The main goal is to establish an affirmative answer for one of the main cases for which the limit Hilbert Kunz multiplicity is even known to exist, namely that of graded ideals in the homogeneous coordinate ring of smooth projective curves. The proof involves more careful estimates of bounds found independently by Brenner and Trivedi on the dimensions of the cohomologies of twists of the syzygy bundle as the characteristic p goes to infinity and uses asymptotic results of Trivedi on the slopes of Harder-Narasimham filtrations of Frobenius pullbacks of bundles. In view of unpublished results of Gessel and Monsky, the case of maximal ideals in diagonal hypersurfaces is also discussed in depth.

INTRODUCTION

In 1983, following Kunz's lead in [9], Monsky defined a new multiplicity in positive characteristic – the Hilbert-Kunz (HK) multiplicity as follows: Let R be a ring of characteristic $p > 0$ and $I = (f_1, \dots, f_s)$ an ideal with the length $\ell(R/I)$ finite. Consider the Frobenius powers $I^{[p^n]} = (f_1^{p^n}, \dots, f_s^{p^n})$ of I and define

$$e_{\text{HK}}(I, R) = \lim_{n \rightarrow \infty} \frac{\ell(R/I^{[p^n]})}{(p^n)^{\dim(R)}}$$

Just like the usual Hilbert-Samuel multiplicity, this new multiplicity seems to measure the degree of singularity at a point on a variety. Furthermore, it plays the role for tight closure that ordinary Hilbert-Samuel multiplicity plays for integral closure. But the numbers seem much more complex (they are usually not integers and possibly not

Date: April 15, 2011.

1991 Mathematics Subject Classification. primary 13D40, 14H60.

Key words and phrases. Hilbert-Kunz multiplicity, Harder-Narasimhan filtrations.

The third author was supported in part by NSA Grant #H98230-08-1-0025.

always rational or even algebraic) than usual multiplicities (which are integers) and, despite intense study in recent years, are still not well understood or even computable except in a few cases.

However what little is known seems to indicate that the numbers may get simpler in the limit as the characteristic p goes to infinity, leading to the question of whether a characteristic zero HK multiplicity defined in such a way could have a more transparent meaning or behavior than the one in characteristic p does. More precisely, if R is a \mathbb{Z} -algebra and I an ideal, let R_p be the reduction of R mod p and I_p the extended ideal. If $\ell(R_p/I_p)$ is finite for almost all p , define

$$e_{\text{HK}}^{\infty}(I, R) \stackrel{\text{def}}{=} \lim_{p \rightarrow \infty} e_{\text{HK}}(I_p, R_p)$$

whenever this limit exists, and call it the *limit Hilbert-Kunz multiplicity* of I .

Although experimental results indicate this limit might always exist, very few cases have been established. It is, of course, clear when $e_{\text{HK}}(I_p, R_p)$ is constant for almost all p , such as for the homogeneous maximal ideal in the coordinate rings of plane cubics [3], [12], [15], in certain monomial ideals [2], [5], [6], [20], and in two-dimensional invariant rings under finite group actions [21]. That this is also the case for ideals of finite projective dimension can be seen via local Riemann-Roch theory (private communication with Kurano); it is interesting that in this last case the limit has an intrinsic geometric interpretation in characteristic zero. A few nonconstant cases are known as well: The limit was shown to exist for the homogeneous maximal ideal of diagonal hypersurfaces, in unpublished work of Gessel and Monsky [14] building on [7]. It was also shown to exist for any homogeneous ideal primary to the homogeneous maximal ideal in homogeneous coordinate rings of smooth projective curves by Trivedi in [18] by delicate study of the variation of Harder-Narasimhan filtrations of Frobenius pullbacks of the syzygy bundle relative to the characteristic p . The limit in this case turns out again to have an intrinsic geometric description in characteristic zero.

In this paper, we are interested in the question of whether a simpler limit gives the same result. In particular, is it necessary to use the full HK multiplicity $e_{\text{HK}}(I_p, R_p)$ in each characteristic p ? This value is itself the usually uncomputable limit $\lim_{n \rightarrow \infty} \frac{\ell(R_p/I_p^{[p^n]})}{(p^n)^d}$ where $d = \dim R$. We propose to replace this complex limit with its first term $\frac{\ell(R_p/I_p^{[p]})}{p^d}$ or more generally any fixed degree term as follows:

Question. Assuming $e_{\text{HK}}^\infty(I, R)$ exists, is it true that for any fixed $n \geq 1$

$$e_{\text{HK}}^\infty(I, R) = \lim_{p \rightarrow \infty} \frac{\ell(R_p/I_p^{[p^n]})}{(p^n)^d} \quad ?$$

Informally, in measuring colengths of p^n th twists of the ideal, if p goes to infinity, is it really necessary to first let n go to infinity?

The motivation behind such a modification is that a simpler limit may make it easier to find a geometric interpretation of the limit HK multiplicity in characteristic zero. It would be encouraging to see a simpler limit giving the possible characteristic zero concept. A drawback is that it still does not yield an intrinsic definition of $e_{\text{HK}}^\infty(I, R)$ in a characteristic zero setting.

The main goal of this paper is to establish an affirmative answer to the question for the case of the homogeneous coordinate rings of smooth projective curves. Our proof is based on the proofs in this setting of Brenner [1] and Trivedi [17] of a formula for the HK multiplicity and of Trivedi [18] regarding the existence of $e_{\text{HK}}^\infty(I, R)$, but requires some additional work as we may not assume that the fixed value n is large enough to give strong Harder-Narasimham filtrations of the syzygy bundle (the case $n = 1$ is the most important). Fortunately, the gap can be filled using Trivedi's results mentioned above to yield:

Corollary 3.3 *Let R be a standard-graded flat domain over \mathbb{Z} such that almost all fiber rings $R_p = R \otimes_{\mathbb{Z}} \mathbb{Z}/p\mathbb{Z}$ are geometrically normal 2-dimensional domains and let $I = (f_1, \dots, f_s)$ be a homogeneous R_+ -primary ideal. With the notation as above, for any fixed $n \geq 1$ one has*

$$\frac{\ell(R_p/I_p^{[p^n]})}{(p^n)^2} = e_{\text{HK}}^\infty(I, R) + O\left(\frac{1}{p}\right)$$

We remark that, with this result, the answer to the question above is known to be yes in all the main cases in which $e_{\text{HK}}^\infty(I, R)$ is known to exist so far.

Section 1 contains a review of the background. The groundwork for our main result is done in Section 2 via some lemmas on the asymptotic growth of cohomologies of bundles as the characteristic p goes to infinity. In Section 3 these lemmas are applied to the syzygy bundle, defined in (1.1), to obtain the corollary above.

The remaining part, Section 4, is devoted to a discussion of consequences of Gessel and Monsky's unpublished work [14]. We see that a side-product of their proof is an affirmative answer to the question above for the case of diagonal hypersurfaces. Furthermore, his work shows that the most tempting naive limit in characteristic zero does not give $e_{\text{HK}}^\infty(I, R)$.

Finally, we mention our convention regarding asymptotics throughout the paper: Let $q = p^n$. We emphasize that for the asymptotic notation $O(-)$ used throughout the paper, such as in $O(\frac{q^2}{p})$, $O(q)$, or even $O(1)$, we have fixed $n > 0$ and let $p \rightarrow \infty$ (unlike in [1] and [17], where p is fixed and n is allowed to go to infinity).

1. PRELIMINARIES AND BACKGROUND

In this section, we present the basic set-up and notations and review relevant results on vector bundles.

Basic set-up

Let R be a standard-graded flat domain over \mathbb{Z} such that almost all fiber rings $R_p = R \otimes_{\mathbb{Z}} \mathbb{Z}/p\mathbb{Z}$ are geometrically normal 2-dimensional domains. Let $I = (f_1, \dots, f_s)$ be a homogeneous R_+ -primary ideal with $\deg f_i = d_i$. Let $Y = \text{Proj } R_{\mathbb{Q}}$ where $R_{\mathbb{Q}} = R \otimes_{\mathbb{Z}} \mathbb{Q}$. For each prime p , consider the reduction to characteristic p

$$R_p = R \otimes_{\mathbb{Z}} \mathbb{Z}/p\mathbb{Z}, \quad I_p = IR_p, \quad Y_p = \text{Proj } R_p$$

Due to our assumptions, Y and Y_p are smooth projective curves for almost all p . The corresponding Hilbert-Kunz multiplicity is

$$e_{\text{HK}}(I_p, R_p) \stackrel{\text{def}}{=} \lim_{n \rightarrow \infty} \frac{\ell(R_p/I_p^{[q]})}{q^2}$$

where $q = p^n$. The key idea in [1] and [17] for determining the Hilbert-Kunz multiplicity is to consider the syzygy bundle $\mathcal{S} = \text{Syz}(f_1, \dots, f_s)$ on Y_p (and on Y) given by

$$(1.1) \quad 0 \longrightarrow \mathcal{S} \longrightarrow \bigoplus_{i=1}^s \mathcal{O}(-d_i) \xrightarrow{f_1, \dots, f_s} \mathcal{O} \longrightarrow 0$$

and the pullback of this exact sequence n times along the absolute Frobenius morphism $F : Y_p \rightarrow Y_p$ (with a subsequent twist by $m \in \mathbb{Z}$)

$$(1.2) \quad 0 \rightarrow \mathcal{S}^q(m) \rightarrow \bigoplus_{i=1}^s \mathcal{O}(m - qd_i) \xrightarrow{f_1^q, \dots, f_s^q} \mathcal{O}(m) \rightarrow 0$$

where \mathcal{S}^q denotes the pullback $(F^*)^n(\mathcal{S}) = \text{Syz}(f_1^q, \dots, f_s^q)$.

Remark 1.1. Notice that for simplicity, we use the notation \mathcal{S} for the syzygy bundle over any Y_p , as the characteristic is usually obvious from the context (we study mostly \mathcal{S}^q , not \mathcal{S}). The first sequence is just a reduction mod p of the corresponding sequence in characteristic zero. In particular, \mathcal{S} is the reduction to Y_p of the syzygy bundle on Y .

As R_p is normal, the cokernel of the second map in the associated long exact sequence of cohomology

$$0 \rightarrow H^0(Y_p, \mathcal{S}^q(m)) \rightarrow \bigoplus_{i=1}^s H^0(Y_p, \mathcal{O}(m - qd_i)) \xrightarrow{f_1^q, \dots, f_s^q} H^0(Y_p, \mathcal{O}(m)) \rightarrow \dots$$

is the m -th graded piece of $R_p/I_p^{[q]}$. Brenner [1] and Trivedi [17] exploited this connection to $H^0(Y_p, \mathcal{S}^q(m))$ to determine the Hilbert-Kunz multiplicity of I_p in terms of intrinsic properties of the syzygy bundle, which we review next.

Harder-Narasimhan filtrations

Let X be a smooth projective curve over an algebraically closed field. For any vector bundle \mathcal{V} on X of rank r , the *degree* and *slope* are defined respectively as

$$\deg(\mathcal{V}) \stackrel{\text{def}}{=} \deg(\wedge^r \mathcal{V}) \qquad \mu(\mathcal{V}) \stackrel{\text{def}}{=} \frac{\deg(\mathcal{V})}{r}$$

Slope is additive on tensor products of bundles: $\mu(\mathcal{V} \otimes \mathcal{W}) = \mu(\mathcal{V}) + \mu(\mathcal{W})$. If $f : X' \rightarrow X$ is a finite map of degree q , then $\deg(f^*(\mathcal{V})) = q \deg(\mathcal{V})$ and so $\mu(f^*(\mathcal{V})) = q\mu(\mathcal{V})$.

A bundle \mathcal{V} is called *semistable* if for every subbundle $\mathcal{W} \subseteq \mathcal{V}$ one has $\mu(\mathcal{W}) \leq \mu(\mathcal{V})$. Clearly, bundles of rank 1 are always semistable, and duals and twists of semistable bundles are semistable.

Any bundle \mathcal{V} has a filtration by subbundles

$$0 = \mathcal{V}_0 \subset \mathcal{V}_1 \subset \dots \subset \mathcal{V}_t = \mathcal{V}$$

such that $\mathcal{V}_k/\mathcal{V}_{k-1}$ is semistable and $\mu(\mathcal{V}_k/\mathcal{V}_{k-1}) > \mu(\mathcal{V}_{k+1}/\mathcal{V}_k)$ for each k . This filtration is unique, and it is called the *Harder-Narasimhan (or HN) filtration* of \mathcal{V} .

The *maximal* and *minimal slopes* are defined as

$$\mu_{\max}(\mathcal{V}) \stackrel{\text{def}}{=} \mu(\mathcal{V}_1/\mathcal{V}_0) \qquad \mu_{\min}(\mathcal{V}) \stackrel{\text{def}}{=} \mu(\mathcal{V}_t/\mathcal{V}_{t-1})$$

Remark 1.2. *In positive characteristic, pulling back under the Frobenius morphism F does not necessarily preserve semistability. Therefore, the pullback under F^n of an HN filtration of \mathcal{V} does not always give an HN filtration of $(F^*)^n(\mathcal{V})$. Crucial to the work in [1] and [17] was the existence of a strong HN filtration from [10], i.e., for some n_0 , the HN filtration of $(F^*)^{n_0}(\mathcal{V})$ has the property that all its Frobenius pullbacks are the HN filtrations of $(F^*)^n(\mathcal{V})$, for all $n > n_0$.*

We do not need strong HN filtrations here since for us n is fixed at a given value and cannot be modified, but we do need some relation between the HN filtrations of \mathcal{S} and \mathcal{S}^q . Fortunately, for $p \gg 0$, the following refinement result by Trivedi [18, Lemmas 1.8 and 1.14] applies:

Proposition 1.3 (Trivedi). *Let \mathcal{V} be a bundle of rank r on a smooth projective curve X of genus g over an algebraically closed field of characteristic p with $p > 4(g-1)r^3$. Let $n \geq 1$ and $q = p^n$. If*

$$0 = \mathcal{V}_0 \subset \mathcal{V}_1 \subset \cdots \subset \mathcal{V}_t = \mathcal{V}$$

is the HN filtration of \mathcal{V} , then its pullback

$$0 = (F^*)^n(\mathcal{V}_0) \subset (F^*)^n(\mathcal{V}_1) \subset \cdots \subset (F^*)^n(\mathcal{V}_t) = (F^*)^n(\mathcal{V})$$

can be refined to the HN filtration of $(F^)^n(\mathcal{V})$.*

Furthermore, denoting the k th portion of the refined filtration as follows

$$(F^*)^n(\mathcal{V}_{k-1}) = \mathcal{V}_{k,0} \subset \mathcal{V}_{k,1} \subset \cdots \subset \mathcal{V}_{k,t_k} = (F^*)^n(\mathcal{V}_k)$$

one has that for any i

$$\left| \frac{\mu(\mathcal{V}_{k,i}/\mathcal{V}_{k,i-1})}{q} - \mu(\mathcal{V}_k/\mathcal{V}_{k-1}) \right| \leq \frac{C}{p}$$

where C is a constant depending only on g and r .

In our situation the curves Y and Y_p are not defined over an algebraically closed field, but due to our assumptions the curves $Y_{\overline{\mathbb{Q}}} = Y \times_{\mathbb{Q}} \overline{\mathbb{Q}}$ and $\overline{Y}_p = Y_p \times_{\mathbb{Z}/p\mathbb{Z}} \overline{\mathbb{Z}/p\mathbb{Z}}$

are smooth projective curves over the algebraic closures. In our setting the definition of degree, semistability and the Harder-Narasimhan filtration descends to the original curves. Hence we will move to the algebraic closure and back whenever this is convenient. Moreover, because of the openness of semistability in a family, the Harder-Narasimhan filtration of \mathcal{S} on Y extends to the Harder-Narasimhan filtration almost everywhere, so that the slopes of the quotients in the Harder-Narasimhan filtration of \mathcal{S} on Y_p are constant for almost all p .

2. ASYMPTOTIC LEMMAS FOR BUNDLES

In this section we prove various asymptotic results on the cohomologies of bundles that will be used in the next section for the proof of the main result. Let \mathcal{S} be any bundle on the relative curve $\text{Proj } R \rightarrow \text{Spec } \mathbb{Z}$. Fix $n \geq 0$ and set $q = p^n$ for varying p . We denote the restriction of \mathcal{S} to Y_p again by the symbol \mathcal{S} , as this should cause no confusion in context. We first review the notation that we use to describe concisely the data from the various HN filtrations.

Notation

We continue this practice of introducing notation unadorned by the characteristic p as it will always be obvious from the context.

First, for each p , write the HN filtration of \mathcal{S} as

$$0 = \mathcal{S}_0 \subset \mathcal{S}_1 \subset \cdots \subset \mathcal{S}_t = \mathcal{S}$$

with slopes, normalized slopes, and ranks (for $k = 1, \dots, t$) defined as follows:

$$\mu_k \stackrel{\text{def}}{=} \mu(\mathcal{S}_k/\mathcal{S}_{k-1}) \quad \nu_k \stackrel{\text{def}}{=} \frac{-\mu_k}{\deg Y_p} \quad r_k \stackrel{\text{def}}{=} \text{rank}(\mathcal{S}_k/\mathcal{S}_{k-1})$$

Throughout we will assume that p has been taken to be large enough so that the notations μ_k , ν_k and r_k refer to constants.

Taking pullbacks under the n th Frobenius morphism and setting

$$\mathcal{S}_k^q \stackrel{\text{def}}{=} (F^*)^n(\mathcal{S}_k)$$

gives

$$0 = \mathcal{S}_0^q \subset \mathcal{S}_1^q \subset \cdots \subset \mathcal{S}_t^q = \mathcal{S}^q$$

By Proposition 1.3, for $p \gg 0$, the HN filtration of \mathcal{S}^q can be obtained by refining each containment above, say as

$$\mathcal{S}_{k-1}^q = \mathcal{S}_{k,0} \subset \mathcal{S}_{k,1} \subset \cdots \subset \mathcal{S}_{k,t_k} = \mathcal{S}_k^q$$

We denote the maximal and minimal slopes in this portion as

$$\mu_k^{max} \stackrel{\text{def}}{=} \mu(\mathcal{S}_{k,1}/\mathcal{S}_{k,0}) \quad \text{and} \quad \mu_k^{min} \stackrel{\text{def}}{=} \mu(\mathcal{S}_{k,t_k}/\mathcal{S}_{k,t_k-1})$$

(we will not need the intermediate slopes). Further, we define normalized versions of these slopes as

$$\nu_k^{max} \stackrel{\text{def}}{=} \frac{-\mu_k^{max}}{q \deg Y_p} \quad \text{and} \quad \nu_k^{min} \stackrel{\text{def}}{=} \frac{-\mu_k^{min}}{q \deg Y_p}$$

Note that

$$\mu_1^{max} > \mu_1^{min} > \mu_2^{max} > \mu_2^{min} > \cdots > \mu_k^{max} > \mu_k^{min} > \cdots > \mu_t^{max} > \mu_t^{min}$$

and therefore

$$\nu_1^{max} < \nu_1^{min} < \nu_2^{max} < \nu_2^{min} < \cdots < \nu_k^{max} < \nu_k^{min} < \cdots < \nu_t^{max} < \nu_t^{min}$$

In this situation, Trivedi's result, Proposition 1.3, becomes:

Corollary 2.1 (Trivedi). *With the notations as above, for any k , as $p \rightarrow \infty$*

$$\nu_k^{max} = \nu_k + O\left(\frac{1}{p}\right) \quad \text{and} \quad \nu_k^{min} = \nu_k + O\left(\frac{1}{p}\right)$$

Furthermore, letting ω denote the canonical bundle, we set

$$\theta = \frac{\deg \omega_{Y_p}}{\deg Y_p}$$

which is constant for $p \gg 0$ by the earlier discussion.

Lastly, for any sheaf \mathcal{F} on Y_p we write $h^i(\mathcal{F})$ or $h^i(Y_p, \mathcal{F})$ for $\dim_k H^i(Y_p, \mathcal{F})$.

Asymptotic Lemmas

We first prove a lemma on the cohomology of the twisted bundles $\mathcal{S}^q(m)$ in various ranges of m . Both the lemma and its proof are in direct analogy with Proposition 3.4 of [1], but as now we have that p , not n , is going to infinity, some more care must be taken. In particular, note that we cannot use strong HN filtrations as n is fixed. Instead we compare the filtration to that of the original bundle using the results of Trivedi described in Section 1.

In the proofs of the asymptotic parts of the next few results, we assume that p has been taken large enough so that the genus and degree of Y_p equal those of Y , and we denote them by g and $\deg Y$, respectively. We also assume that p is large enough so that the slopes μ_k and normalized slopes ν_k are constant and that $\deg \omega_{Y_p} = \deg \omega_Y$.

Note that

$$q\nu_k^{\max} \leq q\nu_k^{\min} < q\nu_k^{\min} + \theta \leq q\nu_{k+1}^{\max}$$

Lemma 2.2. *Let \mathcal{S} be a bundle on Y . With the notation above (and setting $\nu_{t+1} = \infty$), one has for $1 \leq k \leq t$:*

(i) *If $m < q\nu_{k+1}^{\max}$, then*

$$H^0(Y_p, \mathcal{S}^q(m)) = H^0(Y_p, \mathcal{S}_k^q(m))$$

In particular, if $m < q\nu_1^{\max}$, then $H^0(Y_p, \mathcal{S}^q(m)) = 0$.

(ii) *If $q\nu_k^{\min} + \theta < m$, then*

$$H^1(Y_p, \mathcal{S}_k^q(m)) = 0$$

(iii) *For $q\nu_k^{\max} \leq m \leq q\nu_k^{\min} + \theta$, one has*

$$\sum_{m=\lceil q\nu_k^{\max} \rceil}^{\lfloor q\nu_k^{\min} + \theta \rfloor} h^1(Y_p, \mathcal{S}_k^q(m)) = O\left(\frac{q^2}{p}\right)$$

In particular, setting $k = t$ and noting that $\mathcal{S}_t = \mathcal{S}$, one sees that (i), (ii) and (iii) yield the following.

Corollary 2.3.

$$\sum_{m=\lceil q\nu_t^{\max} \rceil}^{\infty} h^1(Y_p, \mathcal{S}^q(m)) = O\left(\frac{q^2}{p}\right)$$

Proof of Lemma 2.2. (i) Consider the exact sequence

$$0 \longrightarrow \mathcal{S}_k^q(m) \longrightarrow \mathcal{S}^q(m) \longrightarrow \mathcal{S}^q/\mathcal{S}_k^q(m) \longrightarrow 0.$$

When $m < q\nu_{k+1}^{\max} = \frac{-\mu_{k+1}^{\max}}{\deg Y_p}$, we have

$$\mu_{\max}(\mathcal{S}^q/\mathcal{S}_k^q(m)) = \mu_{\max}(\mathcal{S}^q/\mathcal{S}_k^q) + m \deg Y_p = \mu_{k+1}^{\max} + m \deg Y_p < 0$$

where the second equality is due to the fact that the HN filtration of \mathcal{S}^q/S_k^q is obtained via quotients from the portion of the HN filtration of \mathcal{S}^q that contains \mathcal{S}_k^q . Thus $H^0(Y_p, \mathcal{S}^q/S_k^q(m)) = 0$, and the result follows from the long exact sequence of cohomology.

(ii) By Serre duality,

$$H^1(Y_p, \mathcal{S}_k^q(m)) \cong H^0(Y_p, \mathcal{S}_k^q(m)^\vee \otimes \omega_{Y_p})$$

But when $m > q\nu_k^{\min} + \theta = \frac{-\mu_k^{\min} + \deg \omega_{Y_p}}{\deg Y_p}$, we have

$$\begin{aligned} \mu_{\max}(\mathcal{S}_k^q(m)^\vee \otimes \omega_{Y_p}) &= -\mu_{\min}(\mathcal{S}_k^q(m)) + \mu(\omega_{Y_p}) \\ &= -(\mu_k^{\min} + m \deg Y_p) + \deg \omega_{Y_p} < 0 \end{aligned}$$

and so $H^0(Y_p, \mathcal{S}_k^q(m)^\vee \otimes \omega_{Y_p}) = 0$.

(iii) Since for $p \gg 0$ the bundle \mathcal{S}_k on Y_p is the specialization (reduction mod p) of the corresponding subbundle in the HN filtration of the syzygy bundle in characteristic zero, there exist integers $\alpha_1, \dots, \alpha_s$ (independent of p) and surjections of sheaves on Y_p

$$\bigoplus_{j=1}^s \mathcal{O}(\alpha_j) \longrightarrow \mathcal{S}_k \longrightarrow 0$$

for all $p \gg 0$. Applying the Frobenius pullback $(F^*)^n$, twisting by m , and taking cohomology yields surjections

$$\bigoplus_{j=1}^s H^1(Y_p, \mathcal{O}(q\alpha_j + m)) \longrightarrow H^1(Y_p, \mathcal{S}_k^q(m)) \longrightarrow 0$$

Therefore it is enough to show that for any fixed integer α

$$\sum_{m=\lceil q\nu_k^{\max} \rceil}^{\lfloor q\nu_k^{\min} + \theta \rfloor} h^1(Y_p, \mathcal{O}(q\alpha + m)) = O\left(\frac{q^2}{p}\right)$$

Reindexing and setting $L_0 = q\alpha + \lceil q\nu_k^{\max} \rceil$ and $L_1 = q\alpha + \lfloor q\nu_k^{\min} + \theta \rfloor$ yields the sum

$$\sum_{l=L_0}^{L_1} h^1(Y_p, \mathcal{O}(l))$$

For those p for which $L_0 \geq 0$, this sum is bounded by Remark 2.4 below. So, we may assume that $L_0 < 0$. In that case, Remark 2.4 again yields that the sum of the terms

with $\ell \geq 0$ is bounded independent of p , and so, setting $L = \min(L_1, -1)$, we get

$$\sum_{l=L_0}^{L_1} h^1(Y_p, \mathcal{O}(l)) = \sum_{l=L_0}^L h^1(Y_p, \mathcal{O}(l)) + O(1)$$

In this remaining range, $h^0(Y_p, \mathcal{O}(l)) = 0$ since $l < 0$ and so the Riemann-Roch theorem yields the sum

$$\sum_{l=L_0}^L (-l \deg Y - (1-g)) + O(1) = -\frac{\deg Y}{2}(L-L_0+1)(L+L_0) - (1-g)(L-L_0+1) + O(1)$$

where we have used the following summation formula

$$\sum_{l=a}^b l = \frac{(b-a+1)(b+a)}{2} \quad \text{for any } a \leq b \in \mathbb{Z}$$

Now, since $\nu_k^{\min} = \nu_k + O(\frac{1}{p})$ and $\nu_k^{\max} = \nu_k + O(\frac{1}{p})$ by Corollary 2.1, we have

$$|L + L_0| \leq |L_1| + |L_0| \leq |q\alpha + q\nu_k^{\min} + \theta| + |q\alpha + q\nu_k^{\max}| + 2 = O(q)$$

and more crucially

$$\begin{aligned} 0 \leq L - L_0 + 1 &\leq L_1 - L_0 + 1 = \lfloor q\nu_k^{\min} + \theta \rfloor - \lceil q\nu_k^{\max} \rceil + 1 \\ &\leq q(\nu_k^{\min} - \nu_k^{\max}) + \theta + 1 = O\left(\frac{q}{p}\right) \end{aligned}$$

Plugging these two estimates in above yields the desired result. \square

The following variation of Serre's Vanishing Theorem is used in the proof above.

Remark 2.4. *Note that for a locally free sheaf \mathcal{F} on our family $\text{Proj } R \rightarrow \text{Spec } \mathbb{Z}$ there exists an $M > 0$ (independent of p) such that*

$$H^1(Y_p, \mathcal{F}_p(m)) = 0, \text{ for all } m \geq M$$

and

$$\sum_{m=0}^M h^1(Y_p, \mathcal{F}_p(m)) = O(1)$$

For the generic fiber $Y_{\mathbb{Q}}$ there exists such a bound by Serre vanishing ([8, Theorem III.5.2]). By semicontinuity ([8, Theorem III.12.8]) it follows that $H^1(Y_p, \mathcal{F}_p(M)) = 0$ for almost all primes p , and by the surjections $H^1(Y_p, \mathcal{F}_p(m)) \rightarrow H^1(Y_p, \mathcal{F}_p(m+1))$ this is also true for all larger twists. The second statement follows also from semicontinuity.

As a first step, we now use the lemma above to prove

Lemma 2.5. *For any integer k with $1 \leq k \leq t-1$, let $R = \sum_{i=1}^k r_i$ and $D = \sum_{i=1}^k r_i \nu_i$. Then*

$$\sum_{m=\lceil q\nu_k^{max} \rceil}^{\lceil q\nu_{k+1}^{max} \rceil-1} h^0(Y_p, \mathcal{S}^q(m)) = q^2 \deg Y \left(\frac{R}{2}(\nu_{k+1}^2 - \nu_k^2) - D(\nu_{k+1} - \nu_k) \right) + O\left(\frac{q^2}{p}\right)$$

Proof. By Lemma 2.2(i), in this range for m , one has $h^0(Y_p, \mathcal{S}^q(m)) = h^0(Y_p, \mathcal{S}_k^q(m))$. Applying the Riemann-Roch theorem then gives

$$\sum_{m=\lceil q\nu_k^{max} \rceil}^{\lceil q\nu_{k+1}^{max} \rceil-1} h^0(Y_p, \mathcal{S}^q(m)) = \sum_{m=\lceil q\nu_k^{max} \rceil}^{\lceil q\nu_{k+1}^{max} \rceil-1} (\deg \mathcal{S}_k^q(m) + (\text{rank } \mathcal{S}_k^q)(1-g) + h^1(Y_p, \mathcal{S}_k^q(m)))$$

By parts (ii) and (iii) of Lemma 2.2, $\sum h^1(Y_p, \mathcal{S}_k^q(m)) = O(\frac{q^2}{p})$. Also, since $\text{rank } \mathcal{S}_k^q = \text{rank } \mathcal{S}_k$, one has $\sum (\text{rank } \mathcal{S}_k^q)(1-g) = O(q)$. Furthermore, by additivity of slopes on tensor products

$$\begin{aligned} \deg \mathcal{S}_k^q(m) &= \deg \mathcal{S}_k^q + (\text{rank } \mathcal{S}_k^q)(\deg \mathcal{O}(m)) \\ &= q \deg \mathcal{S}_k + (\text{rank } \mathcal{S}_k)m \deg Y \\ &= q \sum_{i=1}^k r_i \mu_i + m \deg Y \sum_{i=1}^k r_i \\ &= \deg Y \left(-q \sum_{i=1}^k r_i \nu_i + m \sum_{i=1}^k r_i \right) \\ &= \deg Y(mR - qD) \end{aligned}$$

Therefore the sum becomes

$$\begin{aligned} &\sum_{m=\lceil q\nu_k^{max} \rceil}^{\lceil q\nu_{k+1}^{max} \rceil-1} \deg Y(mR - qD) + O\left(\frac{q^2}{p}\right) \\ &= \deg Y \left(\frac{R}{2}(\lceil q\nu_{k+1}^{max} \rceil - \lceil q\nu_k^{max} \rceil)(\lceil q\nu_{k+1}^{max} \rceil + \lceil q\nu_k^{max} \rceil - 1) - qD(\lceil q\nu_{k+1}^{max} \rceil - \lceil q\nu_k^{max} \rceil) \right) + O\left(\frac{q^2}{p}\right) \end{aligned}$$

But $\nu_k^{max} = \nu_k + O(\frac{1}{p})$ for each k by Corollary 2.1, and so the sum indeed simplifies to

$$\deg Y \left(\frac{R}{2} q^2(\nu_{k+1}^2 - \nu_k^2) - Dq^2(\nu_{k+1} - \nu_k) \right) + O\left(\frac{q^2}{p}\right)$$

as desired. \square

3. MAIN RESULT

Now we return to the basic setting of this paper described at the start of Section 1. Recall that pulling back the exact sequence on Y_p

$$0 \longrightarrow \mathcal{S} \longrightarrow \bigoplus_{i=1}^s \mathcal{O}(-d_i) \xrightarrow{f_1, \dots, f_s} \mathcal{O} \longrightarrow 0$$

along the absolute Frobenius morphism n times (with a subsequent twist by $m \in \mathbb{Z}$) yields the long exact sequence of cohomology

$$0 \longrightarrow H^0(Y_p, \mathcal{S}^q(m)) \longrightarrow \bigoplus_{i=1}^s H^0(Y_p, \mathcal{O}(m - qd_i)) \xrightarrow{f_1^q, \dots, f_s^q} H^0(Y_p, \mathcal{O}(m)) \longrightarrow \dots$$

where \mathcal{S}^q denotes the pullback bundle $(F^*)^n(\mathcal{S}) = \text{Syz}(f_1^q, \dots, f_s^q)$. When R_p is normal, one has that $H^0(Y_p, \mathcal{O}(n)) \cong R_n$ for all $n \in \mathbb{N}$, and so the cokernel of f_1^q, \dots, f_s^q is precisely the m -th graded piece of $R_p/I_p^{[q]}$.

For the proof of the main theorem, we will use the results from the previous section to analyze the cohomologies of $\mathcal{S}^q(m)$. As for the cohomologies of the twists of the structure sheaf, we need the following ingredient. Note that although the statement looks like that of Lemma 2.2 of [1], that result cannot be applied here: For one thing, ν_t^{\max} is not a fixed number, and, even more crucially, ours is an asymptotic statement as $p \rightarrow \infty$, not as $n \rightarrow \infty$. Yet the proof is essentially the same, with these modifications in mind.

Lemma 3.1.

$$\sum_{m=0}^{\lceil q\nu_t^{\max} \rceil - 1} h^0(Y_p, \mathcal{O}(m)) = q^2 \frac{\deg Y}{2} \nu_t^2 + O\left(\frac{q^2}{p}\right)$$

$$\sum_{m=0}^{\lceil q\nu_t^{\max} \rceil - 1} h^0(Y_p, \mathcal{O}(m - qd_i)) = q^2 \frac{\deg Y}{2} (\nu_t - d_i)^2 + O\left(\frac{q^2}{p}\right)$$

Proof. As in Section 2, we assume that p has been taken large enough so that the genus and degree of Y_p equal those of Y , and we denote them by g and $\deg Y$, respectively.

We prove the second statement; the proof of the first is similar. By the Riemann-Roch theorem, one has

$$\begin{aligned}
\sum_{m=0}^{\lceil q\nu_t^{max} \rceil - 1} h^0(\mathcal{O}(m - qd_i)) &= \sum_{m=qd_i}^{\lceil q\nu_t^{max} \rceil - 1} h^0(\mathcal{O}(m - qd_i)) \\
&= \sum_{m=0}^{\lceil q\nu_t^{max} \rceil - 1} (m - qd_i) \deg Y + (1 - g) + h^1(\mathcal{O}(m - qd_i)) \\
&= \sum_{l=0}^{\lceil q\nu_t^{max} \rceil - qd_i - 1} (l \deg Y + (1 - g) + h^1(\mathcal{O}(l))) \\
&= \frac{\deg Y}{2} (\lceil q\nu_t^{max} \rceil - qd_i) (\lceil q\nu_t^{max} \rceil - qd_i - 1) \\
&\quad + (1 - g) (\lceil q\nu_t^{max} \rceil - qd_i) + \sum_{l=0}^{\lceil q\nu_t^{max} \rceil - qd_i - 1} h^1(\mathcal{O}(l))
\end{aligned}$$

The last term is $O(1)$ by Remark 2.4. Furthermore, since

$$\lceil q\nu_t^{max} \rceil = q\nu_t^{max} + O(1) = q\nu_t + O\left(q \cdot \frac{1}{p}\right)$$

by Corollary 2.1, the second term is $O(q)$ and the first term becomes

$$q^2 \frac{\deg Y}{2} (\nu_t - d_i)^2 + O\left(\frac{q^2}{p}\right)$$

as desired. \square

We are now ready to compute the desired limit.

Theorem 3.2. *Let R be a standard-graded flat domain over \mathbb{Z} such that almost all fiber rings $R_p = R \otimes_{\mathbb{Z}} \mathbb{Z}/p\mathbb{Z}$ are geometrically normal 2-dimensional domains and let $I = (f_1, \dots, f_s)$ be a homogeneous R_+ -primary ideal. Set r_k and ν_k to be the ranks and normalized slopes of the quotients in the HN filtration of the syzygy bundle over $Y = \text{Proj } R_{\mathbb{Q}}$. For any fixed integer $n \geq 1$, setting $q = p^n$, one has*

$$\frac{\ell(R_p/I_p^{[q]})}{q^2} = \frac{\deg Y}{2} \left(\sum_{k=1}^t r_k \nu_k^2 - \sum_{i=1}^s d_i^2 \right) + O\left(\frac{1}{p}\right)$$

where $R_p = R \otimes_{\mathbb{Z}} \mathbb{Z}/p\mathbb{Z}$, $I_p^{[q]} = (f_1^q, \dots, f_s^q)R_p$.

Proof. The long exact sequence of cohomology for the exact sequence

$$0 \longrightarrow \mathcal{S}^q(m) \longrightarrow \bigoplus_{i=1}^s \mathcal{O}(m - qd_i) \xrightarrow{f_1^q, \dots, f_s^q} \mathcal{O}(m) \longrightarrow 0$$

yields the containment

$$\text{Coker } H^0(f_1^q, \dots, f_s^q) = R_p/I_p^{[q]} \subseteq H^1(Y_p, \mathcal{S}^q(m)).$$

Therefore by Corollary 2.3

$$\ell(R_p/I_p^{[q]}) = \sum_{m=0}^{\infty} \ell(R_p/I_p^{[q]})_m = \sum_{m=0}^{\lceil q\nu_t^{\max} \rceil - 1} \ell(R_p/I_p^{[q]})_m + O\left(\frac{q^2}{p}\right)$$

The beginning of the long exact sequence then yields

$$\ell(R_p/I_p^{[q]}) = \sum_{m=0}^{\lceil q\nu_t^{\max} \rceil - 1} \left(h^0(\mathcal{O}(m)) - \sum_{i=1}^s h^0(\mathcal{O}(m - qd_i)) + h^0(\mathcal{S}^q(m)) \right) + O\left(\frac{q^2}{p}\right)$$

After changing the order of summation, one may apply Lemma 3.1 to get

$$= q^2 \frac{\deg Y}{2} (\nu_t^2 - \sum_{i=0}^s (\nu_t - d_i)^2) + \sum_{m=0}^{\lceil q\nu_t^{\max} \rceil - 1} h^0(\mathcal{S}^q(m)) + O\left(\frac{q^2}{p}\right)$$

Plugging in the result of Lemma 2.5, using the fact that $h^0(\mathcal{S}^q(m)) = 0$ for $m < \lceil q\nu_1^{\max} \rceil$ by Lemma 2.2(i), and simplifying as in Theorem 3.6 of [1] yields the desired result. \square

This finally brings us to our main goal: The expression on the right hand side of the equation in Theorem 3.2 is equal to the limit Hilbert-Kunz multiplicity

$$e_{\text{HK}}^{\infty}(I, R) \stackrel{\text{def}}{=} \lim_{p \rightarrow \infty} e_{\text{HK}}(I_p, R_p)$$

as proved by Trivedi in [18]. Therefore, we obtain the following consequence.

Corollary 3.3. *With the notation as above, for any fixed $n \geq 1$ one has*

$$\frac{\ell(R_p/I_p^{[p^n]})}{(p^n)^2} = e_{\text{HK}}^{\infty}(I, R) + O\left(\frac{1}{p}\right)$$

In particular,

$$e_{\text{HK}}^{\infty}(I, R) = \lim_{p \rightarrow \infty} \frac{\ell(R_p/I_p^{[p^n]})}{(p^n)^2}$$

In fact, Trivedi shows that for these rings

$$e_{\text{HK}}(I_p, R_p) = e_{\text{HK}}^\infty(I, R) + O\left(\frac{1}{p}\right)$$

It is interesting to note that the bound $O(\frac{1}{p})$ on the speed of convergence is of the same order as in Trivedi's result.

Example 3.4. *The following example can be found in Monsky's paper [13]. For the ring $R = \mathbb{Z}/p\mathbb{Z}[x, y, z]/(x^4 + y^4 + z^4)$ and the homogeneous maximal ideal $I = (x, y, z)$, one has*

$$e_{\text{HK}}(I, R) = \begin{cases} 3 + \frac{1}{p^2} & p \equiv 3, 5 \pmod{8} \\ 3 & p \equiv 1, 7 \pmod{8} \end{cases}$$

It is not clear whether all these results are optimal since we have not been able to find an example with the slower convergence rate of $O(\frac{1}{p})$. See also Example 4.2 for diagonal hypersurfaces.

4. DIAGONAL HYPERSURFACES

Unpublished results of Gessel and Monsky [14] show that $e_{\text{HK}}^\infty(\mathfrak{m}, R)$ exists also for any diagonal hypersurface over \mathbb{Z} , that is, a ring of the form

$$R = \frac{\mathbb{Z}[x_1, \dots, x_s]}{(x_1^{d_1} + \dots + x_s^{d_s})}$$

with respect to the homogeneous ideal \mathfrak{m} generated by the variables. In this section we show how the proof simultaneously gives an affirmative answer to the question in our introduction for these rings, i.e., that for any fixed $n \geq 1$

$$e_{\text{HK}}^\infty(\mathfrak{m}, R) = \lim_{p \rightarrow \infty} \frac{\ell(R_p/\mathfrak{m}_p^{[p^n]})}{(p^n)^d}$$

Furthermore, we then use these methods to provide examples to show that a certain naive limit in characteristic zero analogous to the one used in positive characteristic to define the HK multiplicity does not give the same answer in general.

Affirmative answer for diagonal hypersurface rings

We repeat a small part of the arguments from [14] here to show how it yields the result above. It uses the machinery developed by Han and Monsky in [7] for computing HK multiplicities of diagonal hypersurfaces in positive characteristic. For the notation,

we generally refer the reader to their paper, although the necessities are repeated here. For positive integers k_1, \dots, k_s and field $F = \mathbb{Z}/p\mathbb{Z}$ define

$$\begin{aligned} D_F(k_1, \dots, k_s) &= \dim_F F[x_1, \dots, x_{s-1}]/(x_1^{k_1}, \dots, x_{s-1}^{k_{s-1}}, (x_1 + \dots + x_{s-1})^{k_s}) \\ &= \dim_F F[x_1, \dots, x_s]/(x_1^{k_1}, \dots, x_s^{k_s}, x_1 + \dots + x_s) \end{aligned}$$

In [14], Gessel and Monsky show that, for any p and n , there are inequalities

$$(4.1) \quad d_1 \cdots d_s \frac{D_F(\lfloor \frac{p}{d_1} \rfloor, \dots, \lfloor \frac{p}{d_s} \rfloor)}{p^d} \leq \frac{\ell(R_p/\mathfrak{m}_p^{[p^n]})}{(p^n)^d} \leq d_1 \cdots d_s \frac{D_F(\lfloor \frac{p}{d_1} \rfloor + 1, \dots, \lfloor \frac{p}{d_s} \rfloor + 1)}{p^d}$$

As the outside terms are independent of n , taking the limit as n goes to infinity yields inequalities

$$(4.2) \quad d_1 \cdots d_s \frac{D_F(\lfloor \frac{p}{d_1} \rfloor, \dots, \lfloor \frac{p}{d_s} \rfloor)}{p^d} \leq e_{\text{HK}}(\mathfrak{m}_p, R_p) \leq d_1 \cdots d_s \frac{D_F(\lfloor \frac{p}{d_1} \rfloor + 1, \dots, \lfloor \frac{p}{d_s} \rfloor + 1)}{p^d}$$

they then prove that, as p goes to infinity, the outside terms both converge to the same limit, and in fact, both equal

$$g\left(\frac{1}{d_1}, \dots, \frac{1}{d_s}\right) + O\left(\frac{1}{p}\right)$$

for the function $g: [0, 1]^s \rightarrow \mathbb{Q}$ defined as follows: for any numbers $x_1, \dots, x_s \in [0, 1]$, set

$$(4.3) \quad g(x_1, \dots, x_s) = \frac{1}{2^{s-1}(s-1)!} \sum_{\lambda \in \mathbb{Z}} g_\lambda(x_1, \dots, x_s)$$

where

$$(4.4) \quad g_\lambda(x_1, \dots, x_s) = \sum_{\epsilon_i = \pm 1 \text{ and } \sum \epsilon_i x_i \geq 2\lambda} \epsilon_1 \cdots \epsilon_s (\epsilon_1 x_1 + \dots + \epsilon_s x_s - 2\lambda)^{s-1}$$

Note that g is well-defined since $g_\lambda = 0$ for $|\lambda| \gg 0$. But then the middle terms in both inequalities (4.1) and (4.2) go to the same limit (at the same rate) as well.

In summary, we arrive at the following conclusion.

Theorem 4.1 (Monsky). *For any diagonal hypersurface ring*

$$R = \frac{\mathbb{Z}[x_1, \dots, x_s]}{(x_1^{d_1} + \dots + x_s^{d_s})} \quad d_i \geq 2 \text{ for all } i$$

with homogeneous maximal ideal \mathfrak{m} and any fixed n , one has

$$e_{\text{HK}}^\infty(\mathfrak{m}, R) = e_{\text{HK}}(\mathfrak{m}_p, R_p) + O\left(\frac{1}{p}\right) = \frac{\ell(R_p/\mathfrak{m}_p^{[p^n]})}{(p^n)^d} + O\left(\frac{1}{p}\right)$$

Furthermore,

$$e_{HK}^\infty(\mathfrak{m}, R) = g\left(\frac{1}{d_1}, \dots, \frac{1}{d_s}\right)$$

where the function g is defined as above in (4.3) and (4.4).

Note that, as for the case of homogeneous coordinate rings over smooth curves in the previous section (see Corollary 3.3 and the discussion after it), the bounds on the rates of convergence of the various quantities to $e_{HK}^\infty(\mathfrak{m}, R)$ are the same. We do not know in this case either whether the bound $O(\frac{1}{p})$ on the speed of convergence is optimal.

Example 4.2. *The diagonal hypersurface ring in Example 3.4 satisfies*

$$e_{HK}(I, R) = e_{HK}^\infty(I, R) + O\left(\frac{1}{p^2}\right)$$

The same is true of the following example worked out by Chang in her thesis [4] using the techniques from [7]. For the ring $R = \mathbb{Z}/p\mathbb{Z}[w, x, y, z]/(w^4 + x^4 + y^4 + z^4)$ and the homogeneous maximal ideal $I = (w, x, y, z)$, one has

$$e_{HK}(I, R) = \frac{2}{3} \left(\frac{8p^3 + 4p - 12}{2p^3 - p \pm 1} \right)$$

according as $p \equiv 1(4)$ or $p \equiv 3(4)$. Therefore, one finds that

$$e_{HK}(I, R) = \frac{8}{3} + O\left(\frac{1}{p^2}\right)$$

We do not know an example with the slower converge rate of $O(\frac{1}{p})$.

Limits in characteristic zero

Now we turn to using the results of Gessel and Monsky to examine why a certain naive limit in characteristic zero fails to give the same answer. Given a local ring R of equicharacteristic zero with maximal ideal \mathfrak{m} , it might be tempting (in analogy with the definition of HK multiplicity in positive characteristic) to take a set of generators x_1, \dots, x_r of \mathfrak{m} and to look at the following limit (if it exists)

$$e_{\text{naive}}^\infty = \lim_{N \rightarrow \infty} \frac{\ell(R_{\mathbb{Q}}/(x_1^N, \dots, x_r^N))}{N^d}$$

Unfortunately, this does not yield $e_{HK}^\infty(\mathfrak{m}, R)$ in general. In fact, their unpublished work [14] enables one to compute this limit as well for diagonal hypersurfaces. Indeed, if we

set

$$R = \frac{\mathbb{Z}[x_1, \dots, x_s]}{(x_1^{d_1} + \dots + x_s^{d_s})}$$

then by Lemma 1 of [14] in view of Theorem 2.14 of [7], this limit equals the $\lambda = 0$ term of $g(\frac{1}{d_1}, \dots, \frac{1}{d_s})$, that is

$$e_{\text{naive}}^\infty = \frac{1}{2^{s-1}(s-1)!} g_0$$

Therefore, whenever there are nonzero g_λ terms in $g(\frac{1}{d_1}, \dots, \frac{1}{d_s})$ for some $\lambda \neq 0$, one might have $e_{\text{naive}}^\infty \neq e_{\text{HK}}^\infty(\mathfrak{m}, R)$ by Monsky's Theorem 4.1. We give explicit examples below.

We begin with an example in which a minimal set of generators is used for \mathfrak{m} in computing e_{naive}^∞ and yet one still does not obtain $e_{\text{HK}}^\infty(\mathfrak{m}, R)$ as the limit. This is the “smallest” example of which we know.

Example 4.3. *In the notation above, let $s = 5$ and $d_i = 2$ for all i , that is, take the ring*

$$R = \mathbb{Z}[x_1, \dots, x_5]/(x_1^2 + \dots + x_5^2)$$

Then, writing g_λ for $g_\lambda(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}, \frac{1}{2}, \frac{1}{2})$, we have $g_\lambda = 0$ whenever $|\lambda| \geq 2$ and

$$g_1 = g_{-1} = \left(\frac{1}{2} + \frac{1}{2} + \frac{1}{2} + \frac{1}{2} + \frac{1}{2} - 2\right)^4 = \left(\frac{1}{2}\right)^4$$

Monsky's Theorem 4.1 then yields

$$e_{\text{HK}}^\infty(\mathfrak{m}, R) = \frac{2}{4!} \left(g_0 + 2\left(\frac{1}{2}\right)^4\right)$$

whereas

$$e_{\text{naive}}^\infty = \frac{2}{4!} g_0$$

Now we present a simpler example using similar ideas. It has the drawback though that minimal generating sets were not used when computing e_{naive}^∞ .

Example 4.4. *In the notation above, let $s = 3$ and $d_i = 1$ for all i , that is, take the ring*

$$R = \mathbb{Z}[x_1, x_2, x_3]/(x_1 + x_2 + x_3)$$

Then Theorem 2.14 of [7] shows that $R_{\mathbb{Q}}/(x_1^N, x_2^N, x_3^N)$ has dimension equal to $\lceil \frac{3}{4}N^2 \rceil$. (Monsky pointed out to us that this can also be proved by a simple argument involving a matrix of binomial coefficients.) So $e_{\text{naive}}^\infty = \frac{3}{4}$. But, as R is isomorphic to the regular ring $\mathbb{Z}[x_1, x_2]$, we know that $e_{\text{HK}}^\infty(\mathfrak{m}, R) = 1$.

Remark 4.5. *It is interesting to compare and contrast these examples to the one given by Buchweitz and Chen in [3]. In contrast to our discussion above in characteristic 0, their results show that in characteristic p the naive limit does not even necessarily exist, even for a fixed choice of generators of the homogeneous maximal ideal. Specifically, for the ring*

$$R_p = \mathbb{Z}/p\mathbb{Z}[x_1, x_2, x_3]/(x_1 + x_2 + x_3)$$

(namely the reduction to characteristic p of the ring in Example 4.4 above) they show that the limit

$$\lim_{N \rightarrow \infty} \frac{\ell(R_p/(x_1^N, x_2^N, x_3^N))}{N^2}$$

does not exist. Indeed for the subsequence $N = p^n$ the limit is just the HK multiplicity, which equals 1 since R_p is regular, but for the subsequence $N = 2p^n$ the limit turns out to equal $\frac{3}{4}$ by an elementary computation.

More generally, the study in characteristic p of how the length of

$$F[x, y]/(f^i, g^j, h^k),$$

where F is a field, depends on i, j and k when f, g and h are fixed was carried out by Teixeira in his thesis [16]; the answer involves “ p -fractals”.

REFERENCES

- [1] H. Brenner, *The rationality of the Hilbert-Kunz multiplicity in graded dimension two*, Math. Ann. **334** (2006), 91–110.
- [2] W. Bruns, *Conic divisor classes over a normal monoid algebra*, Contemp. Math. **390** Commutative algebra and algebraic geometry, (2005), 63–71.
- [3] R.-O. Buchweitz, Q. Chen, *Hilbert-Kunz functions of cubic curves and surfaces* J. Algebra **197** (1997), 246–267.
- [4] Sh.-t. Chang, *The asymptotic behavior of Hilbert-Kunz functions and their generalizations*, thesis, University of Michigan, 1993.
- [5] A. Conca, *Hilbert-Kunz function of monomial ideals and binomial hypersurfaces*, Manuscripta Math. **90** (1996), 287–300.
- [6] K. Eto, *Multiplicity and Hilbert-Kunz multiplicity of monoid rings*, Tokyo J. Math. **25** (2002), 241–245.
- [7] C. Han, P. Monsky, *Some surprising Hilbert-Kunz functions*, Math. Z. **214** (1993), 119–135.
- [8] R. Hartshorne, *Algebraic geometry*, Graduate Texts in Mathematics **52**, Springer-Verlag, New York-Heidelberg, 1977.
- [9] E. Kunz, *Characterization of regular local rings for characteristic p* , Amer. J. Math. **91** (1969), 772–784.

- [10] A. Langer, *Semistable sheaves in positive characteristic*, Ann. Math. **159** (2004), 251–276.
- [11] P. Monsky, *The Hilbert-Kunz function*, Math. Ann. **263** (1983), 43–49.
- [12] P. Monsky, *The Hilbert-Kunz function of a characteristic 2 cubic*, J. Algebra **197** (1997), 268–277.
- [13] P. Monsky, *The Hilbert-Kunz multiplicity of an irreducible trinomial*, J. Algebra **304** (2006), 1101–1107.
- [14] I. M. Gessel, P. Monsky, *The limit as $p \rightarrow \infty$ of the Hilbert-Kunz multiplicity of $\sum x_i^{d_i}$* , arXiv:1007.2004.
- [15] K. Pardue, Thesis, Brandeis University, 1993.
- [16] P. Teixeira, Thesis, Brandeis University, 2002.
- [17] V. Trivedi, *Semistability and Hilbert-Kunz multiplicities for curves*, J. Algebra **284** (2005), 627–644.
- [18] V. Trivedi, *Hilbert-Kunz multiplicity and reduction mod p* , Nagoya Math. J. **185** (2007), 123–141.
- [19] V. Trivedi, *Strong semistability and Hilbert-Kunz multiplicity for singular plane curves*, Contemp. Math. **390** Commutative algebra and algebraic geometry, (2005), 165–173.
- [20] K.-I. Watanabe, *Hilbert-Kunz multiplicity of toric rings*, Proc. Inst. Nat. Sci. (Nihon Univ.) **35** (2000), 173–177.
- [21] K.-I. Watanabe, K. Yoshida, *Hilbert-Kunz multiplicity, McKay correspondence and good ideals in two-dimensional rational singularities*, Manuscripta Math. **104** (2001), 275–294.

FACHBEREICH FÜR MATHEMATIK UND INFORMATIK, UNIVERSITÄT OSNABRÜCK, OSNABRÜCK,
GERMANY

E-mail address: `hbrenner@uni-osnabrueck.de`

MATHEMATICS DEPARTMENT, SYRACUSE UNIVERSITY, SYRACUSE, NY 13224, USA

Current address: Department of Mathematics, University of Louisville, Louisville, KY 40292, USA

E-mail address: `jinjia.li@louisville.edu`

MATHEMATICS DEPARTMENT, SYRACUSE UNIVERSITY, SYRACUSE, NY 13224, USA

E-mail address: `clamille@syr.edu`