

4-3-2007

Complete Shrinking Ricci Solitons have Finite Fundamental Group

William Wylie

University of California - Los Angeles

Follow this and additional works at: <https://surface.syr.edu/mat>



Part of the [Mathematics Commons](#)

Recommended Citation

Wylie, William, "Complete Shrinking Ricci Solitons have Finite Fundamental Group" (2007). *Mathematics - Faculty Scholarship*. 125.
<https://surface.syr.edu/mat/125>

This Article is brought to you for free and open access by the Mathematics at SURFACE. It has been accepted for inclusion in Mathematics - Faculty Scholarship by an authorized administrator of SURFACE. For more information, please contact surface@syr.edu.

Complete Shrinking Ricci Solitons have Finite Fundamental Group

William Wylie

ABSTRACT. We show that if a complete Riemannian manifold supports a vector field such that the Ricci tensor plus the Lie derivative of the metric with respect to the vector field has a positive lower bound, then the fundamental group is finite. In particular, it follows that complete shrinking Ricci solitons and complete smooth metric measure spaces with a positive lower bound on the Bakry-Emery tensor have finite fundamental group. The method of proof is to generalize arguments of Garcia-Rio and Fernandez-Lopez in the compact case.

1. Introduction

In this paper we are interested in studying complete Riemannian manifolds (M, g) with a vector field X such that, for some $\lambda > 0$,

$$(1.1) \quad \text{Ric}_g + \mathcal{L}_X g \geq \lambda g$$

where $\mathcal{L}_X g$ is the Lie derivative of g with respect to the vector field X .

This class of manifolds includes two well known subclasses. The first is the class of complete manifolds such that $\text{Ric}_g + \mathcal{L}_X g = \lambda g$. This is the class of complete Ricci Solitons. The study of Ricci solitons is important in understanding many aspects of Ricci flow [CLN]. The second subclass is the class of smooth metric measure spaces with Bakry-Emery tensor bounded below by λ . This class consists of smooth manifolds that satisfy (1.1) for some gradient vector field $X = \nabla f$. We refer the reader to Lott [L] and Qian [Q] for topological results concerning the Bakry-Emery tensor.

Fernández-López and García-Río [FG] have proven the following Myers type theorem: If a complete manifold (M, g) satisfies (1.1) and the vector field X has bounded norm, then M is compact. It is an immediate corollary that any compact manifold satisfying (1.1) has finite fundamental group. In the case where X is a gradient vector field this was proven earlier by Lott [L].

The assumption of bounded $\|X\|$ is necessary to show M is compact since, for example, Euclidean space with the vector field $X(v) = v, \forall v \in \mathbb{R}^n$ satisfies (1.1). However, we show that the fundamental group must be finite in the noncompact case.

1991 *Mathematics Subject Classification.* 53C20.

Key words and phrases. Ricci Soliton, noncompact manifold, fundamental group.

THEOREM 1.1. *If M is a complete Riemannian manifold satisfying (1.1) then M has finite fundamental group.*

Naber [N] has independently proven Theorem 1.1 under the additional assumption that $X = \nabla f$ and the Ricci curvature is bounded.

There are complete manifolds with positive Ricci curvature and infinite fundamental group. It is an obvious consequence of Theorem 1.1 that the Ricci tensor of these manifolds can not be perturbed by a Lie derivative term to be bounded away from zero.

The proof of Theorem 1.1 is similar to the arguments of Fernández-López and García-Río [FG]. They show that if M satisfies (1.1) and $\|X\|$ is bounded then the integral of the Ricci curvature along every geodesic is infinite. By the Ambrose theorem [A] this implies that the manifold is compact. The main idea of this paper is to replace the Ambrose theorem with an estimate of Hamilton [H] (See Lemma 2.2 below). This estimate allows us to obtain an upper bound on the distance between two points that depends only on the value of $\|X\|$ at each point and an upper bound on the Ricci curvature in a neighborhood of each point (Theorem 2.3). Applying the upper bound to a point in the universal cover of M and its image under a deck transformation yields Theorem 1.1.

2. Proof of Theorem 1.1

We make the following definition to aid the exposition.

DEFINITION 2.1. For any point $p \in M$ define

$$H_p = \max \{0, \sup \{\text{Ric}_y(v, v) : y \in B(p, 1), \|v\| = 1\}\}.$$

We can now state the main lemma.

LEMMA 2.2. *Let (M, g) be a complete Riemannian manifold, let $p, q \in M$ such that $r = d(p, q) > 1$ and let γ be the minimal geodesic from p to q parametrized by arclength, then*

$$\int_0^r \text{Ric}(\gamma'(s), \gamma'(s)) ds \leq 2(n-1) + H_p + H_q.$$

Lemma 2.2 was used by Hamilton [H] to study the change in the distance function on a Riemannian manifold evolving by Ricci flow and also appears in Perelman ([P], Lemma 8.1). We include the proof for completeness.

PROOF. By the second variation of arclength formula, for any piecewise smooth function ϕ with $\phi(0) = \phi(r) = 0$,

$$(2.1) \quad 0 \leq \int_0^r ((n-1)(\phi'(s))^2 - \phi^2(s)\text{Ric}(\gamma'(s), \gamma'(s))) ds.$$

Let ϕ be the function

$$\phi(s) = \begin{cases} s & 0 \leq s \leq 1 \\ 1 & 1 \leq s \leq r-1 \\ r-s & r-1 \leq s \leq r \end{cases}$$

Then, since $\phi(s) = 1$ and $\phi'(s) = 0$ for $1 \leq s \leq r-1$, (2.1) becomes

$$\begin{aligned} 0 \leq & \int_0^1 ((n-1)(\phi'(s))^2) ds + \int_{r-1}^r ((n-1)(\phi'(s))^2) ds - \int_0^1 (\phi^2(s)\text{Ric}(\gamma', \gamma')) ds \\ & - \int_{r-1}^r (\phi^2(s)\text{Ric}(\gamma'(s), \gamma'(s))) ds - \int_1^{r-1} \text{Ric}(\gamma'(s), \gamma'(s)) ds. \end{aligned}$$

Adding $\int_0^r \text{Ric}(\gamma'(s), \gamma'(s)) ds$ to both sides of the equation yields

$$\begin{aligned} \int_0^r \text{Ric}(\gamma'(s), \gamma'(s)) ds & \leq \int_0^1 ((n-1)(\phi'(s))^2) ds + \int_{r-1}^r ((n-1)(\phi'(s))^2) ds \\ (2.2) \quad & + \int_0^1 ((1-\phi^2(s)) \text{Ric}(\gamma'(s), \gamma'(s))) ds \\ & + \int_{r-1}^r ((1-\phi^2(s)) \text{Ric}(\gamma'(s), \gamma'(s))) ds. \end{aligned}$$

We now work out the terms on the right hand side of equation (2.2). Since $|\phi'(s)| = 1$ for $0 \leq s \leq 1$ and $r-1 \leq s \leq r$ we have

$$(2.3) \quad \int_0^1 ((n-1)(\phi'(s))^2) ds + \int_{r-1}^r ((n-1)(\phi'(s))^2) ds = 2(n-1).$$

Moreover, $0 \leq \phi \leq 1$ and $\text{Ric}(\gamma'(s), \gamma'(s)) \leq H_p$ for $0 \leq s \leq 1$, therefore

$$(2.4) \quad \int_0^1 ((1-\phi^2(s)) \text{Ric}(\gamma'(s), \gamma'(s))) ds \leq H_p.$$

Similarly, since $\text{Ric}(\gamma'(s), \gamma'(s)) \leq H_q$ for $r-1 \leq s \leq r$,

$$(2.5) \quad \int_{r-1}^r ((1-\phi^2(s)) \text{Ric}(\gamma'(s), \gamma'(s))) ds \leq H_q.$$

Thus, combining (2.2), (2.3), (2.4), and (2.5) gives the lemma. \square

Using Lemma 2.2 and the arguments in [FG], we can now derive an upper bound on the distance between two points that depends only on $\|X\|$ and H .

THEOREM 2.3. *If (M, g) is a complete manifold satisfying (1.1) then, for any $p, q \in M$,*

$$(2.6) \quad d(p, q) \leq \max \left\{ 1, \frac{1}{\lambda} (2(n-1) + H_p + H_q + 2\|X_p\| + 2\|X_q\|) \right\}.$$

PROOF. Assume that $d(p, q) > 1$ and let γ be the minimal geodesic from p to q . Applying Lemma 2.2 we have

$$(2.7) \quad \int_0^r \text{Ric}(\gamma'(s), \gamma'(s)) ds \leq 2(n-1) + H_p + H_q.$$

On the other hand, by equation (1.1)

$$\begin{aligned} \int_0^r \text{Ric}(\gamma'(s), \gamma'(s)) ds & \geq \int_0^r (\lambda g(\gamma'(s), \gamma'(s)) - \mathcal{L}_X g(\gamma'(s), \gamma'(s))) ds \\ (2.8) \quad & \geq \lambda d(p, q) + 2g_p(X, \gamma'(0)) - 2g_q(X, \gamma'(r)) \\ & \geq \lambda d(p, q) - 2\|X_p\| - 2\|X_q\|. \end{aligned}$$

Where, in the last step, we have used

$$\mathcal{L}_X g(\gamma'(s), \gamma'(s)) = 2 \frac{d}{ds} g(X, \gamma'(s)).$$

Combining (2.7) and (2.8) and solving for $d(p, q)$ gives (2.6). \square

PROOF OF THEOREM 1.1. Let \widetilde{M} be the universal cover of M . \widetilde{M} satisfies (1.1) for the pullback metric and pullback vector field, \widetilde{X} . Fix \tilde{p} in \widetilde{M} and let $h \in \pi_1(M)$ identified as a deck transformation on \widetilde{M} . Note that $B(\tilde{p}, 1)$ and $B(h(\tilde{p}), 1)$ are isometric, thus $H_{\tilde{p}} = H_{h(\tilde{p})}$. Also, $\|\widetilde{X}_{\tilde{p}}\| = \|\widetilde{X}_{h(\tilde{p})}\|$ so by applying Theorem 2.3 to the points \tilde{p} and $h(\tilde{p})$ we obtain

$$d(\tilde{p}, h(\tilde{p})) \leq \max \left\{ 1, \frac{2}{\lambda} \left(n - 1 + H_{\tilde{p}} + 2\|\widetilde{X}_{\tilde{p}}\| \right) \right\} \quad \forall h \in \pi_1(M).$$

Since the right hand side is independent of h , this proves the theorem. \square

Acknowledgements: I would like to thank the authors of [FG] for providing me with a copy of their work, Ben Chow for encouraging me to study Ricci solitons and for his interest in this work, and Guofang Wei and Peter Petersen for many helpful discussions. This work was partially completed while at MSRI.

References

- [A] W. Ambrose, *A theorem of Myers*, Duke Math. J. **24** (1957), 345–348. MR MR0089464 (19,680c)
- [CLN] B. Chow, P. Lu, and L. Ni, *Hamilton's ricci flow*, Graduate Studies in Mathematics, vol. 77, American Mathematical Society, Providence, RI, 2006.
- [FG] M. Fernández-López and E. García-Río, *A remark on compact ricci solitons*, preprint.
- [H] R. Hamilton, *The formation of singularities in the Ricci flow*, Surveys in differential geometry, Vol. II (Cambridge, MA, 1993), Int. Press, Cambridge, MA, 1995, pp. 7–136. MR MR1375255 (97e:53075)
- [L] J. Lott, *Some geometric properties of the Bakry-Émery-Ricci tensor*, Comment. Math. Helv. **78** (2003), no. 4, 865–883. MR MR2016700 (2004i:53044)
- [N] A. Naber, *Some geometry and analysis on Ricci solitons*, arXiv:math.DG/0612532.
- [P] G. Perelman, *The entropy formula for the Ricci flow and its geometric applications*, arXiv: math.DG/0211159.
- [Q] Z. Qian, *Estimates for weighted volumes and applications*, Quart. J. Math. Oxford Ser. (2) **48** (1997), no. 190, 235–242. MR MR1458581 (98e:53058)

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, LOS ANGELES, CA 90095
E-mail address: wylie@math.ucla.edu