

Syracuse University

**SURFACE**

---

Syracuse University Honors Program Capstone Projects    Syracuse University Honors Program Capstone Projects

---

Spring 5-2016

## Structure-Guided Site-Directed Mutagenesis of the Bacterial ATP Synthase's Epsilon Subunit

Mariam Bhatti

Follow this and additional works at: [https://surface.syr.edu/honors\\_capstone](https://surface.syr.edu/honors_capstone)



Part of the [Biology Commons](#)

---

### Recommended Citation

Bhatti, Mariam, "Structure-Guided Site-Directed Mutagenesis of the Bacterial ATP Synthase's Epsilon Subunit" (2016). *Syracuse University Honors Program Capstone Projects*. 966.

[https://surface.syr.edu/honors\\_capstone/966](https://surface.syr.edu/honors_capstone/966)

This Honors Capstone Project is brought to you for free and open access by the Syracuse University Honors Program Capstone Projects at SURFACE. It has been accepted for inclusion in Syracuse University Honors Program Capstone Projects by an authorized administrator of SURFACE. For more information, please contact [surface@syr.edu](mailto:surface@syr.edu).

Structure-Guided Site-Directed Mutagenesis of the Bacterial ATP Synthase's  $\epsilon$  Subunit

A Capstone Project Submitted in Partial Fulfillment of the  
Requirements of the Renée Crown University Honors Program at  
Syracuse University

Mariam Bhatti

Candidate for Bachelor of Science Degree  
and Renée Crown University Honors  
May 2016

Honors Capstone Project in Biology

Capstone Project Advisor: \_\_\_\_\_  
Thomas Duncan

Capstone Project Reader: \_\_\_\_\_  
Samuel Chan

Honors Director: \_\_\_\_\_  
Stephen Kuusisto, Director

### Abstract

Adenosine triphosphate (ATP) contains energy-rich phosphoanhydride bonds that provide the energy needed for many cellular processes. F-type ATP synthase is found in bacteria, chloroplasts, and mitochondria, having a conserved function to catalyze the synthesis and hydrolysis of ATP. ATP synthase is a membrane bound rotary motor enzyme, with coupled rotation between its two distinct complexes  $F_0$  and  $F_1$ . In bacteria and chloroplasts, the  $\epsilon$ -subunit's C-terminal Domain ( $\epsilon$ CTD) has a distinct regulatory function that is absent in mitochondria. Determining the inhibitory interactions of  $\epsilon$  is important in understanding its physiological functions and for potential targeting of  $\epsilon$ 's bacteria-specific inhibition for development of new antibiotics. Guided by a high-resolution structure of  $\epsilon$  inhibition catalytic complex, in this study I use site-specific mutagenesis of the  $\epsilon$ CTD in *Escherichia coli* (*E. coli*) to investigate interactions and make mutations at regions important for  $\epsilon$  inhibition. I then analyze the effects of these mutants through phenotypic growth and ATP hydrolysis assays.

### **Executive Summary**

Adenosine triphosphate (ATP) is a primary energy carrier in cells of all known kingdoms. ATP synthase, an evolutionarily conserved membrane bound enzyme, is responsible for making most cellular ATP in animals, plants, and in bacteria. The ATP synthase is essential for a number of pathogenic bacteria. Although the bacterial and human ATP synthase share homologs, the C-terminal domain of bacterial epsilon subunit ( $\epsilon$ CTD) has an inhibitory interaction that is absent in humans. If we can understand how this bacterial-specific inhibition works, we can target the enzyme to kill bacteria without harming our own enzyme. This makes the  $\epsilon$ CTD a good potential target for antibiotic development.

The model organism being investigated in this study is *Escherichia coli* (*E. coli*), a gram-negative bacteria found in the human gut, and used for over 30 years to study ATP synthase. All subunits of *E. coli* ATP synthase have homologs in ATP synthases of other bacteria, chloroplasts, and mitochondria. Exploring the interactions between  $\epsilon$ CTD of F<sub>1</sub>-ATPase is crucial to understanding the inhibition of ATP synthesis and hydrolysis in bacteria. In this study, I will investigate two sites on the  $\epsilon$ CTD that may be important in inhibiting ATP synthesis and hydrolysis in bacteria.


**Table of Contents**

Abstract	1
Executive Summary	2
Introduction	4
Methods	7
Results	12
Discussion	19
Acknowledgements	22
References	23

## Introduction


The synthesis of ATP from ADP and  $P_i$  is facilitated by a proton transport mechanism found in the membrane embedded  $F_0$  complex. Proton motive force (pmf) is the electrochemical gradient of protons utilized by  $F_0$  to drive synthesis of ATP at catalytic sites located in the  $F_1$  complex (Fig. 1; Duncan 2004). *In vitro*,  $F_1$  can be dissociated from the membrane as a water-soluble form that can only hydrolyze ATP.

The  $F_1$  complex is composed of 5 distinct subunits:  $\alpha_3\beta_3\gamma_1\delta_1\varepsilon_1$ . The  $F_0$  complex, located in the membrane lipid bilayer, is composed of 3 distinct subunits:  $a_1$ ,  $b_2$ , and  $c_{10}$ . Specifically, the  $a$ - and  $c$ -subunit play a role in catalyzing proton transport through proton binding at the  $c$ -subunit ring proton transport sites, facilitating rotation relative to the  $a$ - and  $b$ - subunits.



**Figure 1: Model of *E. coli* ATP synthase rotary motor enzyme:** Black arrows show the direction of proton transport and subunit rotation during ATP synthesis (Duncan 2014 review).

The  $\epsilon$ -subunit of  $F_1$  has been further studied for its role in inhibition of enzyme activity. Epsilon consists of two domains: the N-terminal domain ( $\epsilon$ NTD) and the C-terminal domain ( $\epsilon$ CTD) (Fig. 2). Studies have revealed that the  $\epsilon$ CTD is not required for oxidative phosphorylation but instead is involved in the regulation and inhibition of enzyme activity (Feniouk 2006). The  $\epsilon$ CTD has two known conformations: extended and compact. The extended conformation or inhibitory state of  $\epsilon$ CTD ( $\epsilon_x$  state) inserts into the central catalytic cavity of the  $F_1$  complex, interacting with 5 other subunits (Cingoloni and Duncan 2011). The  $\epsilon$ CTD in  $\epsilon_x$  state inhibits the enzyme by preventing the rotation of the  $\gamma$ -subunit. The active form of the enzyme has a compact conformation of the  $\epsilon$ CTD ( $\epsilon_c$  state) (Wilkins et al. 1998). When soluble  $F_1$  is diluted to low concentrations,  $\epsilon$  can dissociate from the enzyme, which releases the inhibition. The primary focus of my research project is to use site-directed mutagenesis to identify residues in the  $\epsilon$ CTD that are most important for inhibitory interactions of the  $\epsilon$ CTD with the  $\beta$ - and  $\gamma$ - subunits.



**Figure 2, Extended and compact conformations of epsilon.** Image **a** shows the  $F_1$  complex with  $\alpha$  subunits omitted to visualize  $\epsilon$ 's extended state (magenta), and the position of the  $\epsilon$ CTD in the compact state (gray helices). Image **b** shows  $\epsilon$  alone with the two helices in the extended conformation. Image **c** shows the compact conformation of  $\epsilon$ . Adapted from Cingolani and Duncan (2011).

A recent structure of  $F_1$  from *E. coli* has provided relevant insights on the catalytic sites and subunit interactions of the enzyme (Cingolani and Duncan 2011). In Tom Duncan's lab, previous research demonstrated that the  $\epsilon_x$  state is prevented from forming until hydrolysis of ATP at a catalytic site. This conformational change is significant in showing how  $\epsilon$ CTD inhibits  $F_1$  (Shah et al. 2013). Since  $\epsilon$ -inhibition is bacterial-specific, the  $\epsilon$ CTD could be a good target for antibiotic development by discovering compounds that mimic or strengthen inhibition on ATP synthase by the  $\epsilon$ CTD.

The aim of this research project is to understand the inhibitory behavior of the  $\epsilon$ CTD in the  $F_1$  complex of  $F_0F_1$ . Guided by a high-resolution structure of the  $\epsilon$ -inhibited catalytic complex (Cingolani and Duncan 2011), site-specific mutagenesis of the  $\epsilon$ CTD will be used to investigate interactions thought to be important for inhibition. Mutating residues of the enzyme can provide information on specific interactions necessary for ATP synthesis, hydrolysis, and inhibition. Using information available from the *E. coli*  $F_1$  structure, I will explore interactions between the  $\epsilon$ CTD and other  $F_1$  subunits of the enzyme that are likely to play a role in inhibition. The target sites for mutagenesis are in the C-terminal domain of the  $\epsilon$ -subunit.

Previous existing mutations include  $\epsilon\Delta 5$ , with the last five residues of the  $\epsilon$ -CTD deleted and  $\epsilon 88\text{stop}$ , with the whole inhibitory domain deleted (Shah 2015). These mutants will be useful for comparison with new mutants. Specific residues in two sites on the  $\epsilon$ CTD will be mutated to observe changes in functional and possibly disruptive interactions with other subunits. Three single



Mariam Bhatti  
Capstone

point mutations will be made at alanine97 ( $\epsilon$ A97) residue in helix 1. The second site is on helix 2, lysine123 ( $\epsilon$ K123). The primary focus is to use site-directed mutagenesis to identify residues in the  $\epsilon$ -CTD that are most important for inhibitory interactions of  $\epsilon$ -CTD with the  $\gamma$ - and  $\beta$ - subunits.

## Methods

### Construction of pMB $\epsilon$ 01 plasmid:

The vector for separate expression of  $\epsilon$ -subunit is modified version from pXH302s (Xiong and Vik 1995). An insert encoding an affinity tag (6xHis.Nsi) was introduced into the vector at the start of the *atpC* gene for  $\epsilon$ . The final plasmid pXH302s+NsiI.6xHis was named pMB $\epsilon$ 01. The NsiI site will be useful for moving the 6xHis-tagged  $\epsilon$  into other constructs. The method used to introduce the insert was fusion PCR (Ho et al 1989). Two initial PCR reactions (PCR 1 and PCR 2) amplified the upstream and downstream regions of the plasmid. The final fusion PCR reaction (PCR 3) created the desired insert. Then a ligation reaction (Wu and Wallace 1989) with vector and insert generated pMB $\epsilon$ 01. The unique NsiI site was used to screen plasmids from transformed colonies by restriction digest: a linearized plasmid showed successful insertion of the NsiI-containing fusion product. DNA sequence analysis (Upstate Medical University core Facility) confirmed that the plasmid contained the correct *atpC* gene with the added region including the NsiI site and encoding the 6xHis tag.

### SEQ:

```
 $\epsilon$ :      M   H   H   H   H   H   H   G   H   M>Epsilon
atpC:   ATG CAT CAC CAT CAT CAC CAC GGT CAT ATG
        (NsiI)                               (NdeI)
```

Structural identification of residues to mutate in the  $\epsilon$ CTD:

Using the molecular modeling program Chimera (Pettersen et al, 2004), I was able to analyze the epsilon-inhibited structure of the F<sub>1</sub> complex. Individual amino acids in the sequence were studied based on distances between individual atoms, angles/torsions, hydrogen bonds, and any clashes/contacts that may be important in the structure and function of the F<sub>1</sub> complex. I identified two residues, based on what is currently known about the molecular structure of ATP synthase, that appear to play important roles in  $\epsilon$ CTD:  $\epsilon$ A97 and  $\epsilon$ K123. “In silico” mutagenesis was used to introduce possible amino acid substitutions (rotomers) and study the contacts/clashes to determine the substitution that should be most effective at each residue.

Site-Directed Mutagenesis of the *atpC* gene for Epsilon:

Using mutagenic primers for  $\epsilon$ A97, the following mutants were made in pMB $\epsilon$ 01:  $\epsilon$ A97M,  $\epsilon$ A97Q,  $\epsilon$ A97L. Mutagenic primers were designed by first checking common codon usage in *E. coli*. The plasmid was linearized in a double-digest reaction with AfeI and PstI, and 5' and 3' PCR fragments were generated. Gibson Assembly (Gibson 2009) was used to combine the vector and insert for the  $\epsilon$ A97L mutant. A more traditional cloning method as described for pMB $\epsilon$ 01 construction was used for  $\epsilon$ A97Q,  $\epsilon$ A97M,  $\epsilon$ 88stop, and  $\epsilon$  $\Delta$ 5 mutants. Mutant plasmids were transformed into competent DH5 $\alpha$  cells and transformant plasmids were screened by DNA sequencing to confirm the presence of the desired mutation and the absence of any undesired changes. Outstanding mutagenesis projects are  $\epsilon$ L123M,  $\epsilon$ L123Q,  $\epsilon$ L123E.

Phenotypic Assay measuring Respiratory Growth:

Phenotypic effects of the mutations were observed after transforming each pMB $\epsilon$ 01 mutant into the XH1 expression strain (Xiong and Vik 1995). The XH1 expression strain has a deletion of the chromosomal epsilon gene but expresses all other subunits of the enzyme. This phenotypic assay measures growth of the mutants in liquid culture with succinate as a non-fermentable carbon source (Shah and Duncan 2015). *E. coli* is a facultative anaerobe, and can grow by aerobic respiration or by fermentation through glycolysis. Growth in liquid culture can provide a more sensitive ranking of the effects of the mutations on the *in vivo* assembly of functional ATP synthases. ATP synthesis function is required for growth on succinate. Thus, if cells do not grow, assembly of ATP synthase has been disturbed or enzymatic activity is compromised. For XH1 cells transformed with pMB $\epsilon$ 01 (wild-type or each  $\epsilon$  mutant), bacterial colonies were inoculated in 10 mL of Luria Bertani broth (LB) + ampicillin (0.1 mg/ml) overnight. Cells were then diluted again into 10 mL LB + ampicillin to an initial  $A_{600} = 0.1$ . Cultures were grown at 37 °C until  $A_{600} = 0.8$ , then cells were diluted 100-fold in minimal medium containing succinate as carbon source. Growth was measured every 15 minutes using a Biotek Synergy HT plate reader at 37 °C, with 0.4 mL triplicates of each strain in a 48-well microplate (Shah and Duncan 2015).

Preparation of inverted membrane vesicles for functional assays:

*E. coli* cells can be put under high pressure in a French Pressure cell and, as the pressure is slowly released, the pressure shift disrupts the cells (protocol in Shah and Duncan 2015). Cell membrane fragments turn inside out as a result, and the F<sub>1</sub> complex is exposed to the external

Mariam Bhatti  
Capstone

environment. Modified Lowry assay protocol was used to determine membrane protein concentration (Peterson 1977).

#### ATP Hydrolysis:

Assays to measure ATP hydrolysis by membranes were done by a photometric “coupled enzymes” assay (protocol in Shah and Duncan 2015). Assays testing the effects of lauryl dimethylamine-N-oxide (LDAO) and Dicyclohexylcarbodiimide (DCCD) were included as before (Shah and Duncan 2015). LDAO is known to alleviate epsilon inhibition, therefore activating the enzyme. DCCD covalently modifies aspartate61 residues in the c-ring of  $F_0$ , which are required for proton transport. Therefore, DCCD irreversibly inhibits ATP synthesis and hydrolysis by blocking proton transport through  $F_0$ .


**Table 1: Primers used in site-directed mutagenesis**

<b>Construct</b>	<b>Primers used in PCR: Sequence 5' to 3'</b>
pMBε01	<p><b>PCR reaction 1:</b> pMBε01 forward: C AAA CTG GAG ACT GTC <u>ATG CAT CAC CAT CAT CAC CAC</u> GGT GGC CAT ATG GCA ATG ACT TAC CAC XH downstream: GAC TGG CTT TTG TGC TTT TCA AGC CGG</p> <p><b>PCR reaction 2:</b> pMBε01 reverse: GTG GTA AGT CAT TGC CAT ATG GCC ACC <u>GTG GTG ATG ATG GTG ATG</u> <u>CAT GAC AGT CTC CAG TTT G</u> XH upstream: GAG CGT CGA TTT TTG TGA TGC TCG TC</p> <p><b>FPCR reaction 3:</b> pMBε01 forward and pMBε01 reverse</p>
εA97L	<p>εA97L forward: GAA GCG GCC ATG GAA <u>CTG</u> AAA CGT AAG GCT GAA GAG εA97L reverse: CTC TTC AGC CTT ACG TTT <u>CAG</u> TTC CAT GGC GCG CGC TTC</p>
εA97M	<p>εA97M forward: GAA GCG CGC GCC ATG GAA <u>ATG</u> AAA CGT AAG GCT GAA GAG εA97M reverse: CTC TTC AGC CTT ACG TTT <u>CAT</u> TTC CAT GGC GCG CGC TTC</p>
εA97Q	<p>εA97Q forward: GAA GCG CGC GCC ATG GAA <u>CAG</u> AAA CGT AAG GCT GAA GAG εA97Q reverse: CTC TTC AGC CTT ACG TTT <u>CTG</u> TTC CAT GGC GCG CGC TTC</p>
εΔ5	<p>εΔ5 forward: GCT GCG CGT TAT CGA GTT GTA ACA CCG GCT TGA AAA GCA C εΔ5 reverse: GTG CTT TTC AAG CCG GTG TTA CAA CTC GAT AAC GCG CAG C</p>
ε88stop	<p>ε88stop forward: GCC GAC ACC GCA ATT CGT GGC CAA TAA CAC CGG CTT GAA AAG C ε88stop reverse: GCT TTT CAA GCC GGT GTT ATT GGC CAC GAA TTG CGG TGT CGG C</p>


## Results

### Epsilon Mutants:

In order to examine the role of  $\epsilon$ -helix1/ $\gamma$  interaction for forming inhibited states, we targeted alanine97 ( $\epsilon$ A97) in the  $\epsilon$ CTD's first helix. Mutations were avoided that would likely disrupt the compact state. In  $\epsilon$ 's extended state,  $\epsilon$ A97 side chain contacts four residues of  $\gamma$  ( $\gamma$ P151,  $\gamma$ Q135,  $\gamma$ A134,  $\gamma$ G150). The structure of  $\epsilon$ -inhibited  $F_1$  indicates tight packing of the  $\epsilon$ A97 side chain in a pocket formed by these  $\gamma$  residues (Fig. 3). Mutations at this first target site should address whether the  $\gamma/\epsilon$ -helix 1 contact is important for transition into the  $\epsilon$ -inhibited state, and I will test this by introducing bulky residues that should disrupt this interface. Mutating  $\epsilon$ A97 to an amino acid with a larger side chain should disrupt this tight packing. Hydrophobic/nonpolar amino acid like leucine or methionine and a polar uncharged amino acid such as glutamine were considered as possible mutants. In the compact state,  $\epsilon$ A97 does not contact  $\epsilon$ 's second helix or the  $\epsilon$ NTD (Fig. 4). Mutating  $\epsilon$ A97 should not disrupt the compact conformation of  $\epsilon$ .



**Figure 3, In  $\epsilon$ 's extended conformation (magenta),  $\epsilon$ A97 (ball/stick) packs against several  $\gamma$  residues (yellow, space-filling). Van der Waal contacts of  $\epsilon$ A97 with  $\gamma$  are shown with green lines. The position of  $\epsilon$ A97 indicates tight packing with four residues of gamma. A larger side chain should disrupt the packing in this region.**




**Figure 4,  $\epsilon$ A97 compact conformation.** The side chain of  $\epsilon$ A97 has no contacts in the compact conformation, and faces away from the second helix. A mutation in this region should not disrupt the compact conformation of  $\epsilon$ .

Using Chimera, three potential mutants at  $\epsilon$ A97 were examined: leucine ( $\epsilon$ A97L), glutamine ( $\epsilon$ A97Q), and methionine ( $\epsilon$ A97M). In the extended state,  $\epsilon$ A97L and  $\epsilon$ A97Q have many predicted clashes. Leucine indicates 6-9 clashes, with 1 potential clash in the compact conformation. Glutamine indicates 8-15 clashes with residues on  $\gamma$ . Methionine is also a possible mutant, as it is not too hydrophobic and should not destabilize the  $\alpha$ -helix. Methionine in extended conformation indicates 8-14 clashes. There is a high probability of 0 clashes in the compact conformation for all three amino acid substitutions.

Another target site includes the second helix of  $\epsilon$  and its predominant role in inhibition. A few distinct residues of the  $\epsilon$ CTD appear to form specific hydrogen bonds that may be critical for stabilizing the  $\epsilon$ -inhibited state. Epsilon Lysine123 ( $\epsilon$ K123) in helix two contacts  $\beta$ D372 of subunit  $\beta_1$  and  $\alpha$ A405 of subunit  $\alpha_3$ . There is likely a hydrogen bond and a charge interaction between the side chains of  $\epsilon$ K123 and  $\beta$ D372 (Fig. 5). Lysine is positively charged, so introducing a long non-


Mariam Bhatti  
Capstone

polar amino acid like methionine or polar uncharged glutamine would disrupt this bond. Mutating lysine to a negatively charged amino acid like glutamate would disrupt this bond and introduce an electrostatic repulsion with  $\beta$ D372.



**Figure 5,  $\epsilon$ K123 in  $\epsilon$ 's extended conformation.** Ball and stick models of contacting residues are shown with green lines and apparent hydrogen bonds are shown with orange lines.

In  $\epsilon$ 's compact conformation, the first two carbons of the  $\epsilon$ K123 side chain contact Tyrosine53 and Proline47 in the  $\epsilon$ NTD (Fig. 6). Structural analysis indicates methionine, glutamine, and glutamate should not alter these interactions in the compact conformation. Additionally, these mutants should not disrupt the  $\alpha$ -helical structure of  $\epsilon$ .






**Figure 6, Contacts of  $\epsilon$ K123 in the compact conformation.** The  $C\beta$  and  $C\gamma$  carbons of the  $\epsilon$ K123 side chain have contacts (green lines) with Tyr53 and Pro47 in the  $\epsilon$ NTD. The most favorable rotamers of methionine, glutamine, and glutamate would have their  $C\beta$  and  $C\gamma$  carbons in similar positions, so these mutations should not alter these interactions in the compact conformation.

Using structure editing in Chimera, three potential mutants were modeled at  $\epsilon$ K123: methionine ( $\epsilon$ K123M), glutamine ( $\epsilon$ K123Q), and glutamate ( $\epsilon$ K123E). In the extended conformation, 1 likely rotamer of  $\epsilon$ K123M had a high probability to clash with  $\alpha$ A405- $C\beta$  of subunit  $\alpha_3$ . Another favorable  $\epsilon$ K123M rotamer is oriented away from this  $\alpha$ A405 and from  $\beta$ D372 of  $\beta_1$ , indicating the  $\epsilon$ K123M substitution may occur without significant steric disruptions. In  $\epsilon$ 's extended state, the  $\epsilon$ K123Q mutation indicates only one clash with  $\alpha$ A405 of subunit  $\alpha_3$ . The  $\epsilon$ K123Q mutation in  $\epsilon$ 's extended conformation indicates a high probability of having no clashes, but a small probability of having clashes with  $\alpha$ A405. In the compact conformation,  $\epsilon$ K123M indicates low probability of clashes with the  $\epsilon$ NTD, while  $\epsilon$ K123Q indicates a low probability of clashing with Tyrosine53 on the  $\epsilon$ NTD and Glutamine127 on  $\epsilon$ 's second helix. However, the compact conformation also indicates probability of forming clashes with Tyr 53 on  $\epsilon$ NTD and Glutamine 127 on  $\epsilon$ 's second  $\alpha$  helix. Compact confirmation also indicates probability of forming clashes with Tyr 53 on  $\epsilon$ NTD. Thus, the potential for mutagenesis of  $\epsilon$ K123 in  $\epsilon$ 's second  $\alpha$ -helix is promising. However, the  $\epsilon$ K123 mutants have not been created yet, so the remaining results will focus on the  $\epsilon$ A97 mutants.


#### Phenotypic Assay measuring Respiratory Growth

As shown in a recent study, aerobic growth is inhibited when five C-terminal residues are deleted from the  $\epsilon$  subunit of *E. coli*'s ATP synthase (Shah and Duncan 2015). This growth inhibition is due to a reduced capacity for ATP synthesis *in vivo*. Whereas the prior study over-

expressed the entire ATP synthase, our results (Figs. 7, 8) confirm a similar or slightly greater reduction in growth yield when  $\epsilon\Delta 5$  is expressed separately and all other  $F_0F_1$  subunits are expressed from the chromosomal *atp* operon. The new  $\epsilon A97$  mutants tested here are expected to disrupt the interaction of  $\epsilon$ -helix1 with  $\gamma$ . If the interaction of  $\epsilon$ -helix1 with  $\gamma$  is critical for epsilon to achieve the extended state, disrupting it will prevent inhibition of the enzyme. Aerobic growth assays of  $\epsilon A97$  mutants in liquid medium with succinate as the sole carbon source show that the  $\epsilon A97$  mutants grew almost as well as wild-type cells (Figs. 7, 8). The  $\epsilon A97$  mutants grew as well as or better than the  $\epsilon 88stop$  mutant, which has its entire C-terminal domain deleted.. Overall, these results indicate that  $F_0F_1$  is assembled properly in  $\epsilon A97$  mutants and retains near-normal ATP synthesis *in vivo*. Hence,  $\epsilon A97$  mutations do not lead to increased inhibition of the enzyme as seen with  $\epsilon\Delta 5$ .



**Fig 7, Phenotypic Respiratory Growth Curves.** Growth kinetics are shown in defined medium that contains succinate as sole carbon source. Cells were grown for at least 24 hours, with OD measurements taken every 15 minutes.




**Fig 8, Relative Growth Yield of Mutants as compared to wild-type:** Wild-type's growth plateau was set as 100% and values for the relative growth yields of mutants are shown. Error bars represent standard error of the mean (SEM) from 4 separate assays.

ATPase Activities of wild-type and mutant membranes:

ε97 mutants show higher activity than WT, this could be because mutants are inherently activated compared to WT, but might also be due to higher levels of F<sub>o</sub>F<sub>1</sub> expressed in the mutant membranes. LDAO is an amine oxide detergent that has been shown to increase ATPase activity by disrupting the epsilon inhibition of the enzyme. Therefore, it can indicate the extent of inhibition by ε. We subjected all membranes to coupled ATPase assays with and without LDAO to estimate the levels of ε-mediated inhibition. WT showed 1.9x activation by LDAO (p=0.002), which is consistent with previous studies. The εA97M and εA97Q mutants showed almost no activation by LDAO, suggesting those membranes lack intrinsic capacity for ε

inhibition.. The  $\epsilon$ A97L data were too scattered to conclude if it is more similar to WT or the other two  $\epsilon$ A97 mutants. LDAO activation data for  $\epsilon\Delta 5$  and  $\epsilon 88\text{stop}$  are scattered but similar to results with those mutants shown by Shah and Duncan, 2015 (2X for  $\epsilon\Delta 5$ , 1.4X for  $\epsilon 88\text{stop}$ ).

DCCD is an inhibitor of ATP synthase that covalently binds to the Aspartate 61 (Asp61) residue of the c-subunits. This residue is essential for proton transport through  $F_0$ . Thus, pre-incubation of membranes with DCCD for 30 min inhibits proton transport through  $F_0$  and, when  $F_1$  is well coupled to  $F_0$  in the membranes, inhibits the majority of membrane ATPase activity. We found that the ATPase activity of all mutant membranes is as sensitive to inhibition by DCCD as are WT membranes. Thus, the  $\epsilon$ A97 mutations do not cause any significant uncoupling between the  $F_1$  and  $F_0$  motors in the ATP synthase.



**Figure 9, *In-vitro* membrane ATPase activities.** Direct assays (clear bars) and LDAO activation (shaded bars) are shown. Below each sample, the number of separate assays is shown in

parentheses.(direct, +LDAO). Error bars represent the SEM. For each membrane sample, the value above the bars is the ratio of activity +LDAO to the direct activity.

## Discussion

My project focused on understanding the inter-subunit and intra-subunit interactions of the epsilon subunit in bacterial F-type ATP synthase. We generated three mutants that had single point mutations at  $\epsilon$ A97 with the help of site-directed mutagenesis. The mutants were subjected to phenotypic growth assays in medium containing a non-fermentable carbon source, succinate. Since non-fermentative growth requires ATP synthesis by  $F_0F_1$ , any defect in growth would be a result of impaired ATP synthase function. Our assays did not show any major defect in growth in the  $\epsilon$ A97 mutants. The growth yield was slightly reduced but was still better than  $\epsilon$ 88stop, which has the entire  $\epsilon$ CTD deleted.

ATP hydrolysis activity of the mutants was examined with the help of a coupled enzyme assay. In order to take a closer look at  $\epsilon$ -mediated inhibition, the samples were treated with LDAO, which relieves  $\epsilon$ -mediated inhibition, increasing the ATPase activity. The three mutants showed less activation than WT membranes. The reduced activation of  $\epsilon$ A97 mutants was similar to that seen with  $\epsilon$ 88stop, which suggests that  $\epsilon$ A97 mutations may disrupt the ability of  $\epsilon$  to achieve the inhibited state. In order to take a closer look at the effect of the  $\epsilon$ A97 mutants on the enzymatic activity, the inhibition of ATPase activity by  $\epsilon$ A97M/Q/L should be directly measured by quantifying  $F_0 F_1$  in membranes by immuno-blotting (as in Shah and Duncan, 2015).

Residue  $\epsilon$ A97 is present in helix1, and contacts the  $\gamma$  subunit in the  $\epsilon$ -inhibited state or  $\epsilon$ -helix2 in  $\epsilon$ 's compact conformation. Residue  $\epsilon$ K123 is present in  $\epsilon$ -helix2, which contacts  $\alpha$

Mariam Bhatti  
Capstone

and  $\beta$  subunits in the inhibitory state, but contacts  $\epsilon$ -helix1 in  $\epsilon$ 's compact state. The proton flow through  $F_0$  is coupled with enzymatic reactions by  $F_1$ . However, in reality, there is some amount of slippage observed even in WT enzyme whereby some protons 'slip' through without resulting in work. The mutations in the above mentioned locations might result in increased leakage of protons or decreased coupling. As a result, the mutant enzymes may have an impaired capacity to generate proton motive force. This is unlikely for two reasons: first, mutant cells showed nearly normal growth on the respiratory substrate succinate; second, DCCD modification of  $F_0$  inhibited mutant membrane ATPase as effectively as for WT, indicating normal  $F_0$ - $F_1$  coupling. To further confirm this, a fluorescent assay could be used to directly measure the kinetics of proton pumping by the mutant membranes (as in Shah and Duncan 2015).

In order to study the effects of the mutations on interaction of the  $\epsilon$ CTD with  $F_1$ , another approach should be used for future studies: to correlate inhibition with  $F_1$  binding and dissociation of  $\epsilon$ .  $F_1$  can be isolated from  $F_0F_1$ -ATPase in a soluble form and  $\epsilon$  becomes a dissociable inhibitor.  $F_1$  can be depleted of endogenous  $\epsilon$  by an anti- $\epsilon$  immuno-affinity column. Thus, inhibition of isolated  $F_1(-\epsilon)$  can be quantified by adding each isolated  $\epsilon$  mutant. As shown by Shah and Duncan (2013), affinity-tagged  $\epsilon$  (+/- mutants) can be overexpressed for this purpose. Thus, the inhibitory constant ( $K_I$ ) can be compared between wild type and mutant forms of  $\epsilon$ . The  $\epsilon$ A97 mutations may lead to altered binding of  $\epsilon$  to other  $F_1$  subunits. In wild-type enzyme,  $\epsilon$ 's C-terminal  $\alpha$ -helices alternate rapidly between the compact and extended conformations. In the compact conformation, the helices are in a coiled-coil conformation with each other; whereas in the extended conformation, helix1 establishes contacts with the  $\gamma$  subunit and helix2 establishes contacts with 2  $\alpha$ , 2  $\beta$  and  $\gamma$  subunits. With isolated WT  $F_1$ ,  $\epsilon$  favors the extended conformation and the enzyme is strongly inhibited. This conformational change may be

Mariam Bhatti  
Capstone

prevented if the  $\epsilon$ A97 mutations disrupt the  $\epsilon$ -helix1/ $\gamma$  interaction, and that interaction is necessary for  $\epsilon$ -helix2 to insert inside  $F_1$  to achieve the fully-inhibited state. To test this directly, an established optical method can be used to measure kinetics of binding and dissociation for  $F_1$  and mutant  $\epsilon$ . The extended state of the  $\epsilon$ CTD limits the  $F_1/\epsilon$  dissociation rate so, for example,  $F_1/\epsilon$ 88stop dissociates  $\sim 80X$  faster than  $F_1/wt-\epsilon$  (Shah and Duncan, 2013). Thus, measuring  $F_1/\epsilon$  with the  $\epsilon$ A97 mutants should confirm whether the mutations disrupt access to the extended, inhibitory conformation.

ATP synthesis assays will also be needed for key interpretation of the  $\epsilon$ A97 mutants. These assays can be easily done with the same membrane samples that were used for ATP hydrolysis assays. Other future experiments should include carrying out immunoblotting experiments to determine expression level of  $\epsilon$  in the strains and the content of  $F_0F_1$  in the membranes (anti- $\beta$  blot, as in Shah et al., 2015). A western blot will quantify activity per mg of enzyme rather than activity per mg of membrane. Additionally, outstanding projects include mutations at  $\epsilon$ K123. This site contains possible hydrogen bond and side chain specific interactions that may be important for  $\epsilon$  inhibition..

Overall, my work continues the lab's focus on understanding the complete role of the  $\epsilon$  subunit in the functioning of the bacterial ATP synthase. It is established that  $\epsilon$  is responsible for auto-inhibition of the enzyme. With ATP synthase approved as a target for drugs against pathogenic bacteria like *Mycobacterium tuberculosis*, targeting the auto-inhibitory properties of  $\epsilon$  subunit in pathogens may provide another way of inhibiting their growth and working around the problem of drug resistance. In order to do so, further studies need to be done to elucidate the unique role of the  $\epsilon$  subunit in regulating function of bacterial ATP synthases.

### **Acknowledgements**

First, I would like to thank my advisor, Thomas Duncan for accepting me into his lab my sophomore year. I would not have been able to complete my capstone without your guidance. Thank you for sacrificing weekends to do membrane preps and assay work with me, which was crucial to progress my capstone.

I want to thank my mentor, Nancy Walker who helped me with most of my cloning and mutagenesis work. Thank you for teaching me new strategies and teaching me to be patient with research.

I also want to acknowledge my post-doc Naman Shah for helping me during my initial years in the lab and always being willing to give me feedback and advice.

Thank you Renee Crown Honors program and faculty for choosing to fund my research this past summer and recognizing the potential in my research project.

Thank you Professor Chan, for being my capstone reader and teaching me many of the concepts that I needed to advance my research in your Biochemistry II course that I took my junior year.

I am grateful for all the support I have received from advisors, mentors, and peers during this process. Without the help of all my mentors and advisors at Syracuse University, including Kari Segraves and Kate Hanson, I would not have been able to complete my capstone and I greatly appreciate your help.

Finally, I want to thank my family and friends for their constant support and encouragement.



### Citations

- Bulygin, V.V., Duncan, T.M. & Cross, R.L. Rotor/Stator interactions of the  $\epsilon$  subunit in Escherichia coli ATP synthase and implications for enzyme regulation. *J. Biol. Chem.* 279, 35616-21 (2004).
- Cingolani, Gino, and Thomas M. Duncan. "Structure of the ATP Synthase Catalytic Complex (F1) from Escherichia Coli in an Autoinhibited Conformation." *Nat Struct Mol Biol Nature Structural & Molecular Biology* 18.6 (2011): 701-07. Web.
- Duncan, T.M. The ATP synthase: parts and properties of a rotary motor. in *The Enzymes*, vol. XXIII: Energy Coupling and Molecular Motors Vol. 23 (eds. Hackney, D.D. & Tamanoi, F.) 203–275 (Elsevier, New York, 2004).
- Gibson, Daniel. "One-step Enzymatic Assembly of DNA Molecules up to Several Hundred Kilobases in Size." *Protocol Exchange* (2009). Web.
- Ho, Steffan N., Henry D. Hunt, Robert M. Horton, Jeffrey K. Pullen, and Larry R. Pease. "Site-directed Mutagenesis by Overlap Extension Using the Polymerase Chain Reaction." *Gene* 77.1 (1989): 51-59. Web.
- Peterson, G. L. "A simplification of the protein assay method of Lowry et al. which is more generally applicable." *Anal. Biochem.* 83 (1977): 346-356.
- Richter, M.L. Gamma-epsilon interactions regulate the chloroplast ATP synthase. *Photosynth. Res.* 79, 319–329 (2004).
- Rodgers, A.J. & Wilce, M.C. Structure of the  $\gamma$ - $\epsilon$  complex of ATP synthase. *Nat. Struct. Biol.* 7, 1051-1054 (2000).
- Shah, Naman B, and Thomas M. Duncan. "Aerobic Growth of Escherichia Coli Is Reduced, and ATP Synthesis Is Selectively Inhibited When Five C-terminal Residues Are Deleted from the  $\epsilon$  Subunit of ATP Synthase." *Journal of Biological Chemistry J. Biol. Chem.* 290.34 (2015): 21032-1041. Web.
- Shah, N. B., M. L. Hutcheon, B. K. Haarer, and T. M. Duncan. "F1-atpase of Escherichia coli: the - Inhibited State Forms After ATP Hydrolysis, is Distinct from the ADP-inhibited state, and Responds Dynamically to Catalytic Site Ligands." *Journal of Biological Chemistry* 288.13 (2013): 9383-395. Web.
- Petersen EF, Goddard TD, Huang CC, Couch GS, Greenblatt DM, Meng EC, Ferrin TE. [UCSF Chimera--a visualization system for exploratory research and analysis.](#) *J Comput Chem.* 2004 Oct;25(13):1605-12.

Mariam Bhatti  
Capstone

Wu, Dan Y., and R. Bruce Wallace. "The Ligation Amplification Reaction (LAR)—Amplification of Specific DNA Sequences Using Sequential Rounds of Template-dependent Ligation." *Genomics* 4.4 (1989): 560-69. Web.

Xiong, H. and S. B. Vik. "Construction and plasmid-borne complementation of strains lacking the epsilon subunit of the Escherichia coli F1F0 ATP synthase." *Journal of Bacteriology* 177 (2013): 851-853.